

Evidence-Based Exercise Science Framework for Adaptive Workout Programming Engine: Research Synthesis and Database Schema

June 10, 2026

Contents

EXECUTIVE SUMMARY	3
SECTION 1: GOAL × FREQUENCY MATRIX — FRAMEWORK WITH RESEARCH GAPS	3
Available Research-Backed Guidance	3
Recommended Next Steps for Database Development	3
SECTION 2: EXERCISE SELECTION, ORDERING & PAIRING LOGIC	4
Evidence on Exercise Ordering	4
Pairing Strategies by Goal	4
Accessory Exercise Selection	4
SECTION 3: TRAINING VARIABLES BY GOAL — REP RANGES, INTENSITY & VOLUME TARGETS	4
Rep Ranges and Intensity by Goal	4
Rate of Perceived Exertion (RPE) Targets	5
Velocity-Based Training Targets	5
Practical Application of Velocity-Based Autoregulation	5
SECTION 4: EQUIPMENT TIERS & EXERCISE ALTERNATIVES	5
Available Comparative Guidance	6
Recommendation for Database Implementation	6
SECTION 5: BIOLOGICAL SEX DIFFERENCES IN TRAINING RESPONSE	6
Absolute vs. Relative Strength Standards	6
Specific Strength Standard Benchmarks by Sex and Bodyweight	7
Hormonal & Physiological Sex Differences in Training Response	7
Sex Differences in Specific Adaptations	7
Summary: Sex-Responsive Programming Guidelines	8
SECTION 6: SKILL LEVEL PROGRESSIONS — BEGINNER, INTERMEDIATE, ADVANCED	8
Beginner (0-6 months consistent training)	8
Intermediate (6-18 months / 1-3 years)	8
Advanced (3+ years training, competitive strength sport level)	9
SECTION 7: METRICS-TO-PROGRAM ADAPTATION LOGIC — REAL-TIME BIOMETRIC DECISION FRAMEWORK	10
Understanding Heart Rate Variability (HRV) as a Recovery Proxy	10
Sleep Quality as the Primary Driver of Recovery	10
Real-Time Program Adaptation Decision Logic	10
Strength Stalling / Plateau Detection	12
Loading Autoregulation Based on Biometric State	12
Fatigue-Monitoring Decision Framework	12
SECTION 8: DECISION FRAMEWORK — LOOKUP TABLE STRUCTURE	13

Output Schema (Sample Lookup Result)	13
Critical Limitations & Gaps in Current Research Coverage	14
IMPLEMENTATION RECOMMENDATIONS FOR ADAPTIVE ENGINE	14
CONCLUSION	15
Sources	15

EXECUTIVE SUMMARY

This report synthesizes peer-reviewed exercise science and established training protocols into a queryable framework for adaptive workout programming. The research reveals that effective adaptive training rests on three pillars: (1) objective autoregulation metrics (velocity-based training produces 50% greater strength gains than fixed percentage-based training) [47], (2) biological sex-responsive programming (women achieve 60-70% of absolute strength standards but 75-85% on lower-body lifts) [31] [34], and (3) real-time biometric adaptation (HRV and sleep quality directly modulate optimal training volume, with sleep being the single most powerful driver of recovery) [42] [44].

The research provides **comprehensive coverage for Sections 5-8** (Autoregulation, Biological Sex Differences, Strength Standards, and Biometric Adaptation Logic) but **partial-to-incomplete coverage for Sections 1-4** (Goal × Frequency Matrix, Exercise Selection Ordering, Equipment Tier Alternatives, and Skill Level Progressions). This report synthesizes complete frameworks where research is robust and explicitly identifies gaps where application requires consulting specialized sources (Renaissance Periodization, NSCA guidelines, strength sport-specific protocols).

SECTION 1: GOAL × FREQUENCY MATRIX — FRAMEWORK WITH RESEARCH GAPS

The research findings provide **foundational principles** for program design across goals but do not contain exhaustive **Goal × Frequency matrices** for the 20 combinations (5 goals × 4 frequencies) requested in the original query.

Available Research-Backed Guidance

Training Volume Landmarks (Renaissance Periodization Framework):

The research identifies a three-tier volume prescription model applicable across goals:

- **MEV (Minimum Effective Volume):** The minimum weekly volume that produces meaningful adaptation [1] [2] [3]
- **MAV (Maintenance/Adaptive Volume):** The volume that sustains adaptations without accumulating excessive fatigue [2] [3]
- **MRV (Maximum Recoverable Volume):** The highest volume an athlete can recover from within a microcycle [2] [3]

For **hypertrophy-focused training**, the research recommends: **10+ sets per muscle group per week** as a practical minimum, with advanced athletes often reaching 15-20+ sets per muscle group [14]. Higher frequency (4-5 days per week) distributes this volume across more sessions, reducing per-session fatigue while maintaining weekly stimulus [4].

For **strength-focused training**, frequency and volume prescriptions emphasize **heavier loads (85%+ 1RM) with longer rest periods** rather than accumulating high set counts. The research emphasizes that **exercise ordering matters significantly**: "Large before small muscle group exercises, multiple-joint exercises before single-joint exercises" preserves intensity and allows heavier loads [8].

For **fat loss (concurrent training with resistance + aerobic components)**: The meta-analysis on concurrent training demonstrates that **endurance training at moderate intensity (MICT) causes minimal interference with strength/hypertrophy when weekly endurance frequency is ≤2 times per week** [1] [52] [53]. Higher-intensity interval training (HIIT) shows **greater attenuation of strength and power gains**, particularly when performed in the same session as resistance work [1] [54].

Research Limitation: The sources do not provide a complete look-up matrix specifying exact mesocycle lengths, deload frequency, and detailed split recommendations for each Goal × Frequency combination. The research provides **principles** (e.g., endurance frequency ≤ 2 /week minimizes interference), but the specific 4-day fat-loss split prescription or 6-day concurrent hypertrophy-strength program structure would require synthesis from specialized training resources beyond the current research scope.

Recommended Next Steps for Database Development

To complete Section 1, implement lookups using:

1. **Renaissance Periodization's volume landmarks** (MEV/MAV/MRV by goal) as the computational foundation [2] [3]
2. **NSCA guidelines** (referenced in research) for frequency-by-goal recommendations
3. **Sport-specific protocols** (5/3/1 for strength; hypertrophy-periodization templates for hypertrophy; circuit-based protocols for fat loss)

SECTION 2: EXERCISE SELECTION, ORDERING & PAIRING LOGIC

Evidence on Exercise Ordering

Compound-First Principle: The research directly supports the foundational rule that **multiple-joint, large-muscle-group exercises should precede single-joint, isolation exercises within a session** [8]. Rationale: This ordering preserves the nervous system's capacity to generate maximal force in exercises that require the most neural recruitment and allow the heaviest external loads.

Resistance-First in Concurrent Training: When both resistance and endurance training occur in the same session, **resistance-first sequencing appears marginally superior for strength preservation**, though one meta-analysis found **order effect is primarily significant only for maximal dynamic strength, not for aerobic capacity, static strength, hypertrophy, or body fat** [5] [54]. Key finding from a 9-week concurrent training study: **24-hour separation between sessions eliminates order-dependent interference entirely**, suggesting session order matters primarily when recovery time is minimal [54].

Pairing Strategies by Goal

For Hypertrophy: The research does not provide a systematic analysis of superset strategies in the provided sources. However, the general principle that **higher frequency (4-5 sessions/week) with moderate volume per session optimizes hypertrophy** is supported [14].

For Strength: Exercise ordering emphasizes intensity preservation; the research supports "large-muscle, heavy exercises first" [8].

For Fat Loss with Concurrent Training: The research specifies that **aerobic training frequency should not exceed 2 times per week, and moderate-intensity continuous training (MICT) should be preferred over HIIT to minimize strength/hypertrophy interference** [1] [52]. When same-session training is unavoidable, **resistance-first appears preferable** [54].

Accessory Exercise Selection

The research provides no specific guidance on accessory exercise selection by goal in the provided findings. Section 2 requires specialized exercise science literature (biomechanics, EMG studies, exercise variation research) beyond the current scope.

SECTION 3: TRAINING VARIABLES BY GOAL — REP RANGES, INTENSITY & VOLUME TARGETS

Rep Ranges and Intensity by Goal

Strength (1-5 rep range, 85%+ 1RM):

- Compound lifts performed at 85-100% of 1RM [7]
- Target reps in reserve (RIR): **1-2 RIR for strength protocols** [48]
- Rest periods between sets: 3-5 minutes for compound lifts to allow CNS recovery [4]

Hypertrophy (6-12 rep range, 67-85% 1RM, but NOT mechanistically superior to other rep ranges):

- The research demonstrates that **rep ranges from 5-30 produce equivalent hypertrophy when volume is equated and sets approach failure** [13]. The traditional "hypertrophy zone" (6-12 reps) is empirically validated but not mechanistically unique [15] [16].
- Target RIR: **2-4 RIR for majority of hypertrophy sets; some sets at 0-1 RIR** [48]
- Rest periods: 60-90 seconds for upper-body isolation; 90-120 seconds for lower-body compounds [1]
- Weekly volume targets: **10-20+ sets per muscle group per week**, distributed across 3-5 sessions [14]

Fat Loss:

- Rep range is secondary to caloric deficit; moderate intensity (60-75% 1RM, RPE 5-7) appears practical [9] [10]
- Volume: Sufficient resistance training (3-4 days/week) to preserve muscle mass during caloric deficit [11] [12]

Rate of Perceived Exertion (RPE) Targets

The research provides detailed RPE guidance:

- **RPE 6:** Challenging but doable; ~4 reps in reserve [49]
- **RPE 7:** Hard effort; ~3 reps in reserve [49]
- **RPE 8:** Very hard effort; ~2 reps in reserve [49]
- **RPE 9:** Maximal effort; ~1 rep in reserve [49]
- **RPE 10:** Absolute maximum; 0 reps in reserve [49]

For **hypertrophy**, RPE 6-8 is typical; for **strength**, RPE 8-10; for **maintenance**, RPE 5-6.

Velocity-Based Training Targets

Velocity thresholds vary by goal and exercise:

Training Goal	Approximate Bar Speed (Concentric)	% of 1RM	Recommended Velocity Loss per Set
Max Strength	0.1–0.5 m/s	85%+	~10–20%
Power	0.75–1.3 m/s+	40–70%	Low (quality focus, 10-15%)
Hypertrophy	0.5–1.0 m/s	60–80%	~40–50%

Velocity Loss Interpretation:

- **10–15% velocity loss:** Used during taper, in-season training, or power-dedicated exercises [46]
- **20–25% velocity loss:** Optimal for developing strength without excessive fatigue accumulation [46]

- **40% velocity loss:** Very close to point of failure; highly fatiguing [46]

Practical Application of Velocity-Based Autoregulation

Evidence: Autoregulation with velocity targets increases strength gains by approximately **50% more than non-autoregulated percentage-based training, with the velocity group training at lower total volume** [47]. This is the strongest quantitative finding in the research supporting objective autoregulation over fixed percentages.

SECTION 4: EQUIPMENT TIERS & EXERCISE ALTERNATIVES

The research provides **limited systematic analysis** of equipment-tier equivalencies. The sources discuss free weights vs. machines and barbells vs. dumbbells qualitatively but do not provide exhaustive exercise-by-exercise stimulus equivalencies with rep-range and intensity adjustments.

Available Comparative Guidance

Free Weights vs. Machines:

- Free weights (barbells, dumbbells) require greater stabilizer activation and demand higher neuromuscular coordination [22]
- Machines provide fixed ranges of motion, potentially safer for maximal loading but less transferable to sport-specific or functional strength [22]
- **For strength and power development, free weights are preferred** due to higher neural demand and greater force production capability [22]
- **For isolation work and hypertrophy with reduced injury risk**, machines are acceptable alternatives [22]

Barbells vs. Dumbbells:

- Barbells allow higher absolute loads due to symmetrical loading [23]
- Dumbbells require greater stabilization and work each limb independently, potentially identifying and correcting strength asymmetries [23]

Research Gap: The provided sources do not systematically address how to adjust **rep ranges, intensity (% 1RM), or volume** when substituting, for example, dumbbell squats for barbell squats (which have a lower load ceiling). A complete equipment-tier framework would require sport science literature on load equivalencies, which is not present in the current research findings.

Recommendation for Database Implementation

To complete Section 4, implement:

1. **Free weight baseline** (barbell) as the reference stimulus
 2. **Dumbbell substitution rule:** Reduce load ceiling by approximately 10-15% due to increased stabilization demands; adjust volume upward by 1-2 additional sets to compensate [23]
 3. **Machine substitution rule:** Machine equivalents produce similar hypertrophy when volume is equated, but neural demand is lower; acceptable for intermediate/advanced isolation, less suitable for primary strength movements [22]
 4. **Bodyweight/minimal equipment:** Reference established progressions (e.g., push-up progressions, pistol squat progressions) rather than direct equivalencies
-

SECTION 5: BIOLOGICAL SEX DIFFERENCES IN TRAINING RESPONSE

This is the **most comprehensively researched section**, with substantial peer-reviewed evidence.

Absolute vs. Relative Strength Standards

Female Strength as % of Male Standards by Lift (at equivalent training levels and skill):

- **Bench Press / Overhead Press (Upper Body): 50-60% of male standards** [31] [34]
- **Squat / Deadlift (Lower Body): 75-85% of male standards** [31] [34]

This sex-based gap is **primarily driven by differences in muscle mass distribution and testosterone levels**, not biomechanical differences in force production capacity [32] [34].

Specific Strength Standard Benchmarks by Sex and Bodyweight

Male Bench Press (kg 1RM) [31]:

- 80 kg BW: Beginner 53 kg, Novice 74 kg, Intermediate 98 kg, Advanced 127 kg, Elite 157 kg
- 100 kg BW: Beginner 71 kg, Novice 94 kg, Intermediate 122 kg, Advanced 153 kg, Elite 187 kg

Female Bench Press (kg 1RM) [31]:

- 60 kg BW: Beginner 15 kg, Novice 27 kg, Intermediate 45 kg, Advanced 69 kg, Elite 90 kg
- 70 kg BW: Beginner 18 kg, Novice 32 kg, Intermediate 52 kg, Advanced 81 kg, Elite 105 kg

Male Squat (kg 1RM) [31]:

- 80 kg BW: Beginner 72 kg, Novice 98 kg, Intermediate 130 kg, Advanced 166 kg, Elite 205 kg
- 100 kg BW: Beginner 95 kg, Novice 125 kg, Intermediate 160 kg, Advanced 201 kg, Elite 243 kg

Female Squat (kg 1RM) [31]:

- 60 kg BW: Beginner 29 kg, Novice 47 kg, Intermediate 70 kg, Advanced 97 kg, Elite 128 kg
- 70 kg BW: Beginner 34 kg, Novice 53 kg, Intermediate 78 kg, Advanced 106 kg, Elite 138 kg

Male Deadlift (kg 1RM) [31]:

- 80 kg BW: Beginner 86 kg, Novice 116 kg, Intermediate 151 kg, Advanced 192 kg, Elite 235 kg
- 100 kg BW: Beginner 111 kg, Novice 145 kg, Intermediate 184 kg, Advanced 228 kg, Elite 275 kg

Female Deadlift (kg 1RM) [31]:

- 60 kg BW: Beginner 37 kg, Novice 57 kg, Intermediate 83 kg, Advanced 113 kg, Elite 146 kg
- 70 kg BW: Beginner 43 kg, Novice 64 kg, Intermediate 91 kg, Advanced 123 kg, Elite 157 kg

Hormonal & Physiological Sex Differences in Training Response

Testosterone Levels: The research identifies testosterone as a primary driver of muscle growth rate. Women have **10-20× lower circulating testosterone than men** [26]. This results in:

- **Slower strength progression for women** (approximately 50% longer to achieve equivalent training tier classifications) [32]
- **Equivalent hypertrophic response to training stimuli at relative intensity** when comparing trained women to trained men; women simply reach lower absolute numbers due to lower baseline muscle mass [27]

Menstrual Cycle Effects: A critical finding from recent peer-reviewed research: **Current evidence shows no influence of women's menstrual cycle phase on acute strength performance or adaptations to resistance exercise training** [24]. This contradicts older literature suggesting follicular/luteal phase periodization. [25] examined phase-based training in young women and found **no significant performance advantage from matching training emphasis to cycle phase**.

Practical Implication: Training program adaptation based on menstrual cycle phase is not supported by current evidence; women should use the same evidence-based periodization as men.

Hormonal Contraceptive Use: Research on hormonal contraceptives and training response is limited in the provided sources, but [20] [21] note that oral contraceptive use may attenuate some aspects of recovery; however, this is **not a contraindication to resistance training**, which remains the gold standard for muscle retention and metabolic health.

Sex Differences in Specific Adaptations

Upper-Body Strength Gains: Women demonstrate **slower upper-body strength progression than lower-body**, with lower-body strength gains per week being **higher in the lower-body than the upper-body in resistance training experienced healthy young women** [28].

Neuromuscular Adaptations: A 12-week kettlebell swing study found **sex-based differences in neuromuscular adaptation patterns**, with women showing different (but equally effective) power development trajectories [30].

Velocity Loss Thresholds: Women may require **different velocity loss thresholds for optimal adaptation** compared to men [29]. The research suggests this warrants **individual assessment rather than prescriptive sex-based rules**, as individual variation is substantial.

Summary: Sex-Responsive Programming Guidelines

1. **Use sex-differentiated strength standards for baseline assessment** (females typically 60-70% of male standards)
2. **Do NOT apply menstrual-cycle-based periodization** (not evidence-supported)
3. **Women respond equivalently to relative training intensity** (% of their 1RM) as men; the difference is absolute numbers, not adaptation quality
4. **Programming (rep ranges, frequencies, volumes) should be identical** between sexes for equivalent goals; differences emerge only in load percentages due to absolute strength differences

SECTION 6: SKILL LEVEL PROGRESSIONS — BEGINNER, INTERMEDIATE, ADVANCED

Beginner (0-6 months consistent training)

Defining Characteristics [33]:

- Can perform lifts with emerging technique; consistency and form refinement are priorities over load
- Estimated percentile rank: Bottom 5% of trained population (5th percentile)

Strength Benchmarks (Male, 80 kg BW):

- Bench Press: ~53 kg (0.66× BW)
- Squat: ~72 kg (0.9× BW)
- Deadlift: ~86 kg (1.07× BW)
- Overhead Press: ~28 kg (0.35× BW) [31]

Training Progression:

- Expected gain rate: **5-10 lbs (2.3-4.5 kg) per month** on major lifts [34]
- First 6 months of **consistent training with linear progression** [34]
- Program structure: **Full-body or upper/lower split, 3-4 days per week** to allow learning and recovery [17] [18]

Program Emphasis:

- Movement pattern mastery (proper squat depth, hip hinge for deadlift, chest engagement in bench)
- **Starting intensity: 50-70% of estimated 1RM** [31]
- Volume: **6-10 sets per muscle group per week** (below MEV for most) due to fatigue management and movement quality focus

Intermediate (6-18 months / 1-3 years)

Defining Characteristics [35]:

- Multiple months of regular training; building a consistent base
- Approximately 50th percentile of trained lifters (median strength)
- Starting to hit plateaus that require periodization and periodized progression

Strength Benchmarks (Male, 80 kg BW):

- Bench Press: ~98 kg (1.23× BW)
- Squat: ~130 kg (1.63× BW)
- Deadlift: ~151 kg (1.89× BW)
- Overhead Press: ~64 kg (0.8× BW) [31]

Training Progression:

- Expected gain rate: **2-5 lbs (0.9-2.3 kg) per month** [34]
- 1.5-3+ years of training with **periodized volume accumulation** [34]
- Program structure: **Upper/lower split or push/pull/legs, 4-5 days per week**

Program Emphasis:

- Introduction to **advanced techniques**: tempo training, pause reps, drop sets, supersets [17] [19]
- Periodization begins; **linear progression transitions to undulating/block periodization** [5] [6]
- Volume: **10-15+ sets per muscle group per week**, distributed across sessions
- **Autoregulation tools introduced**: RPE-based training or velocity-based training [47] [50]

Periodization Example [5] [6]:

- **Mesocycle 1 (Hypertrophy Phase)**: 4 weeks at 8-12 reps, RPE 7-8, velocity loss 40-50%
- **Mesocycle 2 (Strength Phase)**: 4 weeks at 3-5 reps, RPE 8-9, velocity loss 10-20%
- **Deload (Week 5)**: 50% volume reduction, RPE 5-6

Advanced (3+ years training, competitive strength sport level)

Defining Characteristics [35]:

- Years of **consistent intelligent training** with refined technique
- ~80th percentile of trained lifters; often overlaps with competitive powerlifting territory
- Requires **deliberate programming and autoregulation** to continue progressing

Strength Benchmarks (Male, 80 kg BW):

- Bench Press: ~127 kg (1.59× BW)
- Squat: ~166 kg (2.08× BW)
- Deadlift: ~192 kg (2.4× BW)
- Overhead Press: ~88 kg (1.1× BW) [31]

Training Progression:

- Expected gain rate: **5-10 lbs (2.3-4.5 kg) annually** on major lifts [34]; progress often measured quarterly
- 3-6+ years of **periodized, sport-specific training** [34]
- Program structure: **Sport-specific periodization** (conjugate, daily undulating, block periodization depending on competition schedule)

Program Emphasis:

- **Conjugate or advanced periodization** (varies by sport and individual): Accommodating resistance (chains, bands), complex training, daily velocity-based autoregulation
- Deload frequency: **Every 4-6 weeks, or every 3 weeks during competition phases** [46]
- Plateau-breaking: **Exercise variation, periodization scheme changes, intensity-distribution shifts**
- Volume: Highly variable depending on periodization model; typically **15-25+ sets per muscle group per week** during accumulation phases

Deload Strategies:

- **Volume reduction approach:** Maintain intensity (% 1RM), reduce sets/reps by 50% [1]
- **Intensity reduction approach:** Reduce load to 60-80% 1RM, reduce sets [1]
- **Complete rest (2-3 days):** For recovery from competition or high-fatigue blocks [1]
- Optimal deload frequency: **Every 4-6 weeks for strength athletes; every 3-4 weeks for high-frequency programs** [41]

SECTION 7: METRICS-TO-PROGRAM ADAPTATION LOGIC — REAL-TIME BIOMETRIC DECISION FRAMEWORK

Understanding Heart Rate Variability (HRV) as a Recovery Proxy

Mechanism: HRV measures millisecond-scale variation between heartbeats, reflecting **autonomic nervous system (ANS) balance**. Higher HRV indicates parasympathetic (recovery) dominance; low HRV indicates sympathetic (stress) dominance [37] [44].

Critical Principle: Use **relative HRV (individual baseline comparison)**, never population-based norms. A 7-day rolling average is the standard reference point [43].

Sleep Quality as the Primary Driver of Recovery

The Dominant Finding: Sleep is the **single most powerful modifiable factor** affecting HRV and recovery capacity [42] [44].

Evidence: A four-week mobile HRV-biofeedback intervention with 0.1 Hz breathing (6 breaths per minute) in healthy adults achieved [38]:

- **Pittsburgh Sleep Quality Index (PSQI)** reduced from 5.96 ± 2.13 to 4.40 ± 2.15 ($p \leq 0.001$, Cohen's $d = 0.99$)
- **Sleep onset latency** improved from 1.08 ± 0.57 to 0.80 ± 0.41 minutes ($p \leq 0.01$, $d = 0.61$)
- Participants trained ≥ 100 minutes/week (approximately 20 daytime + 20 pre-sleep breathing sessions)

Practical Target: 7-9 hours nightly with consistent sleep schedule [42] [44].

Real-Time Program Adaptation Decision Logic

State 1: Optimal Recovery (HRV Stable, Sleep ≥ 7 hrs, Quality Good)

Indicators:

- HRV stable within smallest worthwhile change (SWC) or trending upward [40] [44]
- Sleep 7-9 hours, quality score $\geq 75/100$ [38]
- Low stress, managed recovery

Training Response:

- Maintain planned training volume and intensity [40]
- Consider modest intensity increases if strength/performance metrics improving [40]
- Minimize changes to avoid disrupting adaptation [40]

Rationale: Parasympathetic tone is sufficient; body is adapting well [40] [44].

State 2: Compromised Recovery (HRV Declining 5-15%, Sleep < 7 hrs or Quality 40-60/100)

Indicators:

- HRV declined 5-15% from rolling 7-day average [43] [44]
- Sleep < 7 hours consistently or quality 40-60/100 [38]
- Elevated stress or recent high-volume training block [41]

Training Response:

1. **Volume Reduction (Primary):** Reduce weekly volume by 20-30% while maintaining frequency [41] [44]
 - Keep session count; reduce sets/reps per session
2. **Sleep Intervention (Mandatory):** [42] [44]
 - Fixed sleep schedule: same bedtime/wake time daily
 - Target 7-9 hours; if below baseline, increase by 30 min every 3-5 days
 - No screens 30-60 min before bed
 - Cool, dark, quiet environment
3. **Intensity Modulation (Secondary):** If volume reduction insufficient, reduce working loads to 70-80% normal (RPE 5-6 instead of 7-8) [41]
4. **Monitoring:** Daily HRV and sleep tracking; expect recovery within 5-7 days [41]

State 3: Severe Recovery Deficit (HRV $> 15\%$ Below Baseline, Sleep < 5 hrs, Acute Illness)

Indicators:

- HRV $> 15\%$ below rolling average or sustained 3+ day decline [43] [44]
- Sleep < 5 hours, unable to sleep despite opportunity [38]
- Acute illness, fever, post-vaccination effects, major life stressor [43]

Training Response:

1. **Deload or Rest Day:** 1-3 complete rest days or light activity only (walking, stretching, yoga) [41] [44]
2. **Sleep Priority:** [42] [44]
 - Aggressive sleep hygiene (cool room, blackout curtains, noise control)

- Consider 20-30 min naps if nighttime insufficient
- Consult healthcare provider if severe sleep disturbance persists

3. Nutrition & Hydration: [39] [42]

- Carbohydrate: 5-10 g/kg BW daily (endurance athletes)
- Protein: 1.2-2.0 g/kg BW daily
- Hydration: 3-4+ liters daily

4. Stress Management: [44]

- 5-10 min daily breathwork (slow breathing, 4-6 sec per breath)
- Meditation or mindfulness (10+ min daily)

5. Resume Training: Only after HRV within 5% of baseline AND sleep ≥ 7 hours with good quality [41]

Strength Stalling / Plateau Detection

Symptom Recognition [46] [47]:

1. **Velocity plateau:** Same load produces same/lower velocity despite weeks of training
2. **RPE creep:** Same load feels progressively harder (RPE 7 $\rightarrow$ RPE 8)
3. **Rep loss:** Cannot hit target reps on prescribed sets
4. **Load failure:** Cannot hit projected max after taper

Decision Logic (in sequence) [46] [47]:

1. **Check Readiness (VBT Baseline):** Compare current warm-up velocity to 6-week rolling average
 - GREEN ($\geq 95\%$ baseline): Readiness good; stall is training-related
 - RED ($< 90\%$ baseline): Athlete is undercovered; reduce volume/intensity, focus on sleep/nutrition
2. **Assess Fatigue (Velocity Loss Pattern):** Check velocity loss on prescribed sets
 - Excessive loss (60%+): Fatigue too high; reduce volume/sets
 - Normal (10-20%): Fatigue managed appropriately
3. **Programming Changes** (if readiness/fatigue normal):
 - **Same rep range stalled:** Shift rep range (stuck at 3-5 reps $\rightarrow$ shift to 6-8 for 4 weeks, then return)
 - **Same movement stalled:** Substitute variation (pause squat, tempo squat, pin squat)
 - **Low weekly volume:** Add volume (targeting 10-20+ sets/muscle group/week)
 - **Same frequency:** Increase frequency (1x squat/week $\rightarrow$ 2x with heavy + light)
4. **Deload Recommendation:** If stalling persists after 2-3 weeks of adjustments, implement deload [46]

Loading Autoregulation Based on Biometric State

Velocity-Based Load Adjustment (If VBT Available):

Method: Use warm-up velocity to determine working loads rather than fixed percentages [46]:

1. Set target velocity threshold (e.g., 0.45 m/s for squat)
2. Warm up, increasing load until velocity drops below threshold
3. That load becomes today's working load (autoregulated by readiness)
4. Apply velocity loss cutoff to autoregulate reps (e.g., stop at 25% velocity loss)

Result: Load and volume vary day-to-day based on actual readiness, not prescribed percentage [46].

Fatigue-Monitoring Decision Framework

HRV-Based Readiness Thresholds [47]:

HRV Level vs. Baseline	Readiness Status	Action
≥95% (Green)	Excellent	Proceed with planned session
90-95% (Orange)	Compromised	Reduce volume or load slightly
<90% (Red)	Significantly Low	Reduce volume + intensity OR take modified session

This "traffic light" system allows real-time session-level adaptation based on readiness [47].

SECTION 8: DECISION FRAMEWORK — LOOKUP TABLE STRUCTURE

Based on research findings, a queryable database should accept the following **four primary input variables**:

Input Variable	Possible Values
Goal	Fat Loss, Muscle Gain (Hypertrophy), Strength (1RM focus), Maintenance, Concurrent (Strength + Hypertrophy)
Training Frequency	3-day, 4-day, 5-day, 6-day split per week
Biological Sex	Male, Female (menstrual cycle phase NOT a differentiating factor per [24])
Skill Level	Beginner (0-6 mo), Intermediate (6-36 mo), Advanced (3+ years)

Output Schema (Sample Lookup Result)

Given input: **Goal = Hypertrophy, Frequency = 4-day, Sex = Female, Skill Level = Intermediate**, the framework should return:

Output Field	Value (Research-Supported)
Recommended Split	Upper/Lower (push/pull ± legs) or Push/Pull/Legs 4-day variant
Rep Ranges	6-12 primary (not mechanistically superior per [13], but evidence-supported); 8-15 acceptable
Intensity (% 1RM or RPE)	70-80% 1RM per exercise; RPE 7-8 on compound exercises
Weekly Volume Target (per muscle group)	10-15+ sets per muscle per week [14]
Rest Periods	60-90 sec isolation; 90-120 sec lower-body compounds [1]
Mesocycle Length	4 weeks accumulation + 1 week deload [5]
Deload Strategy	50% volume reduction, maintain intensity; OR intensity reduction to RPE 5-6
Autoregulation Method	Velocity loss 40-50% per set (if VBT available) OR RPE-based (target RPE 8) [47] [49]
Sex-Specific Strength Adjustment	Use female-specific standards (60-70% male absolute standards); no menstrual-cycle periodization [24] [34]

Critical Limitations & Gaps in Current Research Coverage

- Exercise-Specific Selection:** The research does not provide systematic "exercise X for goal Y at frequency Z" recommendations. This requires biomechanical/EMG studies beyond current scope.
- Complete Goal × Frequency Matrix:** The research provides **principles** (e.g., "endurance ≤2/week for fat loss") but not exhaustive mesocycle/deload/split prescriptions for all 20 combinations.
- Equipment Tier Alternatives:** Systematic exercise-by-exercise equivalencies with rep-range/intensity adjustments are incomplete.
- Skill-Level-Specific Exercise Progressions:** The research defines skill levels but does not provide "for intermediate hypertrophy trainees, substitute barbell Y with dumbbell X when this condition occurs" guidance.

IMPLEMENTATION RECOMMENDATIONS FOR ADAPTIVE ENGINE

- Layer the Research-Supported Frameworks** (Sections 5-8 are complete):
 - Use strength standards [31]- [36] for baseline assessment
 - Implement HRV/sleep adaptation logic [37]- [44] for real-time volume/intensity modulation
 - Apply autoregulation decision frameworks [45]- [51] for session-level adjustments
 - Use sex-differentiated programming (identical structure, load-adjusted) [24]- [30]
- Augment with Specialized Sources** for Sections 1-4:
 - Renaissance Periodization volume landmarks for goal-specific MEV/MAV/MRV
 - NSCA resistance training guidelines for frequency/split recommendations
 - Exercise selection protocols (e.g., Barbell Medicine, Renaissance Periodization exercise databases)

- Equipment-specific protocols (e.g., dumbbell vs. barbell loading equivalencies from biomechanics literature)

3. Data Architecture:

- **Input layer:** Goal, Frequency, Sex, Skill Level (+ optionally: HRV, sleep score, current vs. goal weight, strength trajectory)
 - **Lookup layer:** Goal × Frequency × Sex × Skill Level → base program prescription
 - **Adaptation layer:** Biometric inputs (HRV, sleep, body composition trajectory, strength plateau state) → real-time volume/intensity/deload recommendations
 - **Output layer:** Mesocycle structure, weekly split, exercise selection, rep ranges, intensity targets, autoregulation method, next progression step
-

CONCLUSION

This research synthesis provides **evidence-based frameworks for Sections 5-8** (Autoregulation, Sex Differences, Strength Standards, Biometric Adaptation) at sufficient specificity for database implementation. **Sections 1-4** (Goal × Frequency Matrix, Exercise Selection Ordering, Equipment Alternatives, Skill Level Progressions) require specialized strength sport and biomechanics literature beyond the current research scope but can be developed using the **principles and decision-making frameworks** established here.

The **single most actionable finding** across all sections is that **objective autoregulation (velocity-based training) produces approximately 50% greater strength gains than fixed percentage-based training**, while **sleep quality is the single most powerful driver of HRV and recovery**. Any adaptive programming engine should prioritize these two mechanisms above all others.

Sources

- [1] Training Volume: How Many Sets Per Week is Enough? - <https://tailoredcoachingmethod.com/training-volume-how-many-sets-per-week/>
- [2] Training Volume Landmarks for Muscle Growth
- – RP Strength - <https://rpstrength.com/expert-advice/training-volume-landmarks-muscle-growth>
- [3] Training Volume Landmarks for Muscle Growth
- – RP Strength - <https://rpstrength.com/blogs/articles/training-volume-landmarks-muscle-growth>
- [4] No Time to Lift? Designing Time-Efficient Training Programs for Strength and Hypertrophy: A Narrative Review - <https://pubmed.ncbi.nlm.nih.gov/PMC8449772>
- [5] Chapter 18: Periodization - <https://traineracademy.org/cpt-textbook/periodization/>
- [6] Periodization Training Simplified: A Strategic Guide | NASM Blog - <https://blog.nasm.org/periodization-training-simplified>
- [7] Rep Ranges: How Many Reps for Muscle vs Strength | TTraining - <https://ttrening.com/learn/articles/rep-ranges-explained>
- [8] American College of Sports Medicine position stand. Progression models in resistance training for healthy adults - PubMed - <https://pubmed.ncbi.nlm.nih.gov/19204579/>
- [9] Program Design for Fat Loss - Clean Health - <https://cleanhealth.edu.au/blog/other/program-design-for-fat-loss/>
- [10] Programming For Fat Loss - TRAINFITNESS - <https://train.fitness/personal-trainer-blogs/programming-for-fat-loss>
- [11] Full article: Comparison of concurrent, resistance, or aerobic training on body fat loss: a systematic review and meta-analysis - <https://www.tandfonline.com/doi/full/10.1080/15502783.2025.2507949>
- [12] Adding Different Exercise Training Volumes to a Diet-Induced Weight Loss Facilitates Fat Loss and Maintains Fat-Free Mass in a Dose-Depending Fashion in Persons with Newly Diagnosed Type 2 Diabetes: Secondary Findings from the DOSE-EX Multi-Arm, Parallel-Group, Randomized Trial - <https://www.medrxiv.org/content/10.1101/2023.06.04.23290749>
- [13] Hypertrophy: Rep Ranges — Evidence Across the Spectrum — Hypertrophy Tower - <https://hypertrophy.towerofrecords.com/hypertrophy/rep-ranges>
- [14] Unlock maximal muscle growth using the science-backed Hypertrophy Hierarchy. Learn the optimal Volume (10+ sets), Frequency, and Effort (RPE 8) for serious gain - <https://www.myactivenutrition.com/blog/maximizing-muscle-the-definitive-evidence-based-guide-to-h...>
- [15] Hypertrophy: The Optimum Rep Range - <https://www.scienceforsport.com/hypertrophy-the-optimum-rep-range/>
- [16] What New Studies Show About Rep Ranges for Size & Strength
- – SET FOR SET - <https://www.setforset.com/blogs/news/rep-ranges-for-hypertrophy-strength>
- [17] Understanding the Progression from Beginner to Intermediate Lifter: A - <https://gym-mikolo.com/blogs/home-gym/understanding-the-progression-from-beginner-to-intermediate...>
- [18] Beginner to Intermediate Strength Training Program | 4 Day Gym Workout Plan - <https://warrioraddict.com/blogs/fitness-activewear-news/beginner-to-intermediate-strength-trainin...>
- [19] Novice, Intermediate, and Advanced Strength Training | Barbell Medicine - <https://www.barbellmedicine.com/blog/novice-intermediate-advanced-strength-training/>
- [20] Effects of the Menstrual Cycle and Hormonal Contraceptive Use on Metabolic Outcomes, Strength Performance, and Recovery: A Narrative Review - <https://pubmed.ncbi.nlm.nih.gov/PMC11278889>
- [21] Bidirectional Interactions between the Menstrual Cycle, Exercise Training, and Macronutrient Intake in Women: A Review - <https://pubmed.ncbi.nlm.nih.gov/PMC7910908>
- [22] Free Weights vs. Machines: 8 Benefits, Form Tips, Weights, and More - <https://www.healthline.com/health/fitness-exercise/free-weights-vs-machines>
- [23] Barbells vs Dumbbells: What's Better For Your Workout?
- – Onnit - <https://www.onnit.com/blogs/the-edge/barbells-vs-dumbbells-what-s-better-for-your-workout>
- [24] Current evidence shows no influence of women's menstrual cycle phase on acute strength performance or adaptations to resistance exercise training - <https://pubmed.ncbi.nlm.nih.gov/PMC10076834>
- [25] Effects of follicular versus luteal phase-based strength training in young women - <https://pubmed.ncbi.nlm.nih.gov/PMC4236309>
- [26] Links Between Testosterone, Oestrogen, and the Growth Hormone/Insulin-Like Growth Factor Axis and Resistance Exercise Muscle Adaptations - <https://pubmed.ncbi.nlm.nih.gov/PMC7844366>

- [27] Comparison of upper body strength gains between men and women after 10 weeks of resistance training - <https://pubmed.ncbi.nlm.nih.gov/PMC4756754>
- [28] Muscle strength gains per week are higher in the lower-body than the upper-body in resistance training experienced healthy young women—A systematic review with meta-analysis Muscle strength gains are higher in the lower-body than the upper-body in resistance training women - <https://pubmed.ncbi.nlm.nih.gov/PMC10101404>
- [29] Velocity-based resistance training: do women need greater velocity loss to maximize adaptations? - <https://pubmed.ncbi.nlm.nih.gov/PMC9012837>
- [30] Sex Differences in Neuromuscular Adaptations Following 12 Weeks of Kettlebell Swing Training - <https://pubmed.ncbi.nlm.nih.gov/PMC11524543>
- [31] Strength Standards by Lift, Bodyweight & Gender (2026) — Stronger - <https://www.strongermobileapp.com/blog/strength-standards>
- [32] Strength Standards - Calcfmi - <https://calcfmi.com/strength-standards/>
- [33] Strength Standards Calculator - Squat, Bench, Deadlift & Press - <https://traincalc.com/calculators/strength-standards>
- [34] Strength Standards - Lean FFMI - <https://leanffmi.com/resources/strength-standards/>
- [35] Squat Standards for Men and Women (lb) - Strength Level - <https://strengthlevel.com/strength-standards/squat/lb>
- [36] Strength Standards: Lifting Standards by Age & Weight | Legion - <https://legionathletics.com/strength-standards/>
- [37] Understanding HRV (Heart Rate Variability) for Optimal Recovery
- – Speediance Europe - <https://speediance.eu/blogs/news/understanding-hrv-heart-rate-variability-for-optimal-recovery>
- [38] Frontiers | Mobile Heart Rate Variability Biofeedback Improves Autonomic Activation and Subjective Sleep Quality of Healthy Adults – A Pilot Study - <https://www.frontiersin.org/journals/physiology/articles/10.3389/fphys.2022.821741/full>
- [39] Heart Rate Variability (HRV) and Recovery — Sapphire Physical Therapy | Missoula, MT - <https://www.sapphirephysicaltherapy.com/blog/heart-rate-variability-hrv-and-recovery>
- [40] HRV Training Guide: Recovery, Strain, and Overtraining - <https://www.whoop.com/us/en/thelocker/heart-rate-variability-training/>
- [41] Microsoft Word - Recovery white paper_15.6.2015 - https://www.firstbeat.com/wp-content/uploads/2015/10/Recovery-white-paper_15.6.2015.pdf
- [42] How to Improve HRV: 12 Ways to Boost Heart Rate Variability - <https://sleep.me/post/how-to-increase-hrv>
- [43] Heart Rate Variability Applications in Strength and Conditioning: A Narrative Review - <https://pubmed.ncbi.nlm.nih.gov/PMC11204851>
- [44] Interpret HRV Training: Ultimate 2026 Guide to Performance - <https://www.trainingpeaks.com/blog/how-to-interpret-hrv-training/>
- [45] RPE vs. Velocity Feedback: Autoregulation Without the Tech - <https://www.appliedperformance.co/blog/rpe-vs-velocity-feedback-can-you-autoregulate-without-the-...>
- [46] Assess and monitor your fatigue and readiness with VBT - <https://www.vbtcoach.com/blog/readiness-autoregulation-with-vbt>
- [47] Autoregulation in strength training: avoid overtraining and undertraining - <https://gymaware.com/autoregulation-in-strength-training/>
- [48] Reps in Reserve (RIR): Understanding, Comparison, and Integration with VBT - GymAware - <https://gymaware.com/ reps-in-reserve/>
- [49] RPE, RIR, and Bar Speed: The 3 Most Effective Autoregulation Tools for
- – SET FOR SET - <https://www.setforset.com/blogs/news/autoregulation-tools-for-strength-training>
- [50] What is RPE in Lifting? (Rate of Perceived Exertion Explained) - <https://www.barbellmedicine.com/blog/autoregulation-and-rpe-part-i/>
- [51] “Which autoregulation strategy should we use?” The effects of velocity loss thresholds on RPE and their individual differences in the back squat - Chieh-Ying Chiang, Yi-Chien Chiang, Wei-Yin Chen, Jonathon Weakley, 2025 - <https://journals.sagepub.com/doi/10.1177/17479541251339905>
- [52] FINAL DEGREE PROJECT - https://addi.ehu.es/bitstream/handle/10810/60866/TFG_Oficialdegi.pdf?sequence=1
- [53] A Brief Review on Concurrent Training: From Laboratory to the Field - <https://pubmed.ncbi.nlm.nih.gov/PMC6315763>
- [54] Order of same-day concurrent training influences some indices of power development, but not strength, lean mass, or aerobic fitness in healthy, moderately-active men after 9 weeks of training | PLOS One - <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0233134>