

Executive Quality Summary

6

Total Observations


1

Datasets Audited


3

High Severity Flags

Issue Severity Distribution



Top Data Quality Observations



Clinical Risk Analysis

Based on the AI-driven validation, the primary risks identified relate to high-severity clinical anomalies. These observations often stem from legacy data entry rituals or missing interoperability checks. Immediate attention to 'High' severity flags is recommended to ensure patient safety and diagnostic accuracy.

Detailed Audit Logs

DATASET	COLUMN	OBSERVATION	SEVERITY	STATUS
dmsap_test.csv	Parasite_Density	Missing value detected.	High	AUDITED
dmsap_test.csv	All	Duplicate row detected (2 times). (Val: Duplicate	Medium	AUDITED
dmsap_test.csv	Body_Temp	[SAFE IMPROVEMENT] Clinical temperature outlier. (	High	AUDITED
dmsap_test.csv	Age	[SAFE IMPROVEMENT] Age exceeds clinical protocol l	High	AUDITED
dmsap_test.csv	Parasite_Density	[SAFE IMPROVEMENT] Parasitaemia density exceeds li	Medium	AUDITED
dmsap_test.csv	Wt	[SAFE IMPROVEMENT] Weight-Age mismatch detected. (	Medium	AUDITED

Strategic Quality Roadmap

Short-term

Purge duplicate records and backfill missing critical identifiers identified in Section 4.

Mid-term

Implement real-time range validation (e.g., using FHIR standards) at the source of truth.

Long-term

Train the clinical staff on the impact of data quality on downstream AI research outcomes.