



concentrating NaCl (MC2.05.07.24) · poly1: $B(c) = \beta_0 + \beta_1 \cdot c$ · $\beta_0 \times \sigma$ square slice (B as a function of interfacial conc.)


1. mass
[g²]




2. ICP retentate conc.
[mM²]




3. ICP permeate conc.
[mM²]




4. conductivity-probe retentate
[μS²]




5. conductivity-probe permeate
[μS²]



6. cond.-derived retentate conc.
[mM²]



7. cond.-derived permeate conc.
[mM²]



concentrating NaCl · poly1
 $B(c) = \beta_0 + \beta_1 \cdot c$


Axis: β_0 (y) × σ (x)
 $L_p = 9.99$ fixed
other coeffs held at the
poly1 fit (warm start).

▲ = per-channel optimum
-- = fitted β_0 value
lines = log₁₀ WSSE


Square forward-solve slice
(no per-node optimization).
Blank panels = the pinned
coefficients drive $B(c)$ out of
the physical region (no
feasible square solve there).

concentrating NaCl (MC2.05.07.24) · poly1: $B(c) = \beta_0 + \beta_1 \cdot c$ · $\beta_1 \times \sigma$ square slice (B as a function of interfacial conc.)


1. mass
[g²]




2. ICP retentate conc.
[mM²]




3. ICP permeate conc.
[mM²]




4. conductivity-probe retentate
[μS²]




5. conductivity-probe permeate
[μS²]



6. cond.-derived retentate conc.
[mM²]



7. cond.-derived permeate conc.
[mM²]



concentrating NaCl · poly1
 $B(c) = \beta_0 + \beta_1 \cdot c$

Axis: β_1 (y) × σ (x)

$L_p = 9.99$ fixed

other coeffs held at the
poly1 fit (warm start).








▲ = per-channel optimum

-- = fitted β_1 value

lines = log₁₀ WSSE

Square forward-solve slice
(no per-node optimization).
Blank panels = the pinned
coefficients drive $B(c)$ out of
the physical region (no
feasible square solve there).

concentrating NaCl (MC2.05.07.24) · poly2: $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2$ · $\beta_0 \times \sigma$ square slice (B as a function of interfacial conc.)




concentrating NaCl · poly2
 $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2$

Axis: β_0 (y) × σ (x)
 $L_p = 9.98$ fixed
 other coeffs held at the
 poly2 fit (warm start).


▲ = per-channel optimum
 - - = fitted β_0 value
 lines = log₁₀ WSSE

Square forward-solve slice
 (no per-node optimization).
 Blank panels = the pinned
 coefficients drive $B(c)$ out of
 the physical region (no
 feasible square solve there).

concentrating NaCl (MC2.05.07.24) · poly2: $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2$ · $\beta_1 \times \sigma$ square slice (B as a function of interfacial conc.)




concentrating NaCl (MC2.05.07.24) · poly2: $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2 \cdot \beta_2 \times \sigma$ square slice (B as a function of interfacial conc.)




concentrating NaCl (MC2.05.07.24) · poly3: $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2 + \beta_3 \cdot c^3$ · $\beta_0 \times \sigma$ square slice (B as a function of interfacial conc.)


**1. mass
[g²]**




**2. ICP retentate conc.
[mM²]**




**3. ICP permeate conc.
[mM²]**




**4. conductivity-probe retentate
[μS²]**




**5. conductivity-probe permeate
[μS²]**



**6. cond.-derived retentate conc.
[mM²]**



**7. cond.-derived permeate conc.
[mM²]**



concentrating NaCl · poly3
 $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2 + \beta_3 \cdot c^3$


Axis: β_0 (y) × σ (x)
 $L_p = 10.03$ fixed
other coeffs held at the
poly3 fit (warm start).

▲ = per-channel optimum
-- = fitted β_0 value
lines = log₁₀ WSSE


Square forward-solve slice
(no per-node optimization).
Blank panels = the pinned
coefficients drive $B(c)$ out of
the physical region (no
feasible square solve there).

concentrating NaCl (MC2.05.07.24) · poly3: $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2 + \beta_3 \cdot c^3$ · $\beta_1 \times \sigma$ square slice (B as a function of interfacial conc.)


1. mass
[g²]




2. ICP retentate conc.
[mM²]




3. ICP permeate conc.
[mM²]




4. conductivity-probe retentate
[μS²]




5. conductivity-probe permeate
[μS²]



6. cond.-derived retentate conc.
[mM²]



7. cond.-derived permeate conc.
[mM²]



concentrating NaCl · poly3
 $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2 + \beta_3 \cdot c^3$


Axis: β_1 (y) × σ (x)
 $L_p = 10.03$ fixed
other coeffs held at the
poly3 fit (warm start).

▲ = per-channel optimum
-- = fitted β_1 value
lines = log₁₀ WSSE


Square forward-solve slice
(no per-node optimization).
Blank panels = the pinned
coefficients drive $B(c)$ out of
the physical region (no
feasible square solve there).

concentrating NaCl (MC2.05.07.24) · poly3: $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2 + \beta_3 \cdot c^3$ · $\beta_2 \times \sigma$ square slice (B as a function of interfacial conc.)


**1. mass
[g²]**




**2. ICP retentate conc.
[mM²]**




**3. ICP permeate conc.
[mM²]**




**4. conductivity-probe retentate
[μS²]**




**5. conductivity-probe permeate
[μS²]**



**6. cond.-derived retentate conc.
[mM²]**



**7. cond.-derived permeate conc.
[mM²]**



concentrating NaCl · poly3
 $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2 + \beta_3 \cdot c^3$


Axis: β_2 (y) × σ (x)
 $L_p = 10.03$ fixed
other coeffs held at the
poly3 fit (warm start).

▲ = per-channel optimum
- - = fitted β_2 value
lines = $\log_{10}$ WSSE


Square forward-solve slice
(no per-node optimization).
Blank panels = the pinned
coefficients drive $B(c)$ out of
the physical region (no
feasible square solve there).

concentrating NaCl (MC2.05.07.24) · poly3: $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2 + \beta_3 \cdot c^3$ · $\beta_3 \times \sigma$ square slice (B as a function of interfacial conc.)


**1. mass
[g²]**




**2. ICP retentate conc.
[mM²]**




**3. ICP permeate conc.
[mM²]**




**4. conductivity-probe retentate
[μS²]**




**5. conductivity-probe permeate
[μS²]**



**6. cond.-derived retentate conc.
[mM²]**



**7. cond.-derived permeate conc.
[mM²]**



concentrating NaCl · poly3
 $B(c) = \beta_0 + \beta_1 \cdot c + \beta_2 \cdot c^2 + \beta_3 \cdot c^3$


Axis: β_3 (y) × σ (x)
 $L_p = 10.03$ fixed
other coeffs held at the
poly3 fit (warm start).

▲ = per-channel optimum
-- = fitted β_3 value
lines = log₁₀ WSSE


Square forward-solve slice
(no per-node optimization).
Blank panels = the pinned
coefficients drive $B(c)$ out of
the physical region (no
feasible square solve there).

concentrating NaCl (MC2.05.07.24) · sat: $B(c) = B_{\infty} \cdot (1 - e^{(-c/c^*)}) \cdot B_{\infty} \times \sigma$ square slice (B as a function of interfacial conc.)


1. mass
[g²]




2. ICP retentate conc.
[mM²]




3. ICP permeate conc.
[mM²]




4. conductivity-probe retentate
[μS²]




5. conductivity-probe permeate
[μS²]



6. cond.-derived retentate conc.
[mM²]



7. cond.-derived permeate conc.
[mM²]



concentrating NaCl · sat
 $B(c) = B_{\infty} \cdot (1 - e^{(-c/c^*)})$

Axis: B_{∞} (y) × σ (x)
 $L_p = 9.95$ fixed
other coeffs held at the
sat fit (warm start).

▲ = per-channel optimum
-- = fitted B_{∞} value
lines = log₁₀ WSSE

Square forward-solve slice
(no per-node optimization).
Blank panels = the pinned
coefficients drive $B(c)$ out of
the physical region (no
feasible square solve there).

concentrating NaCl (MC2.05.07.24) · sat: $B(c) = B_{\infty} \cdot (1 - e^{(-c/c^*)}) \cdot c^* \times \sigma$ square slice (B as a function of interfacial conc.)

