

Quiz 2

Problem 1 (7 points). Use the following three facts about determinants to compute the determinant of a matrix using row operations.

- If B is diagonal, then $\det(B) = b_{11} \cdot b_{22} \cdots b_{nn}$.
- If B arises from A by a type I row operation, i.e., interchanging two rows, then $\det(B) = -\det(A)$.
- If B arises from A by a type III row operation, i.e., $r_i + ar_j \rightarrow r_i$, that is, row i is replaced by row i plus a scalar multiple of row j , where $i \neq j$. Then $\det(A) = \det(B)$.

Compute $\det(A)$ by:

- Reducing A to a diagonal matrix B using only type I and III operations.
- Keep track of how many row swaps were made.
- Compute $\det(B)$ by multiplying the diagonal elements of B .

$$A = \begin{bmatrix} 8 & 6 & -3 & 20 \\ 4 & 2 & -5 & -7 \\ 8 & 2 & 7 & 20 \\ 4 & 2 & -11 & -4 \end{bmatrix}$$

Show the work for the above computation here.

On your own, don't include this in the quiz, try computing this determinant by expanding on a row or column.


Discuss which method, "expansion along a row or column" or "using elementary row operations" is, in general, a faster method of computing a determinant.

This page left blank

Problem 2 (7 points). Let A be as above, consider $A\mathbf{x} = \mathbf{b}$ where $\mathbf{b} = (31, 2, 21, 11)$. Find x_1 using Cramer's rule. (You may use MATLAB/Octave to compute the determinants, but write out what you are computing.)

|

Problem 3 (7 points). Given vectors $v = \begin{bmatrix} v_0 \\ v_1 \end{bmatrix}$ and $u = \begin{bmatrix} u_0 \\ u_1 \end{bmatrix}$ it is true that $|\det(u, v)| = \left| \det \begin{bmatrix} u_0 & v_0 \\ u_1 & v_1 \end{bmatrix} \right|$ is the area of the parallelogram formed by u and v in the natural way. Thus we know:



$$\text{area} = \frac{1}{2} \det(u, v) = \frac{1}{2} \det \begin{bmatrix} c-a & e-a \\ d-b & f-b \end{bmatrix}$$

Show that the area of a triangle with vertices (a, b) , (c, d) , and (e, f) is given by

$$\text{area} = \frac{1}{2} \det \begin{bmatrix} 1 & 1 & 1 \\ a & c & e \\ b & d & f \end{bmatrix}$$

Problem 4 (14 points). submit the completion certificate for the OnRamp tutorial from MATLAB.