

Name: _____

Quiz 1 - MAT345

Problem 1 (15 points; 3 points each). Decide if each of the following is true or false.

- (a) _____ If A and B are $n \times n$ symmetric matrices, then AB is symmetric.

This is false. $(AB)^T = B^T A^T = BA$, but for symmetry, we would need $(AB)^T = AB$, and there is no reason that $AB = BA$.

In fact, generating two random 2×2 symmetric matrices will most likely give an example. Here is the first pair I generated:

$$A = \begin{bmatrix} 6 & -2 \\ -2 & -4 \end{bmatrix} \quad B = \begin{bmatrix} -4 & 4 \\ 4 & 2 \end{bmatrix} \quad AB = \begin{bmatrix} -32 & 20 \\ -8 & -16 \end{bmatrix}$$

Clearly, A and B are symmetric, while AB is not.

- (b) _____ An echelon form of a matrix is unique.

This is false. There is a unique reduced echelon form. Both

$$E_1 = \begin{bmatrix} 2 & 3 & 1 \\ 0 & 3 & 2 \end{bmatrix}$$

and

$$E_2 = \begin{bmatrix} 2 & 0 & -1 \\ 0 & 3 & 2 \end{bmatrix}$$

are both echelon and each can be got from the other by an elementary row operation.

- (c) _____ If A is a 4×2 matrix, then $A\mathbf{x} = \mathbf{0}$ has at least two free variables.

This is false. A 4×2 matrix corresponds to an *overdetermined* system, usually with no solutions. Had this been a 2×4 matrix, then there must be at least two free variables. This would be an *underdetermined* system.

- (d) _____ $A^2 - B^2 = (A - B)(A + B)$ for all square $n \times n$ matrices A and B .

This is false. $(A - B)(A + B) = (A - B)A + (A - B)B = A^2 - BA + AB - B^2$, but as $AB \neq BA$ can occur, there is no reason that $AB - BA = 0$.

- (e) _____ For A a 4×6 matrix, let E be the matrix so that EA is the result of the row op $R_1 - 3R_2 \rightarrow R_1$ applied to A . Then the first row of E is $[1 \quad -3 \quad 0]$.

This is false. Although I had intended it to be true, somehow, I switched 3 and 4.

I did not intend this to be a trick problem, so I will accept either answer

here.

$$\begin{bmatrix} 1 & -3 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} \begin{bmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \mathbf{a}_3 \\ \mathbf{a}_4 \end{bmatrix} = \begin{bmatrix} (1) \cdot \mathbf{a}_1 + (-3) \cdot \mathbf{a}_2 + (0) \cdot \mathbf{a}_3 + (0) \cdot \mathbf{a}_4 \\ \vdots \end{bmatrix}$$

The correct answer is $\begin{bmatrix} 1 & -3 & 0 & 0 \end{bmatrix}$

Problem 2 (25 points). Solve $A\mathbf{x} = \mathbf{0}$ for

$$A = \begin{bmatrix} 4 & 8 & 5 & -3 \\ 5 & 10 & 2 & -8 \\ -3 & -6 & -4 & 2 \end{bmatrix}$$

Follow the procedure discussed in class

- (10 points) Use elementary row ops to reduce to an echelon matrix.
- (10 points) Write down the resulting triangular system and use back-substitution to solve.
- (5 points) Write out your solution as a linear combination of vectors.

Reduction to echelon form requires 3 row manipulations:

This part is 10/25 points.

$$\begin{aligned} \begin{bmatrix} 4 & 8 & 5 & -3 \\ -3 & -6 & -4 & 2 \\ 5 & 10 & 2 & -8 \end{bmatrix} &\xrightarrow{\substack{R_2 \leftarrow R_2 - (-3/4)R_1 \\ R_3 \leftarrow R_3 - (5/4)R_1}} \begin{bmatrix} 4 & 8 & 5 & -33 \\ 0 & 0 & -1/4 & -1/4 \\ 0 & 0 & -17/4 & -17/4 \end{bmatrix} \\ &\xrightarrow{R_3 \leftarrow R_3 - (17)R_2} \begin{bmatrix} 4 & 8 & 5 & -3 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

Here x_2 and x_4 are free variables, so let $x_2 = s$ and $x_4 = t$, then we get (by back-substitution)

$$\begin{aligned} x_3 &= -t \\ 4x_1 &= -8s - 5(-t) + 3t = -8s + 8t \rightarrow x_1 = -2s + 2t \end{aligned}$$

This is 10/25 points.

So we have

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} -2s + 2t \\ s \\ -t \\ t \end{bmatrix} = s \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} 2 \\ 0 \\ -1 \\ 1 \end{bmatrix}$$

This final answer is the remaining 5/25 points.