

Math 571 - Homework 2

Richard Ketchersid

Problem 2.1 (R:2:2*). A complex number γ is **algebraic** iff γ is a root to a polynomial with integer coefficients. Prove that there are complex numbers that are not algebraic.

The following are two fun facts (not related to the question, but just for intellectual curiosity):

Let $\mathbb{A} \subset \mathbb{C}$ be the set of algebraic numbers.

1. \mathbb{A} is a field.
2. \mathbb{A} is algebraically closed, that is, if α is a root of a polynomial in $\mathbb{A}[x]$, then $\alpha \in \mathbb{A}$. So in the definition of algebraic numbers you can use any ring of coefficients R , $\mathbb{Z} \subseteq R \subseteq \mathbb{A}$.

[Here](#) is a write-up of the proofs, you should have the background to read this, but it is not an easy read.

The intent here was for you to do a counting argument. There are only countably many polynomials with integer coefficients and each has only finitely many roots, hence there are only countably many algebraic numbers.

Definition 1. A set $S \subseteq X$ is **discrete** iff every point in S is isolated.

Problem 2.2 (R:2:5*). Prove the following for discrete $S \subset \mathbb{R}$:

- a) $\text{Lim}(S) \cap S = \emptyset$ and S is countable.

For each $x \in S$ we find integer n_x so that $N_{\frac{1}{n_x}}(x) \cap (S - \{x\}) = \emptyset$. We can find $q_x \in \mathbb{Q} \cap N_{\frac{1}{2n_x}}(x)$. The map $x \mapsto (n_x, q_x)$ is injective since if $x, x' \in S$ and $(n_x, q_x) = (n_{x'}, q_{x'})$, then $|x - x'| \leq |x - q_x| + |q_x - x'| < \frac{1}{2n_x} + \frac{1}{2n_x} = \frac{1}{n_x}$ and hence $x' \in N_{\frac{1}{n_x}}(x)$. So S is countable.

Suppose $x \in \text{Lim}(S)$, then $S \cup \{x\}$ has a non-isolated point, namely, x , so $\text{Lim}(S) \cap S = \emptyset$.

- b) There is countable set $A \subset \mathbb{R}$ so that $\text{Lim}(A) = \text{Cl}(S)$.

From the preceding we can write $S = \{s_i \mid i \in \mathbb{N}\}$ and get $n_0 < n_1 < \dots$ so the $N_{\frac{1}{n_i}}(x) \cap S = \{x\}$.

Let A_i be a countable discrete set in $N_{\frac{1}{n_i}}$, with $\{s_i\} = \text{Lim}(A_i)$. Namely, take a sequence of distinct points converging to s_i .

Let $A = \bigcup_{i \in \mathbb{N}} A_i$. Clearly, $S \subseteq \text{Lim}(A)$ and so $\text{Cl}(S) \subseteq \text{Lim}(A)$.

Now we want to see that $\text{Lim}(A) \subset \text{Cl}(S)$, so let $z \in \text{Lim}(A)$ and $z = \lim_i z_i$ for $z_i \in A$. If $z \in S$, there is nothing to do, so assume $z \notin S$. Each $z_i \in A_j$ for some i , if for any j , $\{z_i \mid z_i \in A_j\}$ is infinite, then clearly $z = x_j = \lim\{z_i\}_{z_i \in A_j}$. So we see that for each j , $\{z_i \mid z_i \in A_j\}$ is finite and, in fact, the same argument shows this to be true for any infinite subset (subsequence) $\{z_{i_j} \mid j \in \mathbb{N}\}$. For any $\delta > 0$ since $\{i \mid z_i \in N_\delta(z)\}$ is infinite, we know that $\{j \mid A_j \cap N_{\delta/2}(z) \neq \emptyset\}$ is infinite. Choosing such a j so that $\frac{1}{n_j} < \delta/2$ will result in $z_k \in N_{\delta/2}(s_j)$ and $z_k \in N_{\delta/2}(z)$ so $s_j \in N_\delta(z)$. Thus $z \in \text{Lim}(S)$. So $z \in S \cup \text{Lim}(S) = \text{Cl}(S)$.

- c) Give an example of a discrete set S where there is no set A such that $\text{Lim}(A) = S$.

Clearly, $\text{Lim}(S) \subseteq \text{Lim}(A)$, since $\text{Lim}(\text{Lim}(A)) \subseteq \text{Lim}(A)$. So just take S with $\text{Lim}(S) = S \neq \emptyset$.

For the following use the definition that I provided for $\text{Cl}(E)$, namely, $\text{Cl}(E) = \bigcap \{F \mid F \text{ is closed and } E \subseteq F\}$.

Problem 2.3 (R:2:6). For X a metric space and $E \subseteq X$, show that

- a) $\text{Cl}(E) = E \cup \text{Lim}(E)$.

First show $\text{Cl}(E) \subseteq E \cup \text{Lim}(E)$. Let $x \in \text{Cl}(E)$. If $x \in E$ we are done. So assume $x \notin E$. Let O be open with $x \in O$. Towards a contradiction, suppose $O \cap E = \emptyset$, then $O^c \supset E$ and is closed and so $\text{Cl}(E) \subseteq O^c$. But then $x \notin \text{Cl}(E)$. A contradiction. So $O \cap E \neq \emptyset$ for all open nbhds of x , thus $x \in \text{Lim}(E)$.

Next we see $E \cup \text{Lim}(E) \subseteq \text{Cl}(E)$. Let $F \supset E$ be closed, it suffices to see that $\text{Lim}(E) \subseteq F$. Let $x \in \text{Lim}(E)$. If $x \notin F$, then F^c is an open nbhd of x with $F^c \cap E = \emptyset$, so $x \notin \text{Lim}(E)$, a contradiction.

- b) $\text{Lim}(E)$ is closed.

From (a), $\text{Cl}(\text{Lim}(E)) = \text{Lim}(E) \cup \text{Lim}(\text{Lim}(E))$. So if we show $\text{Lim}(\text{Lim}(E)) \subseteq \text{Lim}(E)$ we are done. Let $x \in \text{Lim}(\text{Lim}(E))$. Let O be an open nbhd of x , then $O \cap \text{Lim}(E) \neq \emptyset$. Let U be an open nbhd of $y \in O \cap \text{Lim}(E)$ so that $U \subset O$. Then $U \cap E \neq \emptyset$, so $O \cap E \neq \emptyset$ and this is what was required to see that $x \in \text{Lim}(E)$.

- c) Either show or give a counterexample to $\text{Lim}(E) = \text{Lim}(\text{Cl}(E))$.

$\text{Lim}(E) \subseteq \text{Lim}(\text{Cl}(E))$ simply because $A \subseteq B \implies \text{Lim}(A) \subseteq \text{Lim}(B)$. Let $x \in \text{Lim}(\text{Cl}(E))$ and let O be an open nbhd of x , then $O \cap (E \cup \text{Lim}(E)) \neq \emptyset$. If $O \cap E \neq \emptyset$ then we are done. Else $O \cap \text{Lim}(E) \neq \emptyset$. In this case we argue as we did above. Let $y \in \text{Lim}(E) \cap O$. Let $U \subset O$ be nbhd of y , then $U \cap E \neq \emptyset$, so $O \cap E \neq \emptyset$.

Problem 2.4 (R:2:9*). Let X be a metric space, or just any topological space. Are the following true for all $E \subseteq X$? For each either prove the statement true or give a counterexample. For a counterexample you must provide both X and E .

- a) $\text{Int}(E)^c = \text{Cl}(E^c)$.

Let's try to prove this. there are, as usual, two things to prove here.

$\text{Int}(E)^c \subseteq \text{Cl}(E^c)$: Let $x \in \text{Int}(E)^c$, so $x \notin \text{Int}(E)$. This means every neighborhood of x contains points in E^c . This means $x \in \text{Cl}(E^c)$.

$\text{Cl}(E^c) \subseteq \text{Int}(E)^c$: Let $x \in \text{Cl}(E^c)$ so every nbhd of x meets E^c , so $x \notin \text{Int}(E)$, thus $x \in \text{Int}(E)^c$.

The following two arguments came up in class:

Alternate 1:

$$\begin{aligned}
 x \in \text{Cl}(E^c) &\iff \forall N (N \text{ a nbhd of } x \implies N \cap E^c \neq \emptyset) \\
 &\iff \forall N (N \text{ a nbhd of } x \implies N \not\subseteq E) \\
 &\iff \neg \exists N (N \text{ a nbhd of } x \text{ and } N \subseteq E) \\
 &\iff x \notin \text{Int}(E) \\
 &\iff x \in (\text{Int}(E))^c
 \end{aligned}$$

Alternate 2:

$$\begin{aligned}
 \text{cl}(E^c) &= \bigcap \{F \mid F \text{ is closed and } F \supseteq E^c\} \\
 &= \left(\bigcup \{F^c \mid F \text{ is closed and } F \supseteq E^c\} \right)^c \\
 &= \left(\bigcup \{F^c \mid F^c \text{ is open and } F^c \subseteq E\} \right)^c \\
 &= \left(\bigcup \{O \mid O \text{ is open and } O \subseteq E\} \right)^c \\
 &= (\text{Int}(E))^c
 \end{aligned}$$

b) $\text{Cl}(E) = \text{Int}(E^c)^c$?

This is true and we can just apply (a) here. $\text{Cl}(E) = \text{Cl}((E^c)^c) = \text{Int}(E^c)^c$. This clearly also gives $\text{Cl}(E)^c = \text{Int}(E^c)$.

c) $\text{Cl}(E) = \text{Cl}(\text{Int}(E))$?

This is false. Just take $X = \mathbb{R}$ and $E = \mathbb{Q}$, then $\text{Cl}(\mathbb{Q}) = \mathbb{R}$ but $\text{Cl}(\text{Int}(E)) = \text{Cl}(\emptyset) = \emptyset$.

d) $\text{Int}(E) = \text{Int}(\text{Cl}(E))$

This is just as the previous, same counterexample shows this to be false. $\text{Int}(\mathbb{Q}) = \emptyset \neq \text{Int}(\text{Cl}(\mathbb{Q})) = \text{Int}(\mathbb{R}) = \mathbb{R}$.

An open set, E , is called a **regular open set** iff $E = \text{Int}(\text{Cl}(E))$. Similarly, a closed set, E , is **regular closed set** if $E = \text{Cl}(\text{Int}(E))$.

Let O be any open set, then ∂O is nowhere dense, that is, for all open U , there is $U' \subseteq U$ so that $\emptyset \neq U'$ and $U' \cap \partial O = \emptyset$. Let U be open and suppose $U \cap \partial O \neq \emptyset$. Let $U' = U \cap O$. Clearly, $\emptyset \neq U'$ and $U' \cap \partial O = \emptyset$, since $O \cap \partial O = \emptyset$.

Any non-empty closed nowhere-dense set, N , fails to be regular closed, and so N^c fails to be regular open. For example, the circle $S^1 \subset \mathbb{R}^2$ is the boundary of the open unit disk and thus is closed nowhere-dense, hence not regular-closed. Correspondingly, $G = \mathbb{R}^2 - S^1$ is open, but not regular open.

Definition 2. A metric space X is **separable** iff there is a countable $E \subseteq X$ with E dense in X .

Problem 2.5 (R:2:22). Show the \mathbb{R}^k is separable.

It is easy to see that \mathbb{Q}^k is dense in \mathbb{R}^k . One way is the following, use basic open "boxes" of the form $\prod_{i=1}^k (a_i, b_i)$ for the basic open sets, instead of open balls. The fact that $\mathbb{Q} \cap (a_i, b_i) \neq \emptyset$ immediately yields that $\mathbb{Q}^k \cap \prod_{i=1}^k (a_i, b_i) \neq \emptyset$.

Definition 3. A set \mathcal{B} of open sets is called a **base** for X iff for all $x \in X$ and open set U with $x \in U$, there is $V \in \mathcal{B}$ so that $x \in V \subset U$.

Problem 2.6 (R:2:23*). Prove that a metric space is separable iff it has a countable base.

If X is separable, let S be a countable dense set. Consider $N_{\frac{1}{i}}(s)$ for $s \in S$. Let $x \in X$ and O be an open nbhd of x . Take $N_\delta(x) \subseteq O$ and $s \in S$ with $d(s, x) < \delta/4$. Then $x \in N_{\frac{1}{m}}(s) \subseteq O$ with $\frac{1}{m} < \frac{\delta}{4}$. So the sets $N_{\frac{1}{i}}(s)$ do form a countable base.

If $\{O_i \mid i \in \mathbb{N}\}$ is a countable base, then just take $s_i \in O_i$ for all i , then $S = \{s_i \mid i \in \mathbb{N}\}$ is dense.

Problem 2.7 (R:2:24). Prove that if X is a metric space and every infinite sequence has a limit point, then X is separable. (See the hint in the text.)

For each integer $i > 0$ construct a maximal set $S_i = \{x_j^i\}_{j=0}^{k_i}$ so that $d(x_l^i, x_k^i) \geq \frac{1}{i}$. $k_i < \infty$ for all i since otherwise there would be an infinite sequence with no limit. By maximal here we mean that for any $x \in X$ $N_{\frac{1}{i}}(x) \cap S_i \neq \emptyset$, since otherwise we could add x to S_i and maintain the desired separation of elements.

Let $S = \{x_j^i \mid j \leq k_i \text{ and } i \in \mathbb{N}\}$. S is dense in X since for any $\delta > 0$ and any x , let $i \in \mathbb{N}$ so that $\frac{1}{i} < \delta$, then $S \cap N_\delta(x) \supseteq S \cap N_{\frac{1}{i}}(x) \neq \emptyset$ by construction.