

Math 571 - Homework 4

Richard Ketchersid

Problem 5.1 (B:4.2:2). Show that the following are equivalent for $f : X \rightarrow Y$, where X and Y are metric spaces.

- (a) f is continuous.
- (b) If (p_n) is a sequence from X and $p_n \rightarrow p$, then $f(p_n) \rightarrow f(p)$, equivalently, $\lim_n f(p_n) = f(\lim_n p_n)$.
- (c) If (p_n) is convergent, then $f(p_n)$ is convergent.

Hint: For (c) $\implies$ (b) let $q = \lim_n f(p_n)$ and consider $(p_0, p, p_1, p, p_3, p, \dots) = (\hat{p}_n)$, then $(f(\hat{p}_n)) = (f(p_0), f(p), f(p_1), f(p), \dots)$ must converge.

Problem 5.2 (B:4.2:3). Let X be a metric space.

- (a) Show that there are open subsets A, B of X , such that $A \cup B = X$ and $A \cap B = \emptyset$ if and only if there exists a continuous function $f : X \rightarrow \mathbb{R}$ such that $f(a) = 0$ for all $a \in A$ and $f(b) = 1$ for all $b \in B$.
- (b) Deduce that X is connected if and only if every continuous function $X \rightarrow \{0, 1\}$ is constant.
- (c) Show likewise that X is connected if and only if every continuous function $X \rightarrow \mathbb{Z}$ is constant.

Problem 5.3 (B:4.2:4). (a) Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function such that $f(x) > x$ for all x . For any positive integer n , let f^n denote the n -fold composite $f \circ f \circ \dots \circ f$. Show that for all $x, y \in \mathbb{R}$, there exists a positive integer n such that $f^n(x) > y$.

- (b) Does the conclusion of part (a) remain true if the assumption that f is continuous is removed?

Problem 5.4 (R:4.11). Suppose f is a uniformly continuous mapping of a metric space X into a metric space Y and prove that $\{f(x_n)\}$ is a Cauchy sequence in Y for every Cauchy sequence $\{x_n\}$ in X . Use this result to give an alternative proof of the theorem stated in Exercise 13.

Problem 5.5 (R:4.13). Let E be a dense subset of a metric space X , and let f be a uniformly continuous *real* function defined on E . Prove that f has a continuous extension from E to (see Exercise 5 for terminology). (Uniqueness follows from Exercise 4.)

Could the range space $\mathbb{R}^1$ be replaced by $\mathbb{R}^k$? By any compact metric space? By any complete metric space? By any metric space?

Hint: For each $p \in X$ and each positive integer n , let $V_n(p)$ be the set of all $q \in E$ with $d(p, q) < 1/n$. Use Exercise 9 to show that the intersection of the closures of the sets $f(V_1(p))$, $f(V_2(p))$, ..., consists of a single point, say $g(p)$, of $\mathbb{R}^1$. Prove that the function g so defined on X is the desired extension of f .

Better Hint: Use R:4.11

Problem 5.6 (R:4.18). Define the *Thomae Function* as follows

$$f(x) = \begin{cases} 0 & x \text{ is irrational} \\ \frac{1}{n} & x = \frac{m}{n}, \text{ where } n, m \in \mathbb{Z}, n > 0, \text{ and } \gcd(m, n) = 1 \end{cases}$$

- Show that f is discontinuous at exactly the rational points.
- Show that there is no $g : \mathbb{R} \rightarrow \mathbb{R}$ that is continuous at exactly the irrational points. (Baire Category)

Problem 5.7 (B:4.4:3). Amazingly, there are continuous functions $f : [0, 1] \rightarrow [0, 1]^k$ for any k . Here are a couple of nice videos on this topic:

- A great 3Blue1Brown video: [Hilbert's Curve: Is infinite math useful?](#)
- Eddie Woo discussing several constructions in a high school class: [Space-Filling Curves \(1 of 4: Peano Curve\)](#)

- Let $n > 1$. Show that for every point p of $[0, 1]^n$, the subset obtained by removing p from $[0, 1]^n$ is still connected. On the other hand, point to a result in Rudin showing that $[0, 1]$ does not have this property.
- Show how the existence of a one-to-one continuous map from $[0, 1]$ onto $[0, 1]^n$, together with the above pair of facts, would lead to a contradiction. (Hint: What do you know about one-to-one continuous maps of one compact metric space onto another?)

Problem 5.8 (B:4.3:4). (a) Suppose K is a compact metric space, and $f : K \rightarrow K$ is a map with the property that for every pair of distinct points $x, y \in K$ one has $d(f(x), f(y)) < d(x, y)$. Show that there exists a unique point $p \in K$ such that $f(p) = p$.

- Show by examples that the result of (a) is false if we put any of the noncompact metric spaces $[0, 1)$, $[0, +\infty)$ or $\mathbb{R}$ in place of the compact space K .

Problem 5.9 (R4.20/22). Let X be a metric space, and for any nonempty subset $E \subset X$, define the distance from a point $x \in X$ to E as $\rho_E(x) = \inf_{z \in E} d(x, z)$.

- Prove that $\rho_E(x)$ is a uniformly continuous function on X , and that $\rho_E(x) = 0$ if and only if $x \in \bar{E}$.

(b) Let A and B be disjoint, nonempty closed sets in X . Define:

$$f(p) = \frac{\rho_A(p)}{\rho_A(p) + \rho_B(p)}$$

Prove that f is a continuous function on X with range in $[0, 1]$, such that $f(p) = 0$ precisely on A and $f(p) = 1$ precisely on B .

You could skip this, but it adds context.

The next problem uses a tool that is also at the heart of some of the space-filling curves and is a good capstone to the techniques in this chapter. Consider the space $X = C([0, 1]) = \{f : [0, 1] \rightarrow [0, 1] \mid f \text{ is continuous}\}$. (You have studied this as an inner product space in MAT-512). X is provided with a norm, the L_1 -**norm** or **uniform** norm, which is different from the L_2 -norm, $\|f\|_2 = \sqrt{\int_0^1 f^2 dx}$ coming from the inner-product $\langle f, g \rangle = \int_0^1 fg dx$. The uniform norm is given by

$$\|f\|_1 = \sup\{|f(x)| \mid x \in [0, 1]\}$$

This gives rise, as all norms do, to a metric, $d : X \times X \rightarrow [0, 2]$ given by $d(f, g) = \|f - g\|_1 = \sup\{|f(x) - g(x)| \mid x \in [0, 1]\}$. It should be clear that the 1/2-norms and the resulting metrics are just the natural generalization to $C([0, 1]) \subset \mathbb{R}^{[0, 1]}$ of the corresponding norms/metrics on $\mathbb{R}^k$. (In fact, this is what you naturally do when approximating continuous functions by taking test points, a basic part of both Calculus and computer science.)

Note: By the extreme value theorem, $\sup\{|f(x)| \mid x \in [0, 1]\}$ can be replaced by $\max\{|f(x)| \mid x \in [0, 1]\}$.

Convergence in $C([0, 1])$ under the L_1 -norm is called **uniform convergence**, i.e., convergence in the uniform norm. Clearly, $f_i \rightarrow f$ in $C([0, 1])$ iff for all $\epsilon > 0$, there is $N > 0$, so that for all $i > N$, $\sup\{|f_i(x) - f(x)| \mid x \in [0, 1]\} < \epsilon$.

You must stop skipping if you chose to do so.

We can rephrase this as $f_i \rightarrow f$ uniformly if

$$(\forall \epsilon > 0)(\exists N > 0)(\forall i > N)(\forall x \in [0, 1])(|f_i(x) - f(x)| < \epsilon) \quad (\dagger)$$

Note that $f_i(x) \rightarrow f(x)$ **pointwise** iff

$$(\forall \epsilon > 0)(\forall x \in [0, 1])(\exists N > 0)(\forall i > N)(|f_i(x) - f(x)| < \epsilon)$$

So uniform convergence means we can find for each ϵ and N that **works uniformly** for all $x \in [0, 1]$.

Main Result: If f_i are continuous and $f_i \rightarrow f$ uniformly in the sense of $(\dagger)$, then f is continuous.

Proof. This would have made a nice exercise in itself. For $\epsilon > 0$ and take $N > 0$ so that for all $i > N$, $|f_i(x) - f(x)| < \frac{\epsilon}{3}$ for all $x \in [0, 1]$. Take $i > N$ and find δ so that $0 < |y - x| < \delta \implies |f_i(y) - f_i(x)| < \frac{\epsilon}{3}$. Now take $0 < |y - x| < \delta$, then

$$|f(y) - f(x)| \leq |f(y) - f_i(y)| + |f_i(y) - f_i(x)| + |f_i(x) - f(x)| < \epsilon$$

by the triangle inequality. So we have verified that

$$0 < |y - x| \implies |f(y) - f(x)| < \epsilon$$

Note: If $f_i \rightarrow f$ pointwise, but not uniformly, then f need not be continuous. Consider $f_n(x) = x^n$ on $[0, 1]$, $f(x) = 0$ for $x \in [0, 1)$, but $f(1) = 1$, so f is not continuous. This shows that $f_n \rightarrow f$ pointwise, but not uniformly.

Problem 5.10. Let $C \subseteq [0, 1]$ be the standard Cantor ternary set. Let $f_n : [0, 1] \rightarrow [0, 1]$ be the sequence of continuous, monotonically increasing functions where $f_0(x) = x$, and each f_{n+1} is constructed by modifying f_n so that it is constant on the middle thirds removed in the $(n + 1)$ -th step of the Cantor set construction.

- (a) Prove that the sequence $\{f_n\}$ converges uniformly on $[0, 1]$ to a limit function f (the Cantor function).
- (b) Prove that f is continuous and monotonically non-decreasing on $[0, 1]$.
- (c) Prove that f is surjective onto $[0, 1]$, yet is locally constant on the complement of C (an open set of measure 1).