

● HARD

● ADVANCED MATH

Advanced Math

01

10 questions · ~15 min

SAT QUESTION BANK

2026-05-29

Q1

GRID-IN ADVANCED MATH

The function g is defined by $gx = x + 14t - x$, where t is a constant. In the xy -plane, the graph of $y = gx$ passes through the point $(24, 0)$. What is the value of $g(0)$?

STUDENT-PRODUCED RESPONSE

.	/	.	.
0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

Write the answer above. Fill the matching bubble for each digit, plus the fraction bar or decimal point if needed.

$$y = x^2 + 2x + 1$$

$$x + y + 1 = 0$$

If (x_1, y_1) and (x_2, y_2) are the two solutions to the system of equations above, what is the value of $y_1 + y_2$?

- A. -3
- B. -2
- C. -1
- D. 1

Which expression is equivalent to $\frac{42a}{k} + 42ak$, where $k > 0$?

- A. $\frac{84a}{k}$
- B. $\frac{84ak^2}{k}$
- C. $\frac{42ak + 1}{k}$
- D. $\frac{42ak^2 + 1}{k}$

In the xy -plane, a parabola has vertex $(9, -14)$ and intersects the x -axis at two points. If the equation of the parabola is written in the form $y = ax^2 + bx + c$, where a , b , and c are constants, which of the following could be the value of $a + b + c$?

- A. -23
- B. -19
- C. -14
- D. -12

$$\sqrt{2x+6} + 4 = x + 3$$

What is the solution set of the equation above?

- A. $\{-1\}$
- B. $\{5\}$
- C. $\{-1, 5\}$
- D. $\{0, -1, 5\}$

The expression $4x^2 + bx - 45$, where b is a constant, can be rewritten as $hx + kx + j$, where h , k , and j are integer constants. Which of the following must be an integer?

- A. $\frac{b}{h}$
- B. $\frac{b}{k}$
- C. $\frac{45}{h}$
- D. $\frac{45}{k}$

Q7

GRID-IN ADVANCED MATH

The function f is defined by $f(x) = a^x + b$, where a and b are constants. In the xy -plane, the graph of $y = f(x)$ has an x -intercept at $(2, 0)$ and a y -intercept at $(0, -323)$. What is the value of b ?

STUDENT-PRODUCED RESPONSE

.	/	.	.
0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

Write the answer above. Fill the matching bubble for each digit, plus the fraction bar or decimal point if needed.

$$x^2 - 34x + c = 0$$

In the given equation, c is a constant. The equation has no real solutions if $c > n$. What is the least possible value of n ?

STUDENT-PRODUCED RESPONSE

.	/	.	.
0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

Write the answer above. Fill the matching bubble for each digit, plus the fraction bar or decimal point if needed.

$$0.36x^2 + 0.63x + 1.17$$

The given expression can be rewritten as $a4x^2 + 7x + 13$, where a is a constant. What is the value of a ?

STUDENT-PRODUCED RESPONSE

.	/	.	.
0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

Write the answer above. Fill the matching bubble for each digit, plus the fraction bar or decimal point if needed.

Q10

GRID-IN

ADVANCED MATH

$$h(x) = x^3 + ax^2 + bx + c$$

The function h is defined above, where a , b , and c are integer constants. If the zeros of the function are -5 , 6 , and 7 , what is the value of c ?

STUDENT-PRODUCED RESPONSE

.	/	.	.
0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

Write the answer above. Fill the matching bubble for each digit, plus the fraction bar or decimal point if needed.