

● HARD

● PROBLEM-SOLVING AND DATA ANALYSIS

● ANSWER KEY

Problem-Solving and Data Analysis

02

10 answers · Rationale-rich · Teach from doc

Q1**0.32**

Accept also: 8/25

The correct answer is .32. The volume of a cube is given by the formula $V = s^3$, where s is the length of an edge of the cube. It's given that each edge of the cube has a length of 2.00 centimeters. Substituting 2.00 centimeters for s in the formula $V = s^3$ yields $V = (2.00 \text{ centimeters})^3$, or $V = 8.00$ cubic centimeters. It's given that the cube has a mass of 2.56 grams. Dividing the mass, in grams, of the cube by the volume, in cubic centimeters, of the cube gives its density, in grams per cubic centimeters. Therefore, the density of the cube is $\frac{2.56 \text{ grams}}{8.00 \text{ cubic centimeters}}$, or .32 grams per cubic centimeter. Note that .32 and 8/25 are examples of ways to enter a correct answer.

Q2**D**

D. $d = 33t + 84$

Choice D is correct. On the line of best fit, d increases from approximately 480 to 880 between $t = 12$ and $t = 24$. The slope of the line of best fit is the difference in d -values divided by the difference in t -values, which gives $\frac{880 - 480}{24 - 12} = \frac{400}{12}$, or approximately 33. Writing the equation of the line of best fit in slope-intercept form gives $d = 33t + b$, where b is the y -coordinate of the y -intercept. This equation is satisfied by all points on the line, so $d = 480$ when $t = 12$. Thus, $480 = 33(12) + b$, which is equivalent to $480 = 396 + b$. Subtracting 396 from both sides of this equation gives $b = 84$. Therefore, an equation for the line of best fit could be $d = 33t + 84$.

Choice A is incorrect and may result from an error in calculating the slope and misidentifying the y -coordinate of the y -intercept of the graph as the value of d at $t = 10$ rather than the value of d at $t = 0$. Choice B is incorrect and may result from using the smallest value of t on the graph as the slope and misidentifying the y -coordinate of the y -intercept of the graph as the value of d at $t = 10$ rather than the value of d at $t = 0$. Choice C is incorrect and may result from misidentifying the y -coordinate of the y -intercept as the smallest value of d on the graph.

Q3

C

- C. The sampling method is flawed and may produce a biased estimate of the mean number of children per household in the community.
-

Choice C is correct. In order to use a sample mean to estimate the mean for a population, the sample must be representative of the population (for example, a simple random sample). In this case, Tabitha surveyed 20 families in a playground. Families in the playground are more likely to have children than other households in the community. Therefore, the sample isn't representative of the population. Hence, the sampling method is flawed and may produce a biased estimate.

Choices A and D are incorrect because they incorrectly assume the sampling method is unbiased. Choice B is incorrect because a sample of size 20 could be large enough to make an estimate if the sample had been representative of all the families in the community.

Q4

D

- D. Sample A had a smaller sample size.
-

Choice D is correct. Sample size is an appropriate reason for the margin of error to change. In general, a smaller sample size increases the margin of error because the sample may be less representative of the whole population.

Choice A is incorrect. The margin of error will depend on the size of the sample of recorded votes, not the number of votes that could not be recorded. In any case, the smaller number of votes that could not be recorded for sample A would tend to decrease, not increase, the comparative size of the margin of error. Choice B is incorrect. Since the percent in favor for sample A is the same distance from 50% as the percent in favor for sample B, the percent of favorable responses doesn't affect the comparative size of the margin of error for the two samples. Choice C is incorrect. If sample A had a larger margin of error than sample B, then sample A would tend to be less representative of the population. Therefore, sample A is not likely to have a larger sample size.

Q5**0.7**

Accept also: 7/10

The correct answer is $\frac{35}{50}$. Based on the table, there are a total of 50 tulips, and 35 of these tulips are from site A. The probability of selecting at random a flower from site A, given that the flower is a tulip, is equal to the number of tulips from site A divided by the total number of tulips, which can be written as $\frac{35}{50}$, or $\frac{7}{10}$. Note that 35/50, 7/10, and .7 are examples of ways to enter a correct answer.

Q6**B**B. $p = 5x$

Choice B is correct. It's given that each serving of Crunchy Grain cereal provides 5% of an adult's daily allowance of potassium, so x servings would provide x times 5%. The percentage of an adult's daily allowance of potassium, p , is 5 times the number of servings, x . Therefore, the percentage of an adult's daily allowance of potassium can be expressed as $p = 5x$.

Choices A, C, and D are incorrect and may result from incorrectly converting 5% to its decimal equivalent, which isn't necessary since p is expressed as a percentage. Additionally, choices C and D are incorrect because the context should be represented by a linear relationship, not by an exponential relationship.

Q7**A**

A. The mean of data set A is greater than the mean of data set B.

Choice A is correct. The mean value of a data set is the sum of the values of the data set divided by the number of values in the data set. When a data set is represented in a frequency table, the sum of the values in the data set is the sum of the products of each value and its frequency. For data set A, the sum of products of each value and its frequency is $302 + 344 + 385 + 427 + 469$, or 1,094. It's given that there are 27 values in data set A. Therefore, the mean of data set A is $\frac{1,094}{27}$, or approximately 40.52. Similarly, the mean of data B is $\frac{958}{27}$, or approximately 35.48. Therefore, the mean of data set A is greater than the mean of data set B.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Q8**6.21**

The correct answer is 6.21. It's given that the samples of pumice were cut in the shape of a cube. It's also given that the length of the edge of one of these cubes is 3.000 centimeters. Therefore, the volume of this cube is $3.000 \text{ centimeters}^3$, or 27 cubic centimeters. Since the density of this cube is 0.230 grams per cubic centimeter, it follows that the mass of this cube is $\frac{0.230 \text{ grams}}{1 \text{ cubic centimeter}} \cdot 27 \text{ cubic centimeters}$, or 6.21 grams.

Q9**A****A.** 0.83

Choice A is correct. It's given that an equation for the exponential model shown can be written as $y = ab^x$, where a and b are positive constants. For an exponential model written in this form, if the value of b is greater than 0 but less than 1, the model is decreasing. If the value of b is greater than 1, the model is increasing. The exponential model shown is decreasing. Therefore, the value of b is greater than 0 but less than 1. Of the given choices, only 0.83 is a value greater than 0 but less than 1. Thus, 0.83 is closest to the value of b .

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Q10**D****D.** All trivia teams in the tournament

Choice D is correct. It's given that the organizer selected 55 teams at random from all trivia teams in the tournament. A table is also given displaying the information for the 40 teams in the sample that practiced for at least 3 hours per week. Selecting a sample of a reasonable size at random to use for a survey allows the results from that survey to be applied to the population from which the sample was selected, but not beyond this population. Thus, only the sampling method information is necessary to determine the largest population to which the results of the study can be generalized. Since the organizer selected the sample at random from all trivia teams in the tournament, the largest population to which the results of the study can be generalized is all trivia teams in the tournament.

Choice A is incorrect. The sample was selected at random from all trivia teams in the tournament, not just from the teams that scored an average of 14 or more points per round.

Choice B is incorrect. If a study uses a sample selected at random from a population, the results of the study can be generalized to the population, not just the sample.

Choice C is incorrect. If a study uses a sample selected at random from a population, the results of the study can be generalized to the population, not just a subset of the sample.

Notes

Accept equivalent forms (e.g. $0.25 \equiv 1/4$). For grid-in items, decimal and fraction answers are both valid.

Generated 2026-05-29 · cb-educator-question-bank