

● HARD

● ALGEBRA

● ANSWER KEY

Linear equations in one variable

02

9 answers · Rationale-rich · Teach from doc

Q1**A****A.** Exactly one

Choice A is correct. Distributing 12 on the left-hand side and -3 on the right-hand side of the given equation yields $12x - 36 = -3x - 36$. Adding $3x$ to each side of this equation yields $15x - 36 = -36$. Adding 36 to each side of this equation yields $15x = 0$. Dividing each side of this equation by 15 yields $x = 0$. This means that 0 is the only solution to the given equation. Therefore, the given equation has exactly one solution.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Q2**-22**

The correct answer is -22. The given equation can be rewritten as $-2t + 3 = 38$. Dividing both sides of this equation by -2 yields $t + 3 = -19$. Subtracting 3 from both sides of this equation yields $t = -22$. Therefore, -22 is the value of t that is the solution to the given equation.

Q3**-.9333**Accept also: $-14/15$

The correct answer is $-\frac{14}{15}$. A linear equation in the form $ax + b = cx + d$ has no solution only when the coefficients of x on each side of the equation are equal and the constant terms are not equal. Dividing both sides of the given equation by 2 yields $kx - n = -\frac{28}{30}x - \frac{36}{38}$, or $kx - n = -\frac{14}{15}x - \frac{18}{19}$. Since it's given that the equation has no solution, the coefficient of x on both sides of this equation must be equal, and the constant terms on both sides of this equation must not be equal. Since $\frac{18}{19} < 1$, and it's given that $n > 1$, the second condition is true. Thus, k must be equal to $-\frac{14}{15}$. Note that $-14/15$, $-.9333$, and -0.933 are examples of ways to enter a correct answer.

Q4**403**

The correct answer is 403. For a linear equation in one variable to have infinitely many solutions, the coefficients of the variable must be equal on both sides of the equation and the constant terms must also be equal on both sides of the equation. The given equation can be rewritten as $\frac{43x+7}{4} - \frac{s}{13} = rx - 8$, or $3x + 7 - \frac{s}{13} = rx - 8$. Applying the distributive property to the right-hand side of this equation yields $3x + 7 - \frac{s}{13} = rx - 8r$. For this equation to have infinitely many solutions, the coefficients of x must be equal, so it follows that $3 = r$. Additionally, the constant terms must be equal, which means $7 - \frac{s}{13} = -8r$. Substituting 3 for r in this equation yields $7 - \frac{s}{13} = -83$, or $7 - \frac{s}{13} = -24$. Adding $\frac{s}{13}$ to both sides of this equation yields $7 = -24 + \frac{s}{13}$. Adding 24 to both sides of this equation yields $31 = \frac{s}{13}$. Multiplying both sides of this equation by 13 yields $403 = s$. Therefore, if the equation has infinitely many solutions, the value of s is 403.

Q5**B****B. -2 and 2**

Choice B is correct. Multiplying both sides of the given equation by 313, or 39, yields $39\frac{x+6}{3} = 39\frac{x+6}{13}$, or $13x + 6 = 3x + 6$. Subtracting $3x + 6$ from both sides of this equation yields $10x + 6 = 0$. Dividing both sides of this equation by 10 yields $x + 6 = 0$. Therefore, if $\frac{x+6}{3} = \frac{x+6}{13}$, then the value of $x + 6$ is 0. It follows that of the given choices, the value of $x + 6$ is between -2 and 2.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Q6**D**

D. $\frac{4}{25}m = 5$

Choice D is correct. Since gasoline costs \$4 per gallon, and since Alan's car travels an average of 25 miles per gallon, the expression $\frac{4}{25}$ gives the cost, in dollars per mile, to drive the car.

Multiplying $\frac{4}{25}$ by m gives the cost for Alan to drive m miles in his car. Alan wants to reduce his weekly spending by \$5, so setting $\frac{4}{25}m$ equal to 5 gives the number of miles, m , by which he must reduce his driving.

Choices A, B, and C are incorrect. Choices A and B transpose the numerator and the denominator in the fraction. The fraction $\frac{25}{4}$ would result in the unit miles per dollar, but the question requires a unit of dollars per mile. Choices A and C set the expression equal to 95 instead of 5, a mistake that may result from a misconception that Alan wants to reduce his driving by 5 miles each week; instead, the question says he wants to reduce his weekly expenditure by \$5.

Q7**.9411**

Accept also: .9412, 16/17

The correct answer is $\frac{16}{17}$. It's given that the equation $3kx + 13 = \frac{48}{17}x + 36$ has no solution. A linear equation in the form $ax + b = cx + d$, where a , b , c , and d are constants, has no solution only when the coefficients of x on each side of the equation are equal and the constant terms aren't equal. Dividing both sides of the given equation by 3 yields $kx + 13 = \frac{48}{51}x + \frac{36}{3}$, or $kx + 13 = \frac{16}{17}x + 12$. Since the coefficients of x on each side of the equation must be equal, it follows that the value of k is $\frac{16}{17}$. Note that 16/17, .9411, .9412, and 0.941 are examples of ways to enter a correct answer.

Q8**D**

D. $5n + 10 = 100$

Choice D is correct. It's given that during a certain day, the number of 9-inch frying pans the manufacturing plant makes is n and the number of 7-inch frying pans it makes is 10. It's also given that during this day the number of 10-inch frying pans that the manufacturing plant makes is 4 times the number of 9-inch frying pans, or $4n$. Therefore, the total number of 7-inch, 9-inch, and 10-inch frying pans the manufacturing plant makes is $n + 10 + 4n$, or $5n + 10$. It's given that during this day the manufacturing plant makes 100 frying pans total. Thus, the equation $5n + 10 = 100$ represents this situation.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Q9**B**

B. -3 and 3

Choice B is correct. Multiplying both sides of the given equation by $(7)(9)$, or 63, yields $\left(63\right)\frac{x-5}{7} = \left(63\right)\frac{x-5}{9}$, or $9(x-5) = 7(x-5)$. Subtracting $7(x-5)$ from both sides of this equation yields $2(x-5) = 0$. Dividing both sides of this equation by 2 yields $x-5 = 0$. Therefore, if $\frac{x-5}{7} = \frac{x-5}{9}$, then the value of $x-5$ is 0. It follows that of the given choices, the value of $x-5$ is between -3 and 3.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Choice D is incorrect and may result from conceptual or calculation errors.

Notes

Accept equivalent forms (e.g. $0.25 \equiv 1/4$). For grid-in items, decimal and fraction answers are both valid.

Generated 2026-05-29 · cb-educator-question-bank