

Muon Physics Experiment

Kevin Robb, Bill McNulty
 University of Oklahoma,
 Homer L. Dodge Department of Physics and Astronomy
 440 W. Brooks St, Norman, Oklahoma, USA 73019
 email address: kevin.rob主@ou.edu
 (Dated: November 1, 2019)

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

I. INTRODUCTION

Cosmic rays are entering the earth's upper atmosphere all the time, decaying into many particles that bombard the surface below [1]. One of these such particles is the muon, which is very abundant and simple to detect. Cosmic rays are still not fully understood, but they are protons, neutrons, or other particles with high energy coming from somewhere else in the universe. When these particles enter the earth's atmosphere, they collide with molecules and atoms in the air, undergoing a nuclear reaction or decaying as in Fig. 1.




FIG. 1. Cosmic ray decay. This figure shows a proton, the most common cosmic ray, decaying into two pions (and some other particles), which decay further into 1 muon and 1 neutrino each. Most muons reach the surface before decaying, and their decay is shown in Fig. 2.

Muons produced in this way are not electrically neutral, meaning they will interact electromagnetically with whatever material they come into contact with, losing some energy as they do so. This is how we can detect muons; muons pass through the scintillator and deposit some energy, causing ionization which triggers




FIG. 2. Muon decay. μ^+ decay looks similar with all signs reversed. Each muon produced by cosmic rays as in Fig. 1 decays into an electron and 2 neutrinos. These neutrinos whisk away some of the energy from the muon, but the rest (about a third) goes to an electron. Our apparatus contains a scintillator to detect muon events and a photo-multiplier tube that is triggered by this produced electron.

the photo-multiplier tube (PMT) and creates a measurable pulse. Similarly, a muon that loses enough energy will decay inside the scintillator to produce an electron, triggering the PMT and creating a pulse. In this experiment, we will measure the time differences between pulses and perform an analysis in order to determine the average lifetime of a muon. We will be paring out these enter-decay pulse pairs and disregarding enter-enter pulse pairs caused by two different muons. This experiment can be performed at different elevations to measure the effects of special relativity.

II. THEORY AND BACKGROUND

Muons are produced in the chain of decay from cosmic rays, shown in Fig. 1 and Fig. 2, and have a relatively long lifetime on the order of microseconds, meaning they survive long enough to make it to the surface, unlike the intermediate short-lived pions. The longer

lifetime alone does not allow them to survive to the surface, but their high speed combined with the effects of special relativity dilates their lifetime in the earth's frame of reference. This dilated time is described as

$$\Delta t = \frac{\Delta t_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}, \quad (1)$$

where t is the observed time in the earth's reference frame, t_0 is the proper time (in the muon's reference frame), v is the speed of the muon with respect to the earth, and c is the speed of light. The dilated time is longer than the proper time by a factor depending on the measured lifetime. This factor is $\exp(-t/\tau)$, obtained by putting our data in a histogram and fitting it to

$$N(t) = Be^{-(t/\tau)} + A, \quad (2)$$

where $N(t)$ is the number of counts expected in the bin at time t in the histogram of our data points, A is a vertical shifting parameter corresponding to a uniform distribution of uncorrelated events, and B is a scaling factor relating to the decaying exponential curve of correlated events. In other words, we are looking for the slope of the linear curve when the vertical axis is $\log(N(t))$ instead of $N(t)$. Finding this factor gives us $-B/\tau$, and we can solve for τ , the average lifetime of the muon [2].

After finding the muon lifetime τ , we can further calculate the Fermi coupling constant G_F with

$$\tau = \frac{192\pi^3\hbar^7}{G_F^2 m^5 c^4}, \quad (3)$$

where π , \hbar , and c are known constants and m is the muon mass which we obtain from the Particle Data Group as $105.6583745 \pm 0.0000024$ MeV [3].

III. APPARATUS AND EXPERIMENT

Nam dui ligula, fringilla a, euismod sodales, sollici-

tudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

IV. ANALYSIS AND DISCUSSION

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

V. CONCLUSIONS

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

-
- [1] David Van Baak, *A Conceptual Introduction to 'Muon Physics'* (6/18/2010).
 [2] Thomas Coan and Jingbo Ye *Muon Physics (MP1-A) User's Manual* (TeachSpin Instruction Manuals, South-

- ern Methodist University).
 [3] M. Tanabashi et al.(Particle Data Group), *Phys. Rev. D98, 030001 (2018) and 2019 update* (Lawrence Berkeley National Laboratory, created 8/2/2019).