20 Comparing Two Means

- Most interesting statistical problems involve multiple measured variables. For example, many problems involve comparing two (or more) populations or groups based on two (or more) samples.
- In such situations, each population or group will have its own parameters, and there will often be dependence between parameters. We are often interested in difference or ratios of parameters between groups.

Example 20.1. Do newborns born to mothers who smoke tend to weigh less at birth than newborns from mothers who don't smoke? We'll investigate this question using birthweight (pounds) data on a sample of full term births in North Carolina over a one year period.

Assume birthweights follow a conditional Normal distribution with mean μ_1 for nonsmokers and mean μ_2 for smokers, and standard deviation σ.

Note: our primary goal will be to compare the means μ_1 and μ_2. We're assuming a common standard deviation σ to simplify a little, but we could (and probably should) also let standard deviation vary by smoking status. (We could also assume something other than conditional Normal, but again, we're simplifying.)

1. Suggest a prior distribution.

2. The prior distribution will be a joint distribution on (μ_1, μ_2, σ) triples. We could assume a prior under which μ_1 and μ_2 are independent. But why might we want to assume some prior *dependence* between μ_1 and μ_2? (For some motivation it might help to consider what the frequentist null hypothesis would be.)

3. Regardless of your answers to the previous parts, assume for the prior: μ_1, μ_2, σ are independent; each of μ_1, μ_2 has a Normal(7.5, 0.5) distribution; σ has a Half-Normal(0, 1) distribution. How do you interpret the parameter $\mu_1 - \mu_2$? Plot the prior distribution of $\mu_1 - \mu_2$, and find prior central 50%, 80%, and 98% credible intervals. Also compute the prior probability that $\mu_1 - \mu_2 > 0$.

4. Briefly describe the sample data, summarized in Section 20.1.2.

5. Is it reasonable to assume that the two *samples* are independent? (In this case the y_1 and y_2 samples would be *conditionally independent* given (μ_1, μ_2, σ).)

6. Describe how you would compute the likelihood. For concreteness, how you would you compute the likelihood if there were only 4 babies in the sample: 2 non-smokers with birthweights of 8 pounds and 7 pounds, and 2 smokers with birthweights of 8.3 pounds and 7.1 pounds.

7. Use `brms` to approximate the posterior distribution. Are μ_1 and μ_2 independent according the posterior? Why do you think that is?

8. Plot the posterior distribution of $\mu_1 - \mu_2$ and describe it. Compute and interpret posterior central 50%, 80%, and 98% credible intervals. Also compute and interpret the posterior probability that $\mu_1 - \mu_2 > 0$.

9. If we're interested in $\mu_1 - \mu_2$, why didn't we put a prior directly on $\mu_1 - \mu_2$ rather than on (μ_1, μ_2)?

10. Plot the posterior distribution of μ_1/μ_2, describe it, and find and interpret posterior central 50%, 80%, and 98% credible intervals.

11. Is there some evidence that babies whose mothers smoke tend to weigh less than those whose mothers don't smoke?

12. Can we say that smoking is the cause of the difference in mean weights?

13. Is there some evidence that babies whose mothers smoke tend to weigh *much* less than those whose mothers don't smoke? Explain.

14. One quantity of interest is the **effect size**, which is a way of measuring the magnitude of the difference between groups. When comparing two means, a simple measure of effect size (*Cohen's d*) is

$$\frac{\mu_1 - \mu_2}{\sigma}$$

Plot the posterior distribution of this effect size and describe it. Compute and interpret posterior central 50%, 80%, and 98% credible intervals.

Independence in the data versus in the prior/posterior

It is typical to assume *independence in the data*, e.g., independence of values of the measured variables within and between samples (conditional on the parameters). Whether independence in the data is a reasonable assumption depends on how the data is collected.

But whether it is reasonable to assume prior *independence of parameters* is a completely separate question and is dependent upon our subjective beliefs about any relationships between parameters.

Transformations of parameters

The primary output of a Bayesian data analysis is the full joint posterior distribution on all parameters. Given the joint distribution, the distribution of transformations of the primary parameters is readily obtained.

Effect size for comparing means

When comparing groups, a more important question than "is there a difference?" is "*how large* is the difference?" An **effect size** is a measure of the magnitude of a difference between groups. A difference in parameters can be used to measure the absolute size of the difference in the measurement units of the variable, but effect size can also be measured as a relative difference.

When comparing a numerical variable between two groups, one measure of the population effect size is **Cohens's** d

$$\frac{\mu_1 - \mu_2}{\sigma}$$

The values of any numerical variable vary naturally from unit to unit. The SD of the numerical variable measures the degree to which individual values of the variable vary naturally, so the

SD provides a natural "scale" for the variable. Cohen's d compares the magnitude of the difference in means relative to the natural scale (SD) for the variable.

Some rough guidelines for interpreting $|d|$:

d	0.2	0.5	0.8	1.2	2.0
Effect size	Small	Medium	Large	Very Large	Huge

For example, assume the two population distributions are Normal and the two population standard deviations are equal. Then when the effect size is 1.0 the median of the distribution with the higher mean is the 84th percentile of the distribution with the lower mean, which is a very large difference.

d	0.2	0.5	0.8	1.0	1.2	2.0
Effect size	Small	Medium	Large		Very Large	Huge
Median of population 1 is (blank) percentile of population 2	58th	69th	79th	84th	89th	98th



20.1 Notes

20.1.1 Prior distribution of $\mu_1 - \mu_2$

The prior distribution of $\mu_1 - \mu_2$ is Normal with mean 7.5-7.5 = 0 and standard deviation $\sqrt{0.5^2 + 0.5^2} = 0.707$

```r
qnorm(c(0.01, 0.10, 0.25, 0.75, 0.90, 0.99),
      mean = 7.5 - 7.5,
      sd = sqrt(0.5 ^ 2 + 0.5 ^ 2))
```

```
[1] -1.6449764 -0.9061938 -0.4769363  0.4769363  0.9061938  1.6449764
```

20.1.2 Sample data

```r
data = read_csv("baby_smoke.csv")
```

```
Rows: 999 Columns: 13
-- Column specification --------------------------------------------------------
Delimiter: ","
chr (7): mature, premie, marital, lowbirthweight, gender, habit, whitemom
dbl (6): fage, mage, weeks, visits, gained, weight

i Use `spec()` to retrieve the full column specification for this data.
i Specify the column types or set `show_col_types = FALSE` to quiet this message.
```

```r
head(data)
```

```
# A tibble: 6 x 13
   fage  mage mature    weeks premie visits marital gained weight lowbirthweight
  <dbl> <dbl> <chr>     <dbl> <chr>   <dbl> <chr>    <dbl>  <dbl> <chr>
1    NA    13 younger ~    39 full ~     10 married     38   7.63 not low
2    NA    14 younger ~    42 full ~     15 married     20   7.88 not low
3    19    15 younger ~    37 full ~     11 married     38   6.63 not low
4    21    15 younger ~    41 full ~      6 married     34   8    not low
5    NA    15 younger ~    39 full ~      9 married     27   6.38 not low
6    NA    15 younger ~    38 full ~     19 married     22   5.38 low
# i 3 more variables: gender <chr>, habit <chr>, whitemom <chr>
```

We'll only include full term babies in the analysis. The variables of interest are `weight` and `habit`.

```r
data = data |>
  filter(premie == "full term")
```

habit	n()	mean(weight)	sd(weight)
nonsmoker	739	7.50	1.08
smoker	107	7.17	0.97

```
data |>
  group_by(habit) |>
  summarize(n(), mean(weight), sd(weight)) |>
  kbl(digits = 2) |>
  kable_styling()
```

```
ggplot(data,
       aes(x = weight,
           fill = habit)) +
   geom_histogram(alpha = 0.3,
                  aes(y = after_stat(density)),
                  position = 'identity')
```

```
`stat_bin()` using `bins = 30`. Pick better value `binwidth`.
```

```
ggplot(data,
       aes(x = weight,
           y = habit,
           col = habit)) +
   geom_boxplot()
```





20.1.3 Posterior distribution

```r
library(brms)
```

```
Loading required package: Rcpp

Loading 'brms' package (version 2.23.0). Useful instructions
can be found by typing help('brms'). A more detailed introduction
to the package is available through vignette('brms_overview').


Attaching package: 'brms'

The following object is masked from 'package:stats':

    ar
```

```r
library(tidybayes)
```

```
Attaching package: 'tidybayes'

The following objects are masked from 'package:brms':

    dstudent_t, pstudent_t, qstudent_t, rstudent_t
```

```r
library(bayesplot)
```

```
This is bayesplot version 1.14.0

- Online documentation and vignettes at mc-stan.org/bayesplot

- bayesplot theme set to bayesplot::theme_default()

   * Does _not_ affect other ggplot2 plots

   * See ?bayesplot_theme_set for details on theme setting
```

```
Attaching package: 'bayesplot'

The following object is masked from 'package:brms':

    rhat
```

We want a posterior for μ_1 and μ_2, so we set the intercept to 0 in the linear model. We can use `get_prior` with out model specifications to see what the parameters will be.

```r
get_prior(data = data,
          family = gaussian(),
          weight ~ 0 + habit)
```

```
                prior class           coef group resp dpar nlpar lb ub tag
               (flat)     b
               (flat)     b habitnonsmoker
               (flat)     b    habitsmoker
 student_t(3, 0, 2.5) sigma                                       0
       source
      default
 (vectorized)
 (vectorized)
      default
```

Now we fit the model. The `normal(7.5, 0.5)` prior for μ_1 and μ_2 will be vectorized.

```r
fit <- brm(data = data,
           family = gaussian(),
           weight ~ 0 + habit,
           prior = c(prior(normal(7.5, 0.5), class = b),
                     prior(normal(0, 1), class = sigma)),
           iter = 3500,
           warmup = 1000,
           chains = 4,
           refresh = 0)
```

```
Compiling Stan program...

Start sampling
```

```r
prior_summary(fit)
```

```
            prior class           coef group resp dpar nlpar lb ub tag
 normal(7.5, 0.5)     b
 normal(7.5, 0.5)     b habitnonsmoker
 normal(7.5, 0.5)     b    habitsmoker
     normal(0, 1) sigma                                       0
       source
         user
 (vectorized)
 (vectorized)
         user
```

```r
summary(fit)
```

```
 Family: gaussian
  Links: mu = identity
Formula: weight ~ 0 + habit
   Data: data (Number of observations: 846)
  Draws: 4 chains, each with iter = 3500; warmup = 1000; thin = 1;
         total post-warmup draws = 10000

Regression Coefficients:
               Estimate Est.Error l-95% CI u-95% CI Rhat Bulk_ESS Tail_ESS
habitnonsmoker     7.50      0.04     7.43     7.58 1.00     9993     7584
habitsmoker        7.19      0.10     6.99     7.38 1.00     8723     6750

Further Distributional Parameters:
      Estimate Est.Error l-95% CI u-95% CI Rhat Bulk_ESS Tail_ESS
sigma     1.07      0.03     1.02     1.12 1.00    10160     7338

Draws were sampled using sampling(NUTS). For each parameter, Bulk_ESS
and Tail_ESS are effective sample size measures, and Rhat is the potential
scale reduction factor on split chains (at convergence, Rhat = 1).
```

```r
plot(fit)
```




```
pairs(fit,
      off_diag_args = list(alpha = 0.1))
```



.chain	.iteration	.draw	mu2	mu1	sigma
1	1	1	7.34	7.49	1.09
1	2	2	7.27	7.50	1.04
1	3	3	7.24	7.51	1.06
1	4	4	7.23	7.50	1.05
1	5	5	7.32	7.44	1.09
1	6	6	7.08	7.52	1.09

.chain	.iteration	.draw	mu2	mu1	sigma	mu_diff
1	1	1	7.34	7.49	1.09	0.15
1	2	2	7.27	7.50	1.04	0.23
1	3	3	7.24	7.51	1.06	0.27
1	4	4	7.23	7.50	1.05	0.27
1	5	5	7.32	7.44	1.09	0.13
1	6	6	7.08	7.52	1.09	0.45

```
posterior = fit |>
  spread_draws(b_habitsmoker, b_habitnonsmoker, sigma) |>
  rename(mu1 = b_habitnonsmoker, mu2 = b_habitsmoker)

posterior |> head() |> kbl(digits = 2) |> kable_styling()
```

20.1.4 Posterior distribution of difference in means

```
posterior = posterior |>
  mutate(mu_diff = mu1 - mu2)

posterior |> head() |> kbl(digits = 2) |> kable_styling()
```

```
ggplot(posterior,
       aes(x = mu_diff)) +
  geom_histogram(aes(y = after_stat(density)),
                 col = bayes_col["posterior"], fill = "white", bins = 100) +
  geom_density(linewidth = 1, col = bayes_col["posterior"]) +
  labs(x = "Difference in means (pounds)") +
  theme_bw()
```




```r
mean(posterior$mu_diff)
```

```
[1] 0.315822
```

```r
sd(posterior$mu_diff)
```

```
[1] 0.1082119
```

```r
quantile(posterior$mu_diff,
         c(0.01, 0.10, 0.25, 0.5, 0.75, 0.90, 0.99))
```

```
         1%         10%         25%         50%         75%         90%         99%
0.06890393 0.17628197 0.24189140 0.31563945 0.39102622 0.45509328 0.56207534
```

```r
sum(posterior$mu_diff > 0) / length(posterior$mu_diff)
```

```
[1] 0.9989
```

20.1.5 Posterior distribution of ratio of means

.chain	.iteration	.draw	mu2	mu1	sigma	mu_diff	mu_ratio
1	1	1	7.34	7.49	1.09	0.15	1.02
1	2	2	7.27	7.50	1.04	0.23	1.03
1	3	3	7.24	7.51	1.06	0.27	1.04
1	4	4	7.23	7.50	1.05	0.27	1.04
1	5	5	7.32	7.44	1.09	0.13	1.02
1	6	6	7.08	7.52	1.09	0.45	1.06

```
posterior = posterior |>
  mutate(mu_ratio = mu1 / mu2)

posterior |> head() |> kbl(digits = 2) |> kable_styling()
```

```
ggplot(posterior,
       aes(x = mu_ratio)) +
  geom_histogram(aes(y = after_stat(density)),
                 col = bayes_col["posterior"], fill = "white", bins = 100) +
  geom_density(linewidth = 1, col = bayes_col["posterior"]) +
  labs(x = "Ratio of means") +
  theme_bw()
```



.chain	.iteration	.draw	mu2	mu1	sigma	mu_diff	mu_ratio	effect_size
1	1	1	7.34	7.49	1.09	0.15	1.02	0.13
1	2	2	7.27	7.50	1.04	0.23	1.03	0.22
1	3	3	7.24	7.51	1.06	0.27	1.04	0.25
1	4	4	7.23	7.50	1.05	0.27	1.04	0.26
1	5	5	7.32	7.44	1.09	0.13	1.02	0.11
1	6	6	7.08	7.52	1.09	0.45	1.06	0.41

```
mean(posterior$mu_ratio)
```

```
[1] 1.044161
```

```
sd(posterior$mu_ratio)
```

```
[1] 0.01564791
```

```
quantile(posterior$mu_ratio,
         c(0.01, 0.10, 0.25, 0.5, 0.75, 0.90, 0.99))
```

```
      1%      10%      25%      50%      75%      90%      99%
1.009344 1.024155 1.033327 1.043923 1.054902 1.064448 1.080669
```

20.1.6 Posterior distribution of effect size

```
posterior = posterior |>
  mutate(effect_size = (mu1 - mu2) / sigma)

posterior |> head() |> kbl(digits = 2) |> kable_styling()
```

```
ggplot(posterior,
       aes(x = effect_size)) +
  geom_histogram(aes(y = after_stat(density)),
                 col = bayes_col["posterior"], fill = "white", bins = 100) +
  geom_density(linewidth = 1, col = bayes_col["posterior"]) +
  labs(x = "Effect size") +
  theme_bw()
```




```r
mean(posterior$effect_size)
```

```
[1] 0.2952176
```

```r
sd(posterior$effect_size)
```

```
[1] 0.1015268
```

```r
quantile(posterior$effect_size,
         c(0.01, 0.10, 0.25, 0.5, 0.75, 0.90, 0.99))
```

```
        1%        10%        25%        50%        75%        90%        99%
0.06374755 0.16458943 0.22577629 0.29451560 0.36465077 0.42560646 0.52678586
```