

# HEPP-CPV-project

John Ronayne (), Kevin Maguire (10318135), Sinead Hales(), Dudley Grant (10275291)  
(Dated: November 6, 2013)

## I. INTRODUCTION

Short intro here

## II. $\hat{P}$ , $\hat{C}$ AND $\hat{C}\hat{P}\hat{T}$

## III. CP VIOLATION

CP violation was first observed in the mixing of neutral K-mesons by Christenson, Cronin, Fitch and Turlay in 1964 [1]. They observed the  $\hat{C}\hat{P} = -1$  state  $K_L^0$  decaying to 2 pions, a state with  $\hat{C}\hat{P} = 1$ . Although the fraction of  $K_L^0$  decays violating  $\hat{C}\hat{P}$  in this way is tiny, the discovery was significant.

## IV. KAON MIXING

### A. Neutral Kaon Mixing and $\hat{C}\hat{P}V$

As mentioned  $\hat{C}\hat{P}V$  was first observed in the neutral kaon system. Both direct and indirect  $\hat{C}\hat{P}V$  have been observed but is almost entirely dominated by the indirect process [Zeng, need better reference]. Essential to these mechanisms is the mixing between the neutral Kaon and its anti-particle corresponding to the states  $\langle K^0|$  and  $|\bar{K}^0\rangle$ . These have quark compositions of  $d\bar{s}$  and  $s\bar{d}$ , respectively.

## V. CKM MECHANISM

### Appendix A: Appendix

Difficult calculations in here.

---

[1] "Evidence for the  $2\pi$  Decay of the  $K_2^0$  Meson" - J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay (1964) Phys. Review letters, vol. 13, issue 4