

# HEPP-CPV-project

John Ronayne (10318997), Kevin Maguire (10318135), Sinead Hales(10318221), Dudley Grant (10275291)  
(Dated: November 17, 2013)

## I. INTRODUCTION

Short intro here

## II. $\hat{P}$ , $\hat{C}$ AND $\hat{C}\hat{P}\hat{T}$

## III. CP VIOLATION

CP violation was first observed in the mixing of neutral K-mesons by Christenson, Cronin, Fitch and Turlay in 1964 [1]. They observed the  $\hat{C}\hat{P} = -1$  state  $K_L^0$  decaying to 2 pions, a state with  $\hat{C}\hat{P} = 1$ . Although the fraction of  $K_L^0$  decays violating  $\hat{C}\hat{P}$  in this way is tiny, the discovery was significant.

## IV. CPV IN D-MESON SYSTEM

The quark constituents of the  $D^0(1865)$  and  $\bar{D}^0(1865)$  mesons are  $(c\bar{u})$  and  $(u\bar{c})$ , respectively. This system is unique as it is the only system which undergoes mixing and contains an up-type quark. As opposed to the  $K^0, B^0$  and  $B_S$ , which contain down quarks. This results in different quarks in the mixing box diagrams of these processes, which are illustrated in Fig.(??) and (1). The rates for  $D^0$  mixing are expected to be very small as the mixing process shown is suppressed in two ways. If the intermediate quark is a b, then the decay is doubly Cabbibo suppressed[explain? or has it been explained already?], while if the quark is a d or an s then the process is GIM suppressed[Reference John]. Other processes which may not have the same degree of suppression have been proposed, but there are large uncertainties in the theoretical calculations of their decay rates [14].




FIG. 1: Feynmann diagram showing the process by which the two  $D^0$  states mix. This process is the only known mixing process which contain the  $d,s,b$  quarks in this position [13]

The first stage in detecting CPV in any system is to find mixing between a particle and its anti-particle. Clear evidence for mixing between these states was announced in 2007 and published in 2008 by the BaBar collaboration, followed shortly by the Belle collaboration [11][12]. Results from both experiments show a small amount of  $D^0$  mixing with  $3.9 \sigma$  certainty, at a level of 1%, which is consistent with SM predictions in the order of  $\times 10^{-2}$  [14]. However, measured CP violating parameters were consistent with zero, and thus with no CPV. We discuss here the more recent methods and measurements for mixing parameters.

As in the case of Kaon and B-meson mixing we define the CP eigenstates of the D-meson to be linear combinations of flavour eigenstates.

$$|D_{1,2}^0\rangle = p |D^0\rangle \pm q |\bar{D}^0\rangle$$

Where for normalization  $|p|^2 + |q|^2 = 1$ . The  $|D_1\rangle$  state is a CP even state while the  $|D_2\rangle$  state is a CP odd state. As expected, we will see CPV if  $|p| \neq |q|$

The two parameters which determine the amount of mixing in a system are defined as:

$$x = \frac{\Delta M}{\Gamma} \qquad y = \frac{\Delta \Gamma}{2\Gamma}$$

where  $M$  is mass and  $\Gamma$  is lifetime and  $\Delta A$  :




FIG. 2:

## Appendix A: Appendix

Difficult calculations in here.

- 
- [1] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay (1964) Phys. Review letters, vol. 13, issue 4 - "Evidence for the  $2\pi$  Decay of the  $K_2^0$  Meson"
  - [2] J. Beringer et al. (Particle Data Group), Phys. Rev. D86, 010001 (2012) and 2013 partial update for the 2014 edition - "2013 Review of Particle Physics"
  - [3] B.R Martin, G. Shaw, Wiley(2008) - "Particle Physics, Third Edition"
  - [4] Tatsuya Nakada arXiv:hep-ph/9312290v1 15 Dec 1993 - "Review on CP Violation"
  - [5] <http://large.stanford.edu/courses/2008/ph204/coleman1/>
  - [6] KTeV Collaboration, arXiv:hep-ex/0208007v1 6 Aug 2002 - "Measurements of Direct CP Violation, CPT Symmetry, and Other Parameters in the Neutral Kaon System"
  - [7] Donald H. Perkins, Cambridge University Press, 4th edition - "Introduction to High Energy Physics"
  - [8] Giancarlo D'Ambrosio and Gino Isidori, arXiv:hep-ph/9611284v1 8 Nov 1996 - "CP violation in Kaon Decays"
  - [9] Gjesdal, S. et al. Phys.Lett. B52 (1974) 113 Print-74-1358 (CERN) - "A Measurement of the K(L)-K(s) Mass Difference from the Charge Asymmetry in Semileptonic Kaon Decays"
  - [10] [http://www.hep.phy.cam.ac.uk/~thomson/partIIparticles/handouts/Handout\\_12\\_2011.pdf](http://www.hep.phy.cam.ac.uk/~thomson/partIIparticles/handouts/Handout_12_2011.pdf)
  - [11] The BaBar Collaboration <http://arxiv.org/abs/hep-ex/0703020v1> 1 Apr 2007 - "Evidence for  $D^0 \bar{D}^0$  Mixing"
  - [12] The Belle Collaboration arXiv:hep-ex/0703036v2 1 Apr 2007 - "Evidence for  $D^0 \bar{D}^0$  Mixing"
  - [13] <http://inspirehep.net/record/1209003/files/mix-rick.png>
  - [14] Giulia Casarosa for the BABAR Collaboration - "Studies of CP Violation and Mixing in the D Mesons decays from BABAR" - Journal of Physics: Conference Series 335 (2011) 012043 doi:10.1088/1742-6596/335/1/012043
  - [15] François Goffinet, CP3 Seminar, Unité de Physique Théorique et Mathématique, U.C.L., December 2003
  - [16] Yoav Achiman, "Spontaneous CP Violation in SUSY," Physics Letters B, March 2007
  - [17] Peter Szekeres, "A Course in Modern Mathematical Physics," Cambridge University Press, 2004
  - [18] John Baez and John Huerta, "The Algebra of Grand Unified Theories," Bulletin of the American Mathematical Society, vol 47, May 4 2010

- [19] Howard E. Haber and Ze'ev Surujon, "Group-theoretic Condition for Spontaneous CP Violation," Physical Review D, Volume 86, Issue 7, October 2012
- [20] Bernard de Wit and Eric Laenen, "Field Theory in Particle Physics" Lecture Notes, Universiteit Utrecht, 2009 <http://www.staff.science.uu.nl/wit00103/ftip/Ch11.pdf>
- [21] Paul H. Frampton and Masayasu Harada, "Kaon Spontaneous CP Violation Reevaluated," Physical Review D, Volume 59, Issue 1, March 1998