

Kevin Maguire

April 23, 2014

Layout



Layout

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- \bigcirc $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Condition

-Layout

Layout

Strange Minkrosokian Line Flament

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- \bigcirc $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Condition

20 10 100

Layout

Layout

Singular Lorentz transformation

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions

20 10 100

Layout

Layout

Strange Minkowskian Line Element

SL(2, C) Matrices of the Lorentz Transformation

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- **3** $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions

2014 04 23

Layout

Layout

Strange Minkowskian Line Element

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- **3** $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- **5** The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions

2017 04 23

Layout

Layout

Strange Minkowskian Line Element

SL(2, C) Matrices of the Lorentz Transformation

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- **3** $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- **5** The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions

2014 04 23

-Layout

Layout

Strange Minkowskian Line Element

SL(2, C) Matrices of the Lorentz Transformation

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- **3** $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- **5** The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions



- -Proper is det 1 . preserves the orientation of spacial axes, preserves handedness
- $\bullet\,$ –orthochronous means time is always positive and the direction of time is preserved
- ullet -Think of the standard Lorentz transformation, always two null directions at $x\pm t$

 A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2+y'^2+z'^2-t'^2=x^2+y^2+z^2-t^2$$
,

in the transformation

$$(x,y,z,t) \rightarrow (x',y',z',t')$$

- Take the Proper Orthochronous Lorentz Transformations(POLTs) which form the restricted Lorentz group SO⁺(1,3)
- In general Lorentz transformations



- troduction: Lorentz Transformations by the preservation of the quadratic which form the restricted Lorentz
- Proper is det 1. preserves the orientation of spacial axes, preserves handedness
- –orthochronous means time is always positive and the direction of time is preserved
- Think of the standard Lorentz transformation, always two null directions at $x \pm t$

 A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2+y'^2+z'^2-t'^2=x^2+y^2+z^2-t^2$$
,

in the transformation

$$(x,y,z,t) \rightarrow (x',y',z',t')$$

- Take the Proper Orthochronous Lorentz Transformations(POLTs) which form the restricted Lorentz group $SO^{+}(1,3)$
- In general Lorentz transformations




- -Proper is det 1 . preserves the orientation of spacial axes, preserves handedness
- -orthochronous means time is always positive and the direction of time is preserved
- ullet Think of the standard Lorentz transformation, always two null directions at $x\pm t$

 A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2+y'^2+z'^2-t'^2=x^2+y^2+z^2-t^2$$
,

in the transformation $(x, y, z, t) \rightarrow (x', y', z', t')$

- Take the Proper Orthochronous Lorentz Transformations(POLTs) which form the restricted Lorentz group $SO^+(1,3)$
- In general Lorentz transformations have two invariant null directions



- -Proper is det 1 . preserves the orientation of spacial axes, preserves handedness
- -orthochronous means time is always positive and the direction of time is preserved
- ullet -Think of the standard Lorentz transformation, always two null directions at $x\pm t$


Introduction: Lorentz Transformations


 A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2+y'^2+z'^2-t'^2=x^2+y^2+z^2-t^2$$
,

in the transformation $(x, y, z, t) \rightarrow (x', y', z', t')$

- Take the Proper Orthochronous Lorentz Transformations(POLTs) which form the restricted Lorentz group $SO^+(1,3)$
- In general Lorentz transformations have two invariant null directions





- -Proper is det 1 . preserves the orientation of spacial axes, preserves handedness
- -orthochronous means time is always positive and the direction of time is preserved
- ullet -Think of the standard Lorentz transformation, always two null directions at $x\pm t$


Introduction: Lorentz Transformations

 A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2+y'^2+z'^2-t'^2=x^2+y^2+z^2-t^2$$
,

in the transformation $(x, y, z, t) \rightarrow (x', y', z', t')$

- Take the Proper Orthochronous Lorentz Transformations(POLTs) which form the restricted Lorentz group $SO^+(1,3)$
- In general Lorentz transformations have two invariant null directions



-Layout



Layout

- derive a strange Minkowskian line element
- making a complicated transformation that keeps a single null geodesic fixed look trivial
- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- \bigcirc $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions



- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- \bullet These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coordinate transforms that this is Kasner, but it wont be done here

Start with the Schwarzschild solution

$$\epsilon \mathrm{d}s^2 = \left(1 - \frac{2m}{r}\right)^{-1} \mathrm{d}r^2 + r^2 (\mathrm{d}\theta^2 + \sin^2\theta \mathrm{d}\phi^2) - \left(1 - \frac{2m}{r}\right) \mathrm{d}t^2.$$

• Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

• Make further coordinate transformations to obtain

$$\epsilon ds^{2} = \frac{r^{2}}{\cosh^{2} \mu \xi} (d\xi^{2} + d\eta^{2}) - 2dudr - \left(\mu^{2} - \frac{2k}{r}\right) du^{2}$$

ullet Taking the limit as the energy, $\mu o 0$ gives The Kasner Solution

$$\epsilon ds^2 = r^2 (d\xi^2 + d\eta^2) - 2dudr - \frac{2k}{r} du^2$$





- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coordinate transforms that this is Kasner, but it wont be done here

Strange Minkowskian Line Element

Start with the Schwarzschild solution

$$\epsilon \mathrm{d} s^2 = \left(1 - \frac{2m}{r}\right)^{-1} \mathrm{d} r^2 + r^2 (\mathrm{d} \theta^2 + \sin^2 \theta \mathrm{d} \phi^2) - \left(1 - \frac{2m}{r}\right) \mathrm{d} t^2.$$

• Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

Make further coordinate transformations to obtain

$$\epsilon ds^{2} = \frac{r^{2}}{\cosh^{2} \mu \xi} (d\xi^{2} + d\eta^{2}) - 2dudr - \left(\mu^{2} - \frac{2k}{r}\right) du^{2}$$

ullet Taking the limit as the energy, $\mu o 0$ gives The Kasner Solution

$$\epsilon ds^2 = r^2 (d\xi^2 + d\eta^2) - 2dudr - \frac{2k}{r} du^2$$





- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coordinate transforms that this is Kasner, but it wont be done here

Start with the Schwarzschild solution

$$\epsilon \mathrm{d} s^2 = \left(1 - \frac{2m}{r}\right)^{-1} \mathrm{d} r^2 + r^2 (\mathrm{d} \theta^2 + \sin^2 \theta \mathrm{d} \phi^2) - \left(1 - \frac{2m}{r}\right) \mathrm{d} t^2.$$

• Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

Make further coordinate transformations to obtain

$$\epsilon \mathrm{d}s^2 = \frac{r^2}{\cosh^2 \mu \xi} (\mathrm{d}\xi^2 + \mathrm{d}\eta^2) - 2\mathrm{d}u\mathrm{d}r - \left(\mu^2 - \frac{2k}{r}\right) \mathrm{d}u^2.$$

ullet Taking the limit as the energy, $\mu \to 0$ gives The Kasner Solution

$$\epsilon ds^2 = r^2 (d\xi^2 + d\eta^2) - 2dudr - \frac{2k}{r} du^2.$$





- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coordinate transforms that this is Kasner, but it wont be done here

Strange Minkowskian Line Element

Start with the Schwarzschild solution

$$\epsilon \mathrm{d} s^2 = \left(1 - \frac{2m}{r}\right)^{-1} \mathrm{d} r^2 + r^2 (\mathrm{d} \theta^2 + \sin^2 \theta \mathrm{d} \phi^2) - \left(1 - \frac{2m}{r}\right) \mathrm{d} t^2.$$

• Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

Make further coordinate transformations to obtain

$$\epsilon \mathrm{d}s^2 = \frac{r^2}{\cosh^2 \mu \xi} (\mathrm{d}\xi^2 + \mathrm{d}\eta^2) - 2\mathrm{d}u\mathrm{d}r - \left(\mu^2 - \frac{2k}{r}\right) \mathrm{d}u^2.$$

ullet Taking the limit as the energy, $\mu o 0$ gives The Kasner Solution

$$\epsilon ds^2 = r^2 (d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$





- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coordinate transforms that this is Kasner, but it wont be done here

Strange Minkowskian Line Element

Start with the Schwarzschild solution

$$\epsilon \mathrm{d} s^2 = \left(1 - \frac{2m}{r}\right)^{-1} \mathrm{d} r^2 + r^2 (\mathrm{d} \theta^2 + \sin^2 \theta \mathrm{d} \phi^2) - \left(1 - \frac{2m}{r}\right) \mathrm{d} t^2.$$

• Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

Make further coordinate transformations to obtain

$$\epsilon \mathrm{d}s^2 = \frac{r^2}{\cosh^2 \mu \xi} (\mathrm{d}\xi^2 + \mathrm{d}\eta^2) - 2\mathrm{d}u\mathrm{d}r - \left(\mu^2 - \frac{2k}{r}\right) \mathrm{d}u^2.$$

ullet Taking the limit as the energy, $\mu o 0$ gives The Kasner Solution

$$\epsilon ds^2 = r^2 (d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$



- Its easily shown with suitable coordinate transforms that this is Minkowskian line element
- This is best shown by calculating the geodesic equations after the Eddington-Finkelstein coordinate transforms, all zero if u is proper time along the geodesic

• Then with m = 0 the strange Minkowskian line element is obtained

$$\epsilon ds^2 = r^2 (d\xi^2 + d\eta^2) - 2dudr.$$

• Set r = 0 to give

$$\epsilon ds^2 = 0$$

and this is shown to be a null geodesic.

- Its easily shown with suitable coordinate transforms that this is Minkowskian line element
- This is best shown by calculating the geodesic equations after the Eddington-Finkelstein coordinate transforms, all zero if *u* is proper time along the geodesic

• Then with m = 0 the strange Minkowskian line element is obtained

$$\epsilon ds^2 = r^2 (d\xi^2 + d\eta^2) - 2dudr.$$

• Set r = 0 to give

$$\epsilon \mathrm{d}s^2 = 0$$
.

and this is shown to be a null geodesic.

- Its easily shown with suitable coordinate transforms that this is Minkowskian line element
- This is best shown by calculating the geodesic equations after the Eddington-Finkelstein coordinate transforms, all zero if *u* is proper time along the geodesic

• Then with m = 0 the strange Minkowskian line element is obtained

$$\epsilon ds^2 = r^2 (d\xi^2 + d\eta^2) - 2dudr.$$

• Set r = 0 to give

$$\epsilon \mathrm{d}s^2 = 0$$
.

and this is shown to be a null geodesic.

Layout

2014 04 23

3 Introduction: Lorentz Transformations 3 Strange Minkowskian Line Element 3 Singular Lorentz transformation

LTs that leave one null invariant direction are constructed

Layout

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- \bigcirc $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions



Singular Lorentz Transformation

- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in Cartesian
- Again its clear that r = 0 keeps one direction fixed, as then z=t
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular light transformations.

Singular Lorentz Transformation

• Define an arbitrary complex parameter $\zeta := \xi + i \eta,$ to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2dudr.$$

- The transformation $\zeta \to \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic r = 0 invariant.
- In Cartesian coordinates this transformation becomes

$$x' + iy' = x + iy + w(t - z),$$

$$z' - t' = -r = z - t,$$

$$z' + t' = z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z)$$



Singular Lorentz Transformation

| ular Lorentz Transformation | | |
|---|--|--|
| | | |
| efine an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line ement[3] | | |
| $\epsilon ds^2 = r^2 d\zeta d\overline{\zeta} - 2 ds dr.$ | | |
| he transformation $\zeta \to \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the ngle null geodesic $r = 0$ invariant. | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in Cartesian
- Again its clear that r = 0 keeps one direction fixed, as then z=t
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular light transformations.

Singular Lorentz Transformation

• Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\overline{\zeta} - 2dudr.$$

- The transformation $\zeta \to \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic r = 0 invariant.
- In Cartesian coordinates this transformation becomes

$$x' + iy' = x + iy + w(t - z),$$

 $z' - t' = -r = z - t,$
 $z' + t' = z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z)$



—Singular Lorentz Transformation



- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in Cartesian
- Again its clear that r = 0 keeps one direction fixed, as then z=t
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular largest transformations.

Singular Lorentz Transformation

• Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\overline{\zeta} - 2dudr.$$

- The transformation $\zeta \to \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic r = 0 invariant.
- In Cartesian coordinates this transformation becomes

$$x' + iy' = x + iy + w(t - z),$$

 $z' - t' = -r = z - t,$
 $z' + t' = z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).$



Singular Lorentz Transformation



- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in Cartesian
- Again its clear that r = 0 keeps one direction fixed, as then z=t
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular Lorentz transformations

Singular Lorentz Transformation

• Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\overline{\zeta} - 2dudr.$$

- The transformation $\zeta \to \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic r = 0 invariant.
- In Cartesian coordinates this transformation becomes

$$x' + iy' = x + iy + w(t - z),$$

 $z' - t' = -r = z - t,$
 $z' + t' = z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).$



Singular Lorentz Transformation



- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in Cartesian
- Again its clear that r = 0 keeps one direction fixed, as then z=t
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular Lorentz transformations

Singular Lorentz Transformation

• Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\overline{\zeta} - 2dudr.$$

- The transformation $\zeta \to \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic r = 0 invariant.
- In Cartesian coordinates this transformation becomes

$$x' + iy' = x + iy + w(t - z),$$

 $z' - t' = -r = z - t,$
 $z' + t' = z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).$



-Layout



• shown here that there is a 2 to 1 correspondence between SL(2,C) and POLTs

Layout

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- **9** $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Condition

| $-SL(2,\mathbb{C})$ | Matrices of | the POLT |
|---------------------|-------------|----------|



- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \left(\begin{array}{cc} t - z & x + iy \\ x - iy & t + z \end{array}\right)$$

• This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

• Construct the transformation $A(\vec{x}') = UA(\vec{x})U^{\dagger}$, where

$$U = \left(\begin{array}{cc} \alpha & \beta \\ \gamma & \delta \end{array}\right),$$

is an element of $\mathit{SL}(2,\mathbb{C})$



| entz Transformations and Pure Radiation Fields | - C - C - C - C - C - C - C - C - C - C |
|--|--|
| entz Transionnations and Fure Tradiation Friends | There is a one to one correspondence between points in Mi space-time and Hermitian matrices Construct the following matrix |
| $^{-}$ $SL(2,\mathbb{C})$ Matrices of the POLT | $A = \left(\begin{array}{cc} t-z & x+iy \\ x-iy & t+z \end{array} \right),$ |
| | |
| | |

- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \left(\begin{array}{cc} t - z & x + iy \\ x - iy & t + z \end{array}\right),\,$$

• This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2$$

• Construct the transformation $A(\vec{x}') = UA(\vec{x})U^{\dagger}$, where

$$U = \left(\begin{array}{cc} \alpha & \beta \\ \gamma & \delta \end{array}\right),$$

is an element of $\mathit{SL}(2,\mathbb{C})$





- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

$SL(2,\mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \left(\begin{array}{cc} t - z & x + iy \\ x - iy & t + z \end{array}\right),\,$$

• This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

• Construct the transformation $A(\vec{x}') = UA(\vec{x})U^{\dagger}$, where

$$U = \left(\begin{array}{cc} \alpha & \beta \\ \gamma & \delta \end{array}\right),$$

is an element of $SL(2,\mathbb{C})$





- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

$SL(2,\mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \left(\begin{array}{cc} t - z & x + iy \\ x - iy & t + z \end{array}\right),\,$$

• This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

• Construct the transformation $A(\vec{x}') = UA(\vec{x})U^{\dagger}$, where

$$U = \left(\begin{array}{cc} \alpha & \beta \\ \gamma & \delta \end{array}\right),$$

is an element of $SL(2,\mathbb{C})$





- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

$SL(2,\mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \left(\begin{array}{cc} t - z & x + iy \\ x - iy & t + z \end{array}\right),\,$$

• This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

• Construct the transformation $A(\vec{x}') = UA(\vec{x})U^{\dagger}$, where

$$U = \left(\begin{array}{cc} \alpha & \beta \\ \gamma & \delta \end{array}\right),$$

is an element of $SL(2,\mathbb{C})$



- This is because the determinant of U is 1
- I want to show you an example calculation of U, to do this we write in component form

$SL(2,\mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$ and $A(\vec{x})$ have the same determinant so the above transformation preserves the Lorentz quadratic form, thus it is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t'-z' & x'+iy' \\ x'-iy' & t'+z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t-z & x+iy \\ x-iy & t+z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix}$$

$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t-z)\bar{\alpha} + (x+iy)\bar{\beta} & (t-z)\bar{\gamma} + (x+iy)\bar{\delta} \\ (x-iy)\bar{\alpha} + (t+z)\bar{\beta} & (x-iy)\bar{\gamma} + (t+z)\bar{\delta} \end{pmatrix}.$$

• Thus the general relations

$$t' - z' = (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta},$$

$$x' + iy' = (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta},$$

$$t' + z' = (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}.$$

- This is because the determinant of U is 1
- I want to show you an example calculation of U, to do this we write in component form

$SL(2,\mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$ and $A(\vec{x})$ have the same determinant so the above transformation preserves the Lorentz quadratic form, thus it is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t'-z' & x'+iy' \\ x'-iy' & t'+z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t-z & x+iy \\ x-iy & t+z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$

$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t-z)\bar{\alpha} + (x+iy)\bar{\beta} & (t-z)\bar{\gamma} + (x+iy)\bar{\delta} \\ (x-iy)\bar{\alpha} + (t+z)\bar{\beta} & (x-iy)\bar{\gamma} + (t+z)\bar{\delta} \end{pmatrix}.$$

Thus the general relations

$$t' - z' = (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta},$$

$$x' + iy' = (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta},$$

$$t' + z' = (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}.$$



 $^ SL(2,\mathbb{C})$ Matrices of the POLT



- This is because the determinant of U is 1
- I want to show you an example calculation of U, to do this we write in component form

$SL(2,\mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$ and $A(\vec{x})$ have the same determinant so the above transformation preserves the Lorentz quadratic form, thus it is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t'-z' & x'+iy' \\ x'-iy' & t'+z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t-z & x+iy \\ x-iy & t+z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$

$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t-z)\bar{\alpha} + (x+iy)\bar{\beta} & (t-z)\bar{\gamma} + (x+iy)\bar{\delta} \\ (x-iy)\bar{\alpha} + (t+z)\bar{\beta} & (x-iy)\bar{\gamma} + (t+z)\bar{\delta} \end{pmatrix}.$$

• Thus the general relations

$$t' - z' = (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta},$$

$$x' + iy' = (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta},$$

$$t' + z' = (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}.$$



Example: Singular Lementz transformation $This is singular Lementz transformation from under <math display="block"> \frac{d-d}{d-d} = t-x, \qquad \text{ if } t-d = t-d =$

- Where we have also used det(U) = 1
- because the sign doesn't matter is still a solution Eqn(27)
- A,A' are points in Minkowskian space time, and $\pm U$ are POLTs

Example: Singular Lorentz transformation

• Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

 $x' + iy' = x + iy + w(t - z),$
 $t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$

• Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \qquad \beta = 0$$
 $\gamma = \bar{w}\alpha, \qquad \delta = 0$

• So there are always two possible choices of U

$$U=\pm \left(egin{array}{cc} 1 & 0 \ ar{w} & 1 \end{array}
ight)$$

ullet Thus there is a 2 to 1 correspondence between elements of $SL(2,\mathbb{C})$ and POLTs

20.4

Example: Singular Lorentz transformation

Example. Singular Lorentz transformation \mathbf{v} . This this single Lorentz restrictions from earlier $(-1-\epsilon)$, $(-1-\epsilon)$, (-1-

- ullet Where we have also used det(U)=1
- because the sign doesn't matter is still a solution Eqn(27)
- A,A' are points in Minkowskian space time, and $\pm U$ are POLTs

Example: Singular Lorentz transformation

• Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

 $x' + iy' = x + iy + w(t - z),$
 $t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$

• Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \qquad \beta = 0,$$
 $\gamma = \bar{w}\alpha, \qquad \delta = \alpha.$

• So there are always two possible choices of U

$$U=\pm \left(egin{array}{cc} 1 & 0 \ ar{w} & 1 \end{array}
ight)$$

• Thus there is a 2 to 1 correspondence between elements of $SL(2,\mathbb{C})$ and POLTs

20 10 1100

Example: Singular Lorentz transformation

Example: Singular Lorentz transformation $v \text{ Take the singular largest transformation from eater } f = e^{-x} v = e^{-x}, \qquad e^{-x} e^{-x} = e^{-x}, \qquad e^{-x} e^{-x} e^{-x} = e^{-x} e^$

- Where we have also used det(U) = 1
- because the sign doesn't matter is still a solution Eqn(27)
- A,A' are points in Minkowskian space time, and $\pm U$ are POLTs

Example: Singular Lorentz transformation

• Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

 $x' + iy' = x + iy + w(t - z),$
 $t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$

• Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \qquad \beta = 0,$$
 $\gamma = \bar{\mathbf{w}}\alpha, \qquad \delta = \alpha.$

• So there are always two possible choices of U

$$U=\pm\left(egin{array}{cc} 1 & 0 \ ar{w} & 1 \end{array}
ight)$$

• Thus there is a 2 to 1 correspondence between elements of $SL(2,\mathbb{C})$ and POLTs

20 10

Example: Singular Lorentz transformation

- Where we have also used det(U) = 1
- because the sign doesn't matter is still a solution Eqn(27)
- A,A' are points in Minkowskian space time, and $\pm U$ are POLTs

Example: Singular Lorentz transformation

• Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

 $x' + iy' = x + iy + w(t - z),$
 $t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$

• Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \qquad \beta = 0,$$
 $\gamma = \bar{\mathbf{w}}\alpha, \qquad \delta = \alpha.$

• So there are always two possible choices of U

$$U=\pm\left(egin{array}{cc} 1 & 0 \ ar{w} & 1 \end{array}
ight)$$

ullet Thus there is a 2 to 1 correspondence between elements of $SL(2,\mathbb{C})$ and POLTs

20 10

Example: Singular Lorentz transformation

- Where we have also used det(U) = 1
- because the sign doesn't matter is still a solution Eqn(27)
- A,A' are points in Minkowskian space time, and $\pm U$ are POLTs

Example: Singular Lorentz transformation

• Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

 $x' + iy' = x + iy + w(t - z),$
 $t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$


• Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \qquad \beta = 0,$$
 $\gamma = \bar{\mathbf{w}}\alpha, \qquad \delta = \alpha.$

• So there are always two possible choices of U

$$U=\pm\left(egin{array}{cc} 1 & 0 \ ar{w} & 1 \end{array}
ight)$$

ullet Thus there is a 2 to 1 correspondence between elements of $SL(2,\mathbb{C})$ and POLTs



Layout

Layout

Layout


• Connect Minkowskian space to the 2-sphere by stereographic projection, so we can use points on a 2 sphere to think about LTs

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- \bullet $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- **5** The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions



- As we know, stereographic projection doesn't map the point N at the top of the circle, so that's why we map N to infinity and need to consider the extended complex plane
- It can also be written in terms of θ and ϕ

• Use Stereographic Projection to map \mathbb{S}^2 to the extended complex plane, $\hat{\mathbb{C}}=\mathbb{C}\cup\{\infty\}$




• The algebraic relation for a unit vector is

$$(x,y,z) = \left(\frac{\overline{\zeta} + \zeta}{\overline{\zeta}\zeta + 1}, i\frac{\overline{\zeta} - \zeta}{\overline{\zeta}\zeta + 1}, \frac{\overline{\zeta}\zeta - 1}{\overline{\zeta}\zeta + 1}\right)$$

ullet There is a one to one correspondence between \mathbb{S}^2 and $\hat{\mathbb{C}}$






- As we know, stereographic projection doesn't map the point N at the top of the circle, so that's why we map N to infinity and need to consider the extended complex plane
- It can also be written in terms of θ and ϕ

Fractional Linear Transformation: Stereographic Projection

• Use Stereographic Projection to map \mathbb{S}^2 to the extended complex plane, $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$



• The algebraic relation for a unit vector is

$$(x,y,z) = \left(\frac{\overline{\zeta} + \zeta}{\overline{\zeta}\zeta + 1}, i\frac{\overline{\zeta} - \zeta}{\overline{\zeta}\zeta + 1}, \frac{\overline{\zeta}\zeta - 1}{\overline{\zeta}\zeta + 1}\right)$$

• There is a one to one correspondence between \mathbb{S}^2 and $\hat{\mathbb{C}}$






- As we know, stereographic projection doesn't map the point N at the top of the circle, so that's why we map N to infinity and need to consider the extended complex plane
- It can also be written in terms of θ and ϕ

Fractional Linear Transformation: Stereographic Projection

• Use Stereographic Projection to map \mathbb{S}^2 to the extended complex plane, $\hat{\mathbb{C}}=\mathbb{C}\cup\{\infty\}$



• The algebraic relation for a unit vector is

$$(x,y,z) = \left(\frac{\overline{\zeta} + \zeta}{\overline{\zeta}\zeta + 1}, i\frac{\overline{\zeta} - \zeta}{\overline{\zeta}\zeta + 1}, \frac{\overline{\zeta}\zeta - 1}{\overline{\zeta}\zeta + 1}\right),\,$$

ullet There is a one to one correspondence between \mathbb{S}^2 and $\hat{\mathbb{C}}$






- As we know, stereographic projection doesn't map the point N at the top of the circle, so that's why we map N to infinity and need to consider the extended complex plane
- It can also be written in terms of θ and ϕ

Fractional Linear Transformation: Stereographic Projection

• Use Stereographic Projection to map \mathbb{S}^2 to the extended complex plane, $\hat{\mathbb{C}}=\mathbb{C}\cup\{\infty\}$



• The algebraic relation for a unit vector is

$$(x,y,z) = \left(\frac{\overline{\zeta} + \zeta}{\overline{\zeta}\zeta + 1}, i\frac{\overline{\zeta} - \zeta}{\overline{\zeta}\zeta + 1}, \frac{\overline{\zeta}\zeta - 1}{\overline{\zeta}\zeta + 1}\right),\,$$

ullet There is a one to one correspondence between \mathbb{S}^2 and $\hat{\mathbb{C}}$






- all the points on the 2 sphere are generators of the future null cone in Minkowskian space time
- Can denote an LT by moving three arbitrary points along the surface of the sphere
 as the generators have dimension two, so to match the dim of the LT (it's 6) we
 need three of them
- Extra coord is because we take time into account now, now ζ has two parameters so x is in terms of two parameters, t just defines the direction

Fractional Linear Transformation: Stereographic Projection

• Extend this to Minkowskian space-time[1]




- $(x, y, z, t) \leftrightarrow \mathbb{S}^2 \leftrightarrow \hat{\mathbb{C}}$
- So points in Minkowskian space-time can be written in the form

$$\vec{x} = t \left(\frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1}, 1 \right)$$



- all the points on the 2 sphere are generators of the future null cone in Minkowskian space time
- Can denote an LT by moving three arbitrary points along the surface of the sphere
 as the generators have dimension two, so to match the dim of the LT (it's 6) we
 need three of them
- Extra coord is because we take time into account now, now ζ has two parameters so x is in terms of two parameters, t just defines the direction


Extend this to Minkowskian space-time[1]



- $(x, y, z, t) \leftrightarrow \mathbb{S}^2 \leftrightarrow \hat{\mathbb{C}}$
- So points in Minkowskian space-time can be written in the form

$$\vec{x} = t \left(\frac{\overline{\zeta} + \zeta}{\overline{\zeta}\zeta + 1}, i \frac{\overline{\zeta} - \zeta}{\overline{\zeta}\zeta + 1}, \frac{\overline{\zeta}\zeta - 1}{\overline{\zeta}\zeta + 1}, 1 \right).$$






- all the points on the 2 sphere are generators of the future null cone in Minkowskian space time
- Can denote an LT by moving three arbitrary points along the surface of the sphere
 as the generators have dimension two, so to match the dim of the LT (it's 6) we
 need three of them
- Extra coord is because we take time into account now, now ζ has two parameters so x is in terms of two parameters, t just defines the direction

Fractional Linear Transformation: Stereographic Projection


• Extend this to Minkowskian space-time[1]



- $(x, y, z, t) \leftrightarrow \mathbb{S}^2 \leftrightarrow \hat{\mathbb{C}}$
- So points in Minkowskian space-time can be written in the form

$$\vec{x} = t \left(\frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1}, 1 \right).$$






- all the points on the 2 sphere are generators of the future null cone in Minkowskian space time
- Can denote an LT by moving three arbitrary points along the surface of the sphere
 as the generators have dimension two, so to match the dim of the LT (it's 6) we
 need three of them
- Extra coord is because we take time into account now, now ζ has two parameters so x is in terms of two parameters, t just defines the direction

Fractional Linear Transformation: Stereographic Projection

• Extend this to Minkowskian space-time[1]



- $(x, y, z, t) \leftrightarrow \mathbb{S}^2 \leftrightarrow \hat{\mathbb{C}}$
- So points in Minkowskian space-time can be written in the form

$$\vec{x} = t \left(\frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1}, 1 \right).$$







- These are null directions
- Refer to eqn (27) which should be on the board
- AS we did in the previous example, determine U
- ullet Remember we had $\pm U$ now the signs will cancel in the denominator and numerator

Fractional Linear Transformation

• Make the transformation $\zeta \to \zeta'$ by constructing the matrix $A(\vec{x})$ and determining the matrix U.

$$A(\vec{x}) = \begin{pmatrix} \frac{2t}{\zeta\bar{\zeta}+1} & \frac{2t\zeta}{\zeta\bar{\zeta}+1} \\ \frac{2t\bar{\zeta}}{\zeta\bar{\zeta}+1} & \frac{2t\zeta\bar{\zeta}}{\zeta\bar{\zeta}+1} \end{pmatrix} = c_0 \begin{pmatrix} \frac{1}{\zeta} & \frac{\zeta}{\zeta\zeta} \end{pmatrix},$$

$$c_0{'}\left(\begin{array}{cc} 1 & \zeta' \\ \bar{\zeta}' & \bar{\zeta}'\zeta' \end{array}\right) = \left(\begin{array}{cc} \alpha & \beta \\ \gamma & \delta \end{array}\right) c_0 \left(\begin{array}{cc} 1 & \zeta \\ \bar{\zeta} & \bar{\zeta}\zeta \end{array}\right) \left(\begin{array}{cc} \bar{\alpha} & \bar{\beta} \\ \bar{\gamma} & \bar{\delta} \end{array}\right).$$

• Solve for ζ' to get the fractional linear transformation

$$\zeta' = \frac{(\bar{\gamma} + \bar{\delta}\zeta)}{(\bar{\alpha} + \bar{\beta}\zeta)}$$

• There is a one to one correspondence between POLTs and fractional linear transformations



Fractional Linear Transformation



- These are null directions
- Refer to eqn (27) which should be on the board
- AS we did in the previous example, determine U
- ullet Remember we had $\pm U$ now the signs will cancel in the denominator and numerator

Fractional Linear Transformation

• Make the transformation $\zeta \to \zeta'$ by constructing the matrix $A(\vec{x})$ and determining the matrix U.

$$A(\vec{x}) = \begin{pmatrix} \frac{2t}{\zeta\bar{\zeta}+1} & \frac{2t\zeta}{\zeta\bar{\zeta}+1} \\ \frac{2t\bar{\zeta}}{\zeta\bar{\zeta}+1} & \frac{2t\zeta\bar{\zeta}}{\zeta\bar{\zeta}+1} \end{pmatrix} = c_0 \begin{pmatrix} \frac{1}{\zeta} & \frac{\zeta}{\zeta\zeta} \end{pmatrix},$$

$$c_0{'}\left(egin{array}{cc} 1 & \zeta' \ ar{\zeta'} & ar{\zeta'}\zeta' \end{array}
ight) = \left(egin{array}{cc} lpha & eta \ \gamma & \delta \end{array}
ight) c_0 \left(egin{array}{cc} 1 & \zeta \ ar{\zeta} & ar{\zeta}\zeta \end{array}
ight) \left(egin{array}{cc} ar{lpha} & ar{eta} \ ar{\gamma} & ar{\delta} \end{array}
ight).$$

• Solve for ζ' to get the fractional linear transformation

$$\zeta' = \frac{(\bar{\gamma} + \bar{\delta}\zeta)}{(\bar{\alpha} + \bar{\beta}\zeta)}$$

• There is a one to one correspondence between POLTs and fractional linear transformations



—Fractional Linear Transformation



- These are null directions
- Refer to eqn (27) which should be on the board
- AS we did in the previous example, determine U
- ullet Remember we had $\pm U$ now the signs will cancel in the denominator and numerator

Fractional Linear Transformation

• Make the transformation $\zeta \to \zeta'$ by constructing the matrix $A(\vec{x})$ and determining the matrix U.

$$A(\vec{x}) = \begin{pmatrix} \frac{2t}{\zeta\bar{\zeta}+1} & \frac{2t\zeta}{\zeta\bar{\zeta}+1} \\ \frac{2t\bar{\zeta}}{\zeta\bar{\zeta}+1} & \frac{2t\zeta\bar{\zeta}}{\zeta\bar{\zeta}+1} \end{pmatrix} = c_0 \begin{pmatrix} \frac{1}{\zeta} & \frac{\zeta}{\zeta\zeta} \end{pmatrix},$$

$$c_0{'}\left(egin{array}{cc} 1 & \zeta' \ ar{\zeta'} & ar{\zeta'}\zeta' \end{array}
ight) = \left(egin{array}{cc} lpha & eta \ \gamma & \delta \end{array}
ight) c_0 \left(egin{array}{cc} 1 & \zeta \ ar{\zeta} & ar{\zeta}\zeta \end{array}
ight) \left(egin{array}{cc} ar{lpha} & ar{eta} \ ar{\gamma} & ar{\delta} \end{array}
ight).$$

• Solve for ζ' to get the fractional linear transformation

$$\zeta' = \frac{(\bar{\gamma} + \bar{\delta}\zeta)}{(\bar{\alpha} + \bar{\beta}\zeta)},$$

 There is a one to one correspondence between POLTs and fractional linear transformations



Fractional Linear Transformation



- These are null directions
- Refer to eqn (27) which should be on the board
- ullet AS we did in the previous example, determine U
- ullet Remember we had $\pm U$ now the signs will cancel in the denominator and numerator

Fractional Linear Transformation

• Make the transformation $\zeta \to \zeta'$ by constructing the matrix $A(\vec{x})$ and determining the matrix U.

$$A(\vec{x}) = \begin{pmatrix} \frac{2t}{\zeta\bar{\zeta}+1} & \frac{2t\zeta}{\zeta\bar{\zeta}+1} \\ \frac{2t\bar{\zeta}}{\zeta\bar{\zeta}+1} & \frac{2t\zeta\bar{\zeta}}{\zeta\bar{\zeta}+1} \end{pmatrix} = c_0 \begin{pmatrix} \frac{1}{\zeta} & \frac{\zeta}{\zeta\zeta} \end{pmatrix},$$

$$c_0{'}\left(egin{array}{cc} 1 & \zeta' \ ar{\zeta'} & ar{\zeta'}\zeta' \end{array}
ight) = \left(egin{array}{cc} lpha & eta \ \gamma & \delta \end{array}
ight) c_0 \left(egin{array}{cc} 1 & \zeta \ ar{\zeta} & ar{\zeta}\zeta \end{array}
ight) \left(egin{array}{cc} ar{lpha} & ar{eta} \ ar{\gamma} & ar{\delta} \end{array}
ight).$$

• Solve for ζ' to get the fractional linear transformation

$$\zeta' = \frac{(\bar{\gamma} + \bar{\delta}\zeta)}{(\bar{\alpha} + \bar{\beta}\zeta)},$$

• There is a one to one correspondence between POLTs and fractional linear transformations



Fractional Linear Transformation



- These are null directions
- Refer to eqn (27) which should be on the board
- ullet AS we did in the previous example, determine U
- ullet Remember we had $\pm U$ now the signs will cancel in the denominator and numerator

Fractional Linear Transformation

• Make the transformation $\zeta \to \zeta'$ by constructing the matrix $A(\vec{x})$ and determining the matrix U.

$$A(\vec{x}) = \begin{pmatrix} \frac{2t}{\zeta\bar{\zeta}+1} & \frac{2t\zeta}{\zeta\bar{\zeta}+1} \\ \frac{2t\bar{\zeta}}{\zeta\bar{\zeta}+1} & \frac{2t\zeta\bar{\zeta}}{\zeta\bar{\zeta}+1} \end{pmatrix} = c_0 \begin{pmatrix} \frac{1}{\zeta} & \frac{\zeta}{\zeta\zeta} \end{pmatrix},$$

$$c_0{'}\left(egin{array}{cc} 1 & \zeta' \ ar{\zeta'} & ar{\zeta'}\zeta' \end{array}
ight) = \left(egin{array}{cc} lpha & eta \ \gamma & \delta \end{array}
ight) c_0 \left(egin{array}{cc} 1 & \zeta \ ar{\zeta} & ar{\zeta}\zeta \end{array}
ight) \left(egin{array}{cc} ar{lpha} & ar{eta} \ ar{\gamma} & ar{\delta} \end{array}
ight).$$

• Solve for ζ' to get the fractional linear transformation

$$\zeta' = \frac{(\bar{\gamma} + \bar{\delta}\zeta)}{(\bar{\alpha} + \bar{\beta}\zeta)},$$

• There is a one to one correspondence between POLTs and fractional linear transformations



-Layout

temp

Layout

Strange Minkowskian Line Element

SL(2, C) Matrices of the Lorentz Transformation

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- **3** $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- **1** The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions

 $U = \pm \begin{pmatrix} 1 + \epsilon s & \epsilon b \\ \epsilon c & 1 + \epsilon f \end{pmatrix}$

Infinitesimal Lorentz Transformation

- This was done in a recent lecture(Relativistic QM) so I wont do it
- Where a,b,c,d are complex
- ullet Take many infinitesimal LT steps along a particles trajectory and let ϵ go to zero

Infinitesimal Lorentz Transformation

$$U=\pm\left(egin{array}{cc} 1+\epsilon \mathsf{a} & \epsilon \mathsf{b} \ \epsilon \mathsf{c} & 1+\epsilon \mathsf{f} \end{array}
ight),$$

$$\bar{x}^i = x^i + \epsilon L^i_{j} x^j + O(\epsilon^2),$$

here

$$L^{i}_{j} = \begin{pmatrix} 0 & -2a_{2} & (b_{1} - c_{1}) & (b_{1} + c_{1}) \\ 2a_{2} & 0 & (b_{2} + c_{2}) & (b_{2} - c_{2}) \\ -(b_{1} - c_{1}) & -(b_{2} + c_{2}) & 0 & -2a_{1} \\ (b_{1} + c_{1}) & (b_{2} - c_{2}) & -2a_{1} & 0 \end{pmatrix}$$

$$\frac{d^{2}x^{i}}{ds^{2}} = L^{i}_{j}(s)\frac{dx^{j}}{ds^{j}}.$$

 $U = \pm \begin{pmatrix} 1 + \epsilon s & \epsilon b \\ \epsilon c & 1 + \epsilon f \end{pmatrix}$ $\vec{x}^i = x^i + \epsilon L^i_i x^i + O(\epsilon^2),$

- This was done in a recent lecture(Relativistic QM) so I wont do it
- Where a,b,c,d are complex
- Take many infinitesimal LT steps along a particles trajectory and let ϵ go to zero

$$U = \pm \begin{pmatrix} 1 + \epsilon a & \epsilon b \\ \epsilon c & 1 + \epsilon f \end{pmatrix},$$
$$\bar{x}^i = x^i + \epsilon L^i_i x^j + O(\epsilon^2),$$

where

$$L^{i}_{j} = \begin{pmatrix} 0 & -2a_{2} & (b_{1} - c_{1}) & (b_{1} + c_{1}) \\ 2a_{2} & 0 & (b_{2} + c_{2}) & (b_{2} - c_{2}) \\ -(b_{1} - c_{1}) & -(b_{2} + c_{2}) & 0 & -2a_{1} \\ (b_{1} + c_{1}) & (b_{2} - c_{2}) & -2a_{1} & 0 \end{pmatrix}$$

$$\frac{d^{2}x^{i}}{ds^{2}} = L^{i}_{j}(s)\frac{dx^{j}}{ds}.$$

- This was done in a recent lecture(Relativistic QM) so I wont do it
- Where a,b,c,d are complex
- ullet Take many infinitesimal LT steps along a particles trajectory and let ϵ go to zero

Infinitesimal Lorentz Transformation

$$U = \pm \begin{pmatrix} 1 + \epsilon a & \epsilon b \\ \epsilon c & 1 + \epsilon f \end{pmatrix},$$
$$\bar{x}^i = x^i + \epsilon L^i_j x^j + O(\epsilon^2),$$

where

$$L^{i}_{j} = \begin{pmatrix} 0 & -2a_{2} & (b_{1} - c_{1}) & (b_{1} + c_{1}) \\ 2a_{2} & 0 & (b_{2} + c_{2}) & (b_{2} - c_{2}) \\ -(b_{1} - c_{1}) & -(b_{2} + c_{2}) & 0 & -2a_{1} \\ (b_{1} + c_{1}) & (b_{2} - c_{2}) & -2a_{1} & 0 \end{pmatrix}$$

$$\frac{d^{2}x^{i}}{ds^{2}} = L^{i}_{j}(s)\frac{dx^{j}}{ds}.$$

- This was done in a recent lecture(Relativistic QM) so I wont do it
- Where a,b,c,d are complex
- ullet Take many infinitesimal LT steps along a particles trajectory and let ϵ go to zero

Infinitesimal Lorentz Transformation

$$U = \pm \begin{pmatrix} 1 + \epsilon a & \epsilon b \\ \epsilon c & 1 + \epsilon f \end{pmatrix},$$
$$\bar{x}^i = x^i + \epsilon L^i_j x^j + O(\epsilon^2),$$

where

$$L^{i}_{j} = \begin{pmatrix} 0 & -2a_{2} & (b_{1} - c_{1}) & (b_{1} + c_{1}) \\ 2a_{2} & 0 & (b_{2} + c_{2}) & (b_{2} - c_{2}) \\ -(b_{1} - c_{1}) & -(b_{2} + c_{2}) & 0 & -2a_{1} \\ (b_{1} + c_{1}) & (b_{2} - c_{2}) & -2a_{1} & 0 \end{pmatrix}$$

$$\frac{d^{2}x^{i}}{ds^{2}} = L^{i}_{j}(s)\frac{dx^{j}}{ds}.$$

$$\begin{split} \frac{d}{dt} \left(\gamma(u) u^{(1)} \right) &= -2 z_0 u^{(2)} + \left(b_1 - c_1 \right) u^{(0)} + b_1 + c_1, \\ \frac{d}{dt} \left(\gamma(u) u^{(2)} \right) &= 2 z_0 u^{(1)} + \left(b_2 + c_1 \right) u^{(2)} + b_2 - c_2, \\ \frac{d}{dt} \left(\gamma(u) u^{(2)} \right) &= -\left(b_1 - c_1 \right) u^{(1)} - \left(b_2 + c_2 \right) u^{(2)} - 2 z_1, \\ \frac{d\gamma(u)}{dt} &= \left(b_1 + c_1 \right) u^{(1)} + \left(b_2 - c_2 \right) u^{(2)} - 2 z_1 u^{(2)}. \end{split}$$

Clear that these equations can be expressed in terms of P and Q

Infinitesimal Lorentz Transformation: Lorentz Force

ullet Can rewrite this equation in terms of the particles 3-velocity $ec{u}$, in component form

$$\frac{d}{dt}(\gamma(u)u^{(1)}) = -2a_2u^{(2)} + (b_1 - c_1)u^{(3)} + b_1 + c_1,
\frac{d}{dt}(\gamma(u)u^{(2)}) = 2a_2u^{(1)} + (b_2 + c_2)u^{(3)} + b_2 - c_2,
\frac{d}{dt}(\gamma(u)u^{(3)}) = -(b_1 - c_1)u^{(1)} - (b_2 + c_2)u^{(2)} - 2a_1,
\frac{d\gamma(u)}{dt} = (b_1 + c_1)u^{(1)} + (b_2 - c_2)u^{(2)} - 2a_1u^{(3)}.$$

• Define the 3-vectors

$$\vec{P} = (b_1 + c_1, b_2 - c_2, -2a_1),$$

 $\vec{Q} = (b_2 + c_2, -(b_1 - c_1), -2a_2)$



Infinitesimal L

Infinitesimal Lorentz Transformation: Lorentz Force

$$\begin{split} \frac{d}{dt} [\gamma(u)u^{(k)}] &= -2u_0u^{(k)} + (k_1 - c_1)u^{(k)} + k_2 + c_1, \\ \frac{d}{dt} [\gamma(u)u^{(k)}] &= 2u_0u^{(k)} + (b_2 + c_1)u^{(k)} + k_2 - c_2, \\ \frac{d}{dt} [\gamma(u)u^{(k)}] &= -(k_1 - c_1)u^{(k)} - (k_2 + c_2)u^{(k)} - 2u_0, \\ \frac{d}{dt} [\gamma(u)u^{(k)}] &= -(k_1 - c_1)u^{(k)} - (k_2 + c_2)u^{(k)} - 2u_0, \\ \frac{d}{dt} [\gamma(u)u^{(k)}] &= -(k_1 - c_1)u^{(k)} + (k_2 - c_2)u^{(k)} - 2u_1u^{(k)}, \end{split}$$

 $\vec{P} = (b_1 + c_1, b_2 - c_2, -2s_1),$ $\vec{Q} = (b_2 + c_2, -(b_1 - c_1), -2s_2).$

Clear that these equations can be expressed in terms of P and Q

Infinitesimal Lorentz Transformation: Lorentz Force

• Can rewrite this equation in terms of the particles 3-velocity \vec{u} , in component form

$$\frac{d}{dt}(\gamma(u)u^{(1)}) = -2a_2u^{(2)} + (b_1 - c_1)u^{(3)} + b_1 + c_1,
\frac{d}{dt}(\gamma(u)u^{(2)}) = 2a_2u^{(1)} + (b_2 + c_2)u^{(3)} + b_2 - c_2,
\frac{d}{dt}(\gamma(u)u^{(3)}) = -(b_1 - c_1)u^{(1)} - (b_2 + c_2)u^{(2)} - 2a_1,
\frac{d\gamma(u)}{dt} = (b_1 + c_1)u^{(1)} + (b_2 - c_2)u^{(2)} - 2a_1u^{(3)}.$$

• Define the 3-vectors

$$\vec{P} = (b_1 + c_1, b_2 - c_2, -2a_1),$$

 $\vec{Q} = (b_2 + c_2, -(b_1 - c_1), -2a_2).$



Infinitesi

Infinitesimal Lorentz Transformation: Lorentz Force

$$\begin{split} \frac{d}{dt}(z(a)d^{(i)}) &= -2a_0d^{(i)} + (b_1 - c_1)d^{(i)} + b_1 + c_1, \\ \frac{d}{dt}(z(a)d^{(i)}) &= 2a_0d^{(i)} + (b_2 + c_1)d^{(i)} + b_2 - c_2, \\ \frac{d}{dt}(z(a)d^{(i)}) &= -(b_1 - c_1)d^{(i)} - (b_2 + c_2)d^{(i)} - 2a_1, \\ \frac{d}{dt}(z(a)d^{(i)}) &= -(b_1 - c_1)d^{(i)} - (b_2 + c_2)d^{(i)} - 2a_2, \\ \frac{d}{dt}(z(a)d^{(i)}) &= -(b_1 - c_1)d^{(i)} + (b_2 - c_2)d^{(i)} - 2a_1d^{(i)}, \\ \end{aligned}$$
 $\mathbf{y} \text{ Diris this Yadion}. \end{split}$

 $\vec{P} = (b_1 + c_1, b_2 - c_2, -2s_1),$ $\vec{Q} = (b_2 + c_2, -(b_1 - c_1), -2s_2).$

Clear that these equations can be expressed in terms of P and Q

Infinitesimal Lorentz Transformation: Lorentz Force

ullet Can rewrite this equation in terms of the particles 3-velocity $ec{u}$, in component form

$$\frac{d}{dt}(\gamma(u)u^{(1)}) = -2a_2u^{(2)} + (b_1 - c_1)u^{(3)} + b_1 + c_1,
\frac{d}{dt}(\gamma(u)u^{(2)}) = 2a_2u^{(1)} + (b_2 + c_2)u^{(3)} + b_2 - c_2,
\frac{d}{dt}(\gamma(u)u^{(3)}) = -(b_1 - c_1)u^{(1)} - (b_2 + c_2)u^{(2)} - 2a_1,
\frac{d\gamma(u)}{dt} = (b_1 + c_1)u^{(1)} + (b_2 - c_2)u^{(2)} - 2a_1u^{(3)}.$$

• Define the 3-vectors

$$\vec{P} = (b_1 + c_1, b_2 - c_2, -2a_1),$$

 $\vec{Q} = (b_2 + c_2, -(b_1 - c_1), -2a_2).$



Infinitesimal Lorentz Transformation: Lorentz Force

• temp

Infinitesimal Lorentz Transformation: Lorentz Force

• Writing the equations in terms of these

$$\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$$

- This is the same form as the Lorentz force
- Make the Identification

$$\vec{P} = \frac{q}{m}\vec{E}, \qquad \vec{Q} = \frac{q}{m}\vec{B},$$
 (1)

• To be compatible with special relativity the Lorentz force must depend on \vec{u} in this way. So the Lorentz force is a special case of a charged particle moving along a world line in Minkowskian space-time generated by an infinitesimal Lorentz transformation.

nfinitesimal Lorentz Transformation: Lorentz Force

 $\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$

temp

nfinitesimal Lorentz Transformation: Lorentz Force Infinitesimal Lorentz Transformation: Lorentz Force

a Writing the equations in terms of these $\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$ This is the same form as the Lorentz force

Writing the equations in terms of these

$$\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$$

- This is the same form as the Lorentz force
- Make the Identification

$$= -\frac{q}{m}\vec{E}, \qquad \vec{Q} = -\frac{q}{m}\vec{B}, \tag{1}$$

• To be compatible with special relativity the Lorentz force must depend on \vec{u}

Infinitesimal Lorentz Transformation: Lorentz Force



temp

Infinitesimal Lorentz Transformation: Lorentz Force

• Writing the equations in terms of these

$$\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$$

- This is the same form as the Lorentz force
- Make the Identification

$$\vec{P} = \frac{q}{m}\vec{E}, \qquad \vec{Q} = \frac{q}{m}\vec{B},$$
 (1)

• To be compatible with special relativity the Lorentz force must depend on \vec{u} in this way. So the Lorentz force is a special case of a charged particle moving along a world line in Minkowskian space-time generated by an infinitesimal Lorentz transformation

Infinitesimal Lorentz Transformation: Lorentz Force

temp

Infinitesimal Lorentz Transformation: Lorentz Force

• Writing the equations in terms of these

$$\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$$

- This is the same form as the Lorentz force
- Make the Identification

$$\vec{P} = \frac{q}{m}\vec{E}, \qquad \vec{Q} = \frac{q}{m}\vec{B},$$
 (1)

• To be compatible with special relativity the Lorentz force must depend on \vec{u} in this way. So the Lorentz force is a special case of a charged particle moving along a world line in Minkowskian space-time generated by an infinitesimal Lorentz transformation.



nfinitesimal Lorentz Transformation: Lorentz Force

 $\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$

 $\vec{P} = \frac{q}{-}\vec{E}, \quad \vec{Q} = \frac{q}{-}\vec{B},$

a Writing the equations in terms of these

. This is the same form as the Lorentz force

Make the Identification

Layout

temp

Layout

Strange Minkowskian Line Element

SL(2, C) Matrices of the Lorentz Transformation

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- **3** $SL(2,\mathbb{C})$ Matrices of the Lorentz Transformation
- **1** The Fractional Linear Transformation
- Infinitesimal Lorentz Transformation
- Pure Radiation Conditions





- This is derived from the form of the infinitesimal U and from the fractional linear transformation formula
- This is one of the things I said I would show earlier
- THIS WILL BE VERY IMPORTANT, **write on the board**

• The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- \bullet Fixed points of the system are given by $\zeta=\zeta'$ and correspond to null directions
- ullet With this condition solve the fractional linear transformation for ζ

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0$$

- A quadratic means it has two roots in general
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0$$
.





- This is derived from the form of the infinitesimal U and from the fractional linear transformation formula
- This is one of the things I said I would show earlier
- THIS WILL BE VERY IMPORTANT, **write on the board**

Pure Radiation Conditions

• The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- Fixed points of the system are given by $\zeta=\zeta'$ and correspond to null directions
- ullet With this condition solve the fractional linear transformation for ζ

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0$$

- A quadratic means it has two roots in general
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0$$
.





- This is derived from the form of the infinitesimal U and from the fractional linear transformation formula
- This is one of the things I said I would show earlier
- THIS WILL BE VERY IMPORTANT, **write on the board**

Pure Radiation Conditions

• The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- Fixed points of the system are given by $\zeta=\zeta'$ and correspond to null directions
- ullet With this condition solve the fractional linear transformation for ζ

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0.$$

- A quadratic means it has two roots in general
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0$$
.



| Pure Radiation | Conditions | |
|----------------|------------|--|
| | | |



- This is derived from the form of the infinitesimal U and from the fractional linear transformation formula
- This is one of the things I said I would show earlier
- THIS WILL BE VERY IMPORTANT, **write on the board**

• The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- \bullet Fixed points of the system are given by $\zeta=\zeta'$ and correspond to null directions
- ullet With this condition solve the fractional linear transformation for ζ

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0.$$

- A quadratic means it has two roots in general
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0$$
.





- This is derived from the form of the infinitesimal U and from the fractional linear transformation formula
- This is one of the things I said I would show earlier
- THIS WILL BE VERY IMPORTANT, **write on the board**

Pure Radiation Conditions

• The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- Fixed points of the system are given by $\zeta=\zeta'$ and correspond to null directions
- ullet With this condition solve the fractional linear transformation for ζ

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0.$$

- A quadratic means it has two roots in general
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0$$
.





- This is derived from the form of the infinitesimal U and from the fractional linear transformation formula
- This is one of the things I said I would show earlier
- THIS WILL BE VERY IMPORTANT, **write on the board**

Pure Radiation Conditions

• The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- Fixed points of the system are given by $\zeta=\zeta'$ and correspond to null directions
- ullet With this condition solve the fractional linear transformation for ζ

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0.$$

- A quadratic means it has two roots in general
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0$$
.



• q/m factors suppressed for convenience

Pure Radiation Conditions

ullet The a and b are related to $ec{E}$ and $ec{B}$ through the Lorentz force as

$$a_1 = -\frac{1}{2}E^3,$$
 $b_2 = \frac{1}{2}(E^2 + B^1),$ $c_1 = \frac{1}{2}(E^1 + B^2),$ $a_2 = -\frac{1}{2}B^3,$ $b_1 = \frac{1}{2}(E^1 - B^2),$ $c_2 = \frac{1}{2}(B^1 - E^2).$

 Then the real and imaginary parts of the quadratic condition give us the relations

$$|\vec{E}|^2 = |\vec{B}|^2$$

$$\vec{E} \cdot \vec{B} = 0.$$

• These are the familiar pure radiation conditions. Thus if the world line of a charged particle is generated by an infinitesimal singular Lorentz transformation then the particle is moving in a pure radiation EM field.

Pure Radiation Conditions

The a and b are related to \bar{E} and \bar{B} through the Lorentz force as $a_1 = -\frac{1}{2}E^2, \qquad b_2 = \frac{1}{2}(E^2 + B^1), \qquad c_1 = \frac{1}{2}(E^2 + B^2),$ $a_2 = -\frac{1}{2}B^1, \qquad b_1 = \frac{1}{2}(E^1 - B^2), \qquad c_2 = \frac{1}{2}(B^1 - E^2).$

• q/m factors suppressed for convenience

Pure Radiation Conditions

. The s and b are related to \bar{E} and \bar{B} through the Lorentz force as $\alpha_1=-\frac{1}{2}e^2, \quad b_1=\frac{1}{2}(e^2+\theta^2), \quad \alpha_1=\frac{1}{2}(e^2+\theta^2), \quad \alpha_2=\frac{1}{2}(e^2+\theta^2),$ $\alpha_3=\frac{1}{2}e^2, \quad b_1=\frac{1}{2}e^2-b^2, \quad \alpha_3=\frac{1}{2}e^2-\bar{E}^2).$ Then the real already green of the quadratic condition gives on the relations

Pure Radiation Conditions

• The a and b are related to \vec{E} and \vec{B} through the Lorentz force as

$$a_1 = -\frac{1}{2}E^3,$$
 $b_2 = \frac{1}{2}(E^2 + B^1),$ $c_1 = \frac{1}{2}(E^1 + B^2),$ $a_2 = -\frac{1}{2}B^3,$ $b_1 = \frac{1}{2}(E^1 - B^2),$ $c_2 = \frac{1}{2}(B^1 - E^2).$

 Then the real and imaginary parts of the quadratic condition give us the relations

$$|\vec{E}|^2 = |\vec{B}|^2,$$

$$\vec{F} \cdot \vec{B} = 0.$$

• These are the familiar pure radiation conditions. Thus if the world line of a charged particle is generated by an infinitesimal singular Lorentz transformation then the particle is moving in a pure radiation EM field.



Pure Radiation Conditions

• q/m factors suppressed for convenience

Pure Radiation Conditions

The a and b are related to \tilde{E} and \tilde{B} through the Lorentz force as $s_1 = -\frac{1}{2}E^3$, $b_2 = \frac{1}{2}(E^2 + B^1)$, $c_1 = \frac{1}{2}(E^1 + B^2)$, $a_2 = -\frac{1}{2}B^3$, $b_1 = \frac{1}{2}(E^1 - B^2)$, $c_2 = \frac{1}{2}(B^1 - E^2)$.

 $\vec{E} \cdot \vec{B} = 0$.

These are the familiar pure radiation conditions. Thus if the world line of

Pure Radiation Conditions

• The a and b are related to \vec{E} and \vec{B} through the Lorentz force as

$$a_1 = -\frac{1}{2}E^3,$$
 $b_2 = \frac{1}{2}(E^2 + B^1),$ $c_1 = \frac{1}{2}(E^1 + B^2),$ $a_2 = -\frac{1}{2}B^3,$ $b_1 = \frac{1}{2}(E^1 - B^2),$ $c_2 = \frac{1}{2}(B^1 - E^2).$

• Then the real and imaginary parts of the quadratic condition give us the relations

$$|\vec{E}|^2 = |\vec{B}|^2,$$

$$\vec{E} \cdot \vec{B} = 0.$$

• These are the familiar pure radiation conditions. Thus if the world line of a charged particle is generated by an infinitesimal singular Lorentz transformation then the particle is moving in a pure radiation EM field.



References

- D. Finkelstein "Past-Future Asymmetry of the Gravitational Field of a Poi Particle" - Phys. Rev. Vol 110. (1958)
- http://iournals.aps.org/pr/pdf/10.1103/PhysRev.110.965 P.A. Hogan, C.Barrabès - "Advanced General Relativity: Gravity Waves, I. Robinson "Spherical Gravitational Waves" - Phys. Rev. Lett. 4 (1960)
- http://iournals.aps.ors/prl/pdf/10.1103/PhysRevLett.4.431 P.A Horan, C. Barrabès - "Singular Null Hypersurfaces" - World Scientific
- 5 R. Penrose, W. Rindler "Spinors and Space-Time: Volume 1, Two-Spinor Calculus and Relativistic Fields" - Cambridge University Press, (Feb 1987)
- Various Authors "Space-Time and Geometry: The Alfred Schild Lectures" University of Texas Press (March 21, 2012)

References

- 1 J.L. Synge "Relativity: The Special Theory" North Holland Publishing Company (1965)
- 2 D. Finkelstein "Past-Future Asymmetry of the Gravitational Field of a Point Particle" - Phys.Rev.Vol 110, (1958) http://journals.aps.org/pr/pdf/10.1103/PhysRev.110.965
- 3 P.A. Hogan, C.Barrabès "Advanced General Relativity: Gravity Waves, Spinning Particles and Black Holes" - Oxford University Press (May 2013)
- 4 I. Robinson "Spherical Gravitational Waves" Phys.Rev.Lett. 4 (1960) 431-432
 - http://journals.aps.org/prl/pdf/10.1103/PhysRevLett.4.431
- 5 P.A Hogan, C. Barrabès "Singular Null Hypersurfaces" World Scientific Pub Co Inc (April 2004)
- 6 R. Penrose, W. Rindler "Spinors and Space-Time: Volume 1, Two-Spinor Calculus and Relativistic Fields" - Cambridge University Press, (Feb 1987)
- 7 Tristan Needham "Visual Complex Analysis" Clarendon Press, Oxford (1997)
- 8 Various Authors "Space-Time and Geometry: The Alfred Schild Lectures" -University of Texas Press (March 21, 2012)