

# Singular Lorentz Transformations and Pure Radiation Fields

Kevin Maguire

April 23, 2014

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions



# Introduction: Lorentz Transformations

- A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2,$$

in the transformation

$$(x, y, z, t) \rightarrow (x', y', z', t')$$

- Take the Proper Orthochronous Lorentz Transformations (POLTs) which form the restricted Lorentz group  $SO^+(1, 3)$
- In general Lorentz transformations have two invariant null directions

# Introduction: Lorentz Transformations

- A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2,$$

in the transformation

$$(x, y, z, t) \rightarrow (x', y', z', t')$$

- Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group**  $SO^+(1, 3)$
- In general Lorentz transformations have two invariant null directions

# Introduction: Lorentz Transformations

- A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2,$$

in the transformation

$$(x, y, z, t) \rightarrow (x', y', z', t')$$

- Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group**  $SO^+(1, 3)$
- In general Lorentz transformations have two invariant null directions

# Introduction: Lorentz Transformations


- A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2,$$

in the transformation

$$(x, y, z, t) \rightarrow (x', y', z', t')$$

- Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group**  $SO^+(1, 3)$
- In general Lorentz transformations have two invariant null directions



- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

# Strange Minkowskian Line Element

- Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$

- Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

- Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

- Taking the limit as the energy,  $\mu \rightarrow 0$  gives The **Kasner Solution**

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

# Strange Minkowskian Line Element

- Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$

- Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

- Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

- Taking the limit as the energy,  $\mu \rightarrow 0$  gives The **Kasner Solution**

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

# Strange Minkowskian Line Element

- Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$

- Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

- Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

- Taking the limit as the energy,  $\mu \rightarrow 0$  gives The **Kasner Solution**

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$



# Strange Minkowskian Line Element

- Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$

- Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

- Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

- Taking the limit as the energy,  $\mu \rightarrow 0$  gives The **Kasner Solution**

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

# Strange Minkowskian Line Element

- Then with  $m = 0$  the strange Minkowskian line element is obtained

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2dudr.$$

- Set  $r = 0$  to give

$$\epsilon ds^2 = 0,$$

and this is shown to be a null geodesic.

# Strange Minkowskian Line Element

- Then with  $m = 0$  the strange Minkowskian line element is obtained

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2dudr.$$

- Set  $r = 0$  to give

$$\epsilon ds^2 = 0,$$

and this is shown to be a null geodesic.

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

# Singular Lorentz Transformation

- Define an arbitrary complex parameter  $\zeta := \xi + i\eta$ , to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2du dr.$$

- The transformation  $\zeta \rightarrow \zeta + w$ , where  $w \in \mathbb{C}$  is then trivial and leaves the single null geodesic  $r = 0$  invariant.
- In Cartesian coordinates this transformation becomes

$$x' + iy' = x + iy + w(t - z),$$

$$z' - t' = -r = z - t,$$

$$z' + t' = z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).$$

- Addition of complex numbers is commutative, and  $w$  has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

# Singular Lorentz Transformation

- Define an arbitrary complex parameter  $\zeta := \xi + i\eta$ , to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2dudr.$$

- The transformation  $\zeta \rightarrow \zeta + w$ , where  $w \in \mathbb{C}$  is then trivial and leaves the single null geodesic  $r = 0$  invariant.
- In Cartesian coordinates this transformation becomes

$$x' + iy' = x + iy + w(t - z),$$

$$z' - t' = -r = z - t,$$

$$z' + t' = z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).$$

- Addition of complex numbers is commutative, and  $w$  has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

# Singular Lorentz Transformation

- Define an arbitrary complex parameter  $\zeta := \xi + i\eta$ , to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2du dr.$$

- The transformation  $\zeta \rightarrow \zeta + w$ , where  $w \in \mathbb{C}$  is then trivial and leaves the single null geodesic  $r = 0$  invariant.
- In Cartesian coordinates this transformation becomes

$$x' + iy' = x + iy + w(t - z),$$

$$z' - t' = -r = z - t,$$

$$z' + t' = z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).$$

- Addition of complex numbers is commutative, and  $w$  has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

# Singular Lorentz Transformation

- Define an arbitrary complex parameter  $\zeta := \xi + i\eta$ , to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2du dr.$$

- The transformation  $\zeta \rightarrow \zeta + w$ , where  $w \in \mathbb{C}$  is then trivial and leaves the single null geodesic  $r = 0$  invariant.
- In Cartesian coordinates this transformation becomes

$$x' + iy' = x + iy + w(t - z),$$

$$z' - t' = -r = z - t,$$

$$z' + t' = z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).$$

- Addition of complex numbers is commutative, and  $w$  has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group



- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

# $SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- Construct the transformation  $A(\vec{x}') = UA(\vec{x})U^\dagger$ , where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of  $SL(2, \mathbb{C})$

# $SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- Construct the transformation  $A(\vec{x}') = UA(\vec{x})U^\dagger$ , where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of  $SL(2, \mathbb{C})$

# $SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- Construct the transformation  $A(\vec{x}') = UA(\vec{x})U^\dagger$ , where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of  $SL(2, \mathbb{C})$

# $SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- Construct the transformation  $A(\vec{x}') = UA(\vec{x})U^\dagger$ , where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of  $SL(2, \mathbb{C})$

# $SL(2, \mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$  and  $A(\vec{x})$  have the same determinant so the above transformation preserves the Lorentz quadratic form, thus it is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t' - z' & x' + iy' \\ x' - iy' & t' + z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$
$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t - z)\bar{\alpha} + (x + iy)\bar{\beta} & (t - z)\bar{\gamma} + (x + iy)\bar{\delta} \\ (x - iy)\bar{\alpha} + (t + z)\bar{\beta} & (x - iy)\bar{\gamma} + (t + z)\bar{\delta} \end{pmatrix}.$$

- Thus the **general relations**

$$\begin{aligned} t' - z' &= (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta}, \\ x' + iy' &= (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta}, \\ t' + z' &= (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}. \end{aligned}$$

# $SL(2, \mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$  and  $A(\vec{x})$  have the same determinant so the above transformation preserves the Lorentz quadratic form, thus it is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t' - z' & x' + iy' \\ x' - iy' & t' + z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$
$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t - z)\bar{\alpha} + (x + iy)\bar{\beta} & (t - z)\bar{\gamma} + (x + iy)\bar{\delta} \\ (x - iy)\bar{\alpha} + (t + z)\bar{\beta} & (x - iy)\bar{\gamma} + (t + z)\bar{\delta} \end{pmatrix}.$$

- Thus the **general relations**

$$\begin{aligned} t' - z' &= (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta}, \\ x' + iy' &= (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta}, \\ t' + z' &= (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}. \end{aligned}$$

# $SL(2, \mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$  and  $A(\vec{x})$  have the same determinant so the above transformation preserves the Lorentz quadratic form, thus it is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t' - z' & x' + iy' \\ x' - iy' & t' + z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$
$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t - z)\bar{\alpha} + (x + iy)\bar{\beta} & (t - z)\bar{\gamma} + (x + iy)\bar{\delta} \\ (x - iy)\bar{\alpha} + (t + z)\bar{\beta} & (x - iy)\bar{\gamma} + (t + z)\bar{\delta} \end{pmatrix}.$$

- Thus the **general relations**

$$\begin{aligned} t' - z' &= (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta}, \\ x' + iy' &= (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta}, \\ t' + z' &= (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}. \end{aligned}$$



# Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

$$x' + iy' = x + iy + w(t - z),$$

$$t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \quad \beta = 0,$$

$$\gamma = \bar{w}\alpha, \quad \delta = \alpha.$$

- So there are always **two** possible choices of  $U$

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus **there is a 2 to 1 correspondence between elements of  $SL(2, \mathbb{C})$  and POLTs**

## Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

$$x' + iy' = x + iy + w(t - z),$$

$$t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \quad \beta = 0,$$

$$\gamma = \bar{w}\alpha, \quad \delta = \alpha.$$

- So there are always **two** possible choices of  $U$

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus **there is a 2 to 1 correspondence between elements of  $SL(2, \mathbb{C})$  and POLTs**

## Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

$$x' + iy' = x + iy + w(t - z),$$

$$t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \quad \beta = 0,$$

$$\gamma = \bar{w}\alpha, \quad \delta = \alpha.$$

- So there are always **two** possible choices of  $U$

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus there is a 2 to 1 correspondence between elements of  $SL(2, \mathbb{C})$  and POLTs

## Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

$$x' + iy' = x + iy + w(t - z),$$

$$t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \quad \beta = 0,$$

$$\gamma = \bar{w}\alpha, \quad \delta = \alpha.$$

- So there are always **two** possible choices of  $U$


$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus **there is a 2 to 1 correspondence between elements of  $SL(2, \mathbb{C})$  and POLTs**

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

# Fractional Linear Transformation: Stereographic Projection

- Use **Stereographic Projection** to map  $\mathbb{S}^2$  to the **extended complex plane**,  $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$




- The algebraic relation for a unit vector is

$$(x, y, z) = \left( \frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1} \right),$$

- There is a one to one correspondence between  $\mathbb{S}^2$  and  $\hat{\mathbb{C}}$

# Fractional Linear Transformation: Stereographic Projection

- Use **Stereographic Projection** to map  $\mathbb{S}^2$  to the **extended complex plane**,  $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$




- The algebraic relation for a unit vector is

$$(x, y, z) = \left( \frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1} \right),$$

- There is a one to one correspondence between  $\mathbb{S}^2$  and  $\hat{\mathbb{C}}$

# Fractional Linear Transformation: Stereographic Projection

- Use **Stereographic Projection** to map  $\mathbb{S}^2$  to the **extended complex plane**,  $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$



- The algebraic relation for a unit vector is


$$(x, y, z) = \left( \frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1} \right),$$

- There is a one to one correspondence between  $\mathbb{S}^2$  and  $\hat{\mathbb{C}}$



# Fractional Linear Transformation: Stereographic Projection

- Use **Stereographic Projection** to map  $\mathbb{S}^2$  to the **extended complex plane**,  $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$




- The algebraic relation for a unit vector is

$$(x, y, z) = \left( \frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1} \right),$$

- There is a one to one correspondence between  $\mathbb{S}^2$  and  $\hat{\mathbb{C}}$

# Fractional Linear Transformation: Stereographic Projection

- Extend this to Minkowskian space-time[1]




- $(x, y, z, t) \leftrightarrow \mathbb{S}^2 \leftrightarrow \hat{\mathbb{C}}$
- So points in Minkowskian space-time can be written in the form

$$\vec{x} = t \left( \frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1}, 1 \right).$$

# Fractional Linear Transformation: Stereographic Projection

- Extend this to Minkowskian space-time[1]




- $(x, y, z, t) \leftrightarrow \mathbb{S}^2 \leftrightarrow \hat{\mathbb{C}}$

- So points in Minkowskian space-time can be written in the form

$$\vec{x} = t \left( \frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1}, 1 \right).$$

# Fractional Linear Transformation: Stereographic Projection

- Extend this to Minkowskian space-time[1]



- $(x, y, z, t) \leftrightarrow \mathbb{S}^2 \leftrightarrow \hat{\mathbb{C}}$
- So points in Minkowskian space-time can be written in the form

$$\vec{x} = t \left( \frac{\bar{\zeta} + \zeta}{\bar{\zeta}\zeta + 1}, i \frac{\bar{\zeta} - \zeta}{\bar{\zeta}\zeta + 1}, \frac{\bar{\zeta}\zeta - 1}{\bar{\zeta}\zeta + 1}, 1 \right).$$

# Fractional Linear Transformation

- Make the transformation  $\zeta \rightarrow \zeta'$  by constructing the matrix  $A(\vec{x})$  and determining the matrix  $U$ .

$$A(\vec{x}) = \begin{pmatrix} \frac{2t}{\zeta\bar{\zeta}+1} & \frac{2t\zeta}{\zeta\bar{\zeta}+1} \\ \frac{2t\bar{\zeta}}{\zeta\bar{\zeta}+1} & \frac{2t\zeta\bar{\zeta}}{\zeta\bar{\zeta}+1} \end{pmatrix} = c_0 \begin{pmatrix} 1 & \zeta \\ \bar{\zeta} & \zeta\bar{\zeta} \end{pmatrix},$$

$$c_0' \begin{pmatrix} 1 & \zeta' \\ \bar{\zeta}' & \zeta'\bar{\zeta}' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} c_0 \begin{pmatrix} 1 & \zeta \\ \bar{\zeta} & \zeta\bar{\zeta} \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\beta} \\ \bar{\gamma} & \bar{\delta} \end{pmatrix}.$$

- Solve for  $\zeta'$  to get the fractional linear transformation

$$\zeta' = \frac{(\bar{\gamma} + \bar{\delta}\zeta)}{(\bar{\alpha} + \bar{\beta}\zeta)},$$

- There is a one to one correspondence between POLTs and fractional linear transformations

# Fractional Linear Transformation

- Make the transformation  $\zeta \rightarrow \zeta'$  by constructing the matrix  $A(\vec{x})$  and determining the matrix  $U$ .

$$A(\vec{x}) = \begin{pmatrix} \frac{2t}{\zeta\bar{\zeta}+1} & \frac{2t\zeta}{\zeta\bar{\zeta}+1} \\ \frac{2t\bar{\zeta}}{\zeta\bar{\zeta}+1} & \frac{2t\zeta\bar{\zeta}}{\zeta\bar{\zeta}+1} \end{pmatrix} = c_0 \begin{pmatrix} 1 & \zeta \\ \bar{\zeta} & \zeta\bar{\zeta} \end{pmatrix},$$

$$c_0' \begin{pmatrix} 1 & \zeta' \\ \bar{\zeta}' & \zeta'\bar{\zeta}' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} c_0 \begin{pmatrix} 1 & \zeta \\ \bar{\zeta} & \zeta\bar{\zeta} \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\beta} \\ \bar{\gamma} & \bar{\delta} \end{pmatrix}.$$

- Solve for  $\zeta'$  to get the fractional linear transformation

$$\zeta' = \frac{(\bar{\gamma} + \bar{\delta}\zeta)}{(\bar{\alpha} + \bar{\beta}\zeta)},$$

- There is a one to one correspondence between POLTs and fractional linear transformations

# Fractional Linear Transformation

- Make the transformation  $\zeta \rightarrow \zeta'$  by constructing the matrix  $A(\vec{x})$  and determining the matrix  $U$ .

$$A(\vec{x}) = \begin{pmatrix} \frac{2t}{\zeta\bar{\zeta}+1} & \frac{2t\zeta}{\zeta\bar{\zeta}+1} \\ \frac{2t\bar{\zeta}}{\zeta\bar{\zeta}+1} & \frac{2t\zeta\bar{\zeta}}{\zeta\bar{\zeta}+1} \end{pmatrix} = c_0 \begin{pmatrix} 1 & \zeta \\ \bar{\zeta} & \zeta\bar{\zeta} \end{pmatrix},$$

$$c_0' \begin{pmatrix} 1 & \zeta' \\ \bar{\zeta}' & \zeta'\bar{\zeta}' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} c_0 \begin{pmatrix} 1 & \zeta \\ \bar{\zeta} & \zeta\bar{\zeta} \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\beta} \\ \bar{\gamma} & \bar{\delta} \end{pmatrix}.$$

- Solve for  $\zeta'$  to get the **fractional linear transformation**

$$\zeta' = \frac{(\bar{\gamma} + \bar{\delta}\zeta)}{(\bar{\alpha} + \bar{\beta}\zeta)},$$

- There is a **one to one correspondence between POLTs and fractional linear transformations**

# Fractional Linear Transformation

- Make the transformation  $\zeta \rightarrow \zeta'$  by constructing the matrix  $A(\vec{x})$  and determining the matrix  $U$ .

$$A(\vec{x}) = \begin{pmatrix} \frac{2t}{\zeta\bar{\zeta}+1} & \frac{2t\zeta}{\zeta\bar{\zeta}+1} \\ \frac{2t\bar{\zeta}}{\zeta\bar{\zeta}+1} & \frac{2t\zeta\bar{\zeta}}{\zeta\bar{\zeta}+1} \end{pmatrix} = c_0 \begin{pmatrix} 1 & \zeta \\ \bar{\zeta} & \zeta\bar{\zeta} \end{pmatrix},$$

$$c_0' \begin{pmatrix} 1 & \zeta' \\ \bar{\zeta}' & \zeta'\bar{\zeta}' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} c_0 \begin{pmatrix} 1 & \zeta \\ \bar{\zeta} & \zeta\bar{\zeta} \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\beta} \\ \bar{\gamma} & \bar{\delta} \end{pmatrix}.$$

- Solve for  $\zeta'$  to get the **fractional linear transformation**

$$\zeta' = \frac{(\bar{\gamma} + \bar{\delta}\zeta)}{(\bar{\alpha} + \bar{\beta}\zeta)},$$

- There is a **one to one correspondence between POLTs and fractional linear transformations**



- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

# Infinitesimal Lorentz Transformation

$$U = \pm \begin{pmatrix} 1 + \epsilon a & \epsilon b \\ \epsilon c & 1 + \epsilon f \end{pmatrix},$$

$$\bar{x}^i = x^i + \epsilon L^i_j x^j + O(\epsilon^2),$$

where

$$L^i_j = \begin{pmatrix} 0 & -2a_2 & (b_1 - c_1) & (b_1 + c_1) \\ 2a_2 & 0 & (b_2 + c_2) & (b_2 - c_2) \\ -(b_1 - c_1) & -(b_2 + c_2) & 0 & -2a_1 \\ (b_1 + c_1) & (b_2 - c_2) & -2a_1 & 0 \end{pmatrix}$$

$$\frac{d^2 x^i}{ds^2} = L^i_j(s) \frac{dx^j}{ds}.$$

# Infinitesimal Lorentz Transformation

$$U = \pm \begin{pmatrix} 1 + \epsilon a & \epsilon b \\ \epsilon c & 1 + \epsilon f \end{pmatrix},$$

$$\bar{x}^i = x^i + \epsilon L^i_j x^j + O(\epsilon^2),$$

where

$$L^i_j = \begin{pmatrix} 0 & -2a_2 & (b_1 - c_1) & (b_1 + c_1) \\ 2a_2 & 0 & (b_2 + c_2) & (b_2 - c_2) \\ -(b_1 - c_1) & -(b_2 + c_2) & 0 & -2a_1 \\ (b_1 + c_1) & (b_2 - c_2) & -2a_1 & 0 \end{pmatrix}$$

$$\frac{d^2 x^i}{ds^2} = L^i_j(s) \frac{dx^j}{ds}.$$

# Infinitesimal Lorentz Transformation

$$U = \pm \begin{pmatrix} 1 + \epsilon a & \epsilon b \\ \epsilon c & 1 + \epsilon f \end{pmatrix},$$

$$\bar{x}^i = x^i + \epsilon L^i_j x^j + O(\epsilon^2),$$

where

$$L^i_j = \begin{pmatrix} 0 & -2a_2 & (b_1 - c_1) & (b_1 + c_1) \\ 2a_2 & 0 & (b_2 + c_2) & (b_2 - c_2) \\ -(b_1 - c_1) & -(b_2 + c_2) & 0 & -2a_1 \\ (b_1 + c_1) & (b_2 - c_2) & -2a_1 & 0 \end{pmatrix}$$

$$\frac{d^2 x^i}{ds^2} = L^i_j(s) \frac{dx^j}{ds}.$$

# Infinitesimal Lorentz Transformation: Lorentz Force

- Can rewrite this equation in terms of the particles 3-velocity  $\vec{u}$ , in component form

$$\frac{d}{dt}(\gamma(u)u^{(1)}) = -2a_2u^{(2)} + (b_1 - c_1)u^{(3)} + b_1 + c_1,$$

$$\frac{d}{dt}(\gamma(u)u^{(2)}) = 2a_2u^{(1)} + (b_2 + c_2)u^{(3)} + b_2 - c_2,$$

$$\frac{d}{dt}(\gamma(u)u^{(3)}) = -(b_1 - c_1)u^{(1)} - (b_2 + c_2)u^{(2)} - 2a_1,$$

$$\frac{d\gamma(u)}{dt} = (b_1 + c_1)u^{(1)} + (b_2 - c_2)u^{(2)} - 2a_1u^{(3)}.$$

- Define the 3-vectors

$$\vec{P} = (b_1 + c_1, b_2 - c_2, -2a_1),$$

$$\vec{Q} = (b_2 + c_2, -(b_1 - c_1), -2a_2).$$

# Infinitesimal Lorentz Transformation: Lorentz Force

- Can rewrite this equation in terms of the particles 3-velocity  $\vec{u}$ , in component form

$$\frac{d}{dt}(\gamma(u)u^{(1)}) = -2a_2u^{(2)} + (b_1 - c_1)u^{(3)} + b_1 + c_1,$$

$$\frac{d}{dt}(\gamma(u)u^{(2)}) = 2a_2u^{(1)} + (b_2 + c_2)u^{(3)} + b_2 - c_2,$$

$$\frac{d}{dt}(\gamma(u)u^{(3)}) = -(b_1 - c_1)u^{(1)} - (b_2 + c_2)u^{(2)} - 2a_1,$$

$$\frac{d\gamma(u)}{dt} = (b_1 + c_1)u^{(1)} + (b_2 - c_2)u^{(2)} - 2a_1u^{(3)}.$$

- Define the 3-vectors

$$\vec{P} = (b_1 + c_1, b_2 - c_2, -2a_1),$$

$$\vec{Q} = (b_2 + c_2, -(b_1 - c_1), -2a_2).$$

# Infinitesimal Lorentz Transformation: Lorentz Force

- Writing the equations in terms of these

$$\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$$

- This is the same form as the Lorentz force
- Make the Identification

$$\vec{P} = \frac{q}{m}\vec{E}, \quad \vec{Q} = \frac{q}{m}\vec{B}, \quad (1)$$

- To be compatible with special relativity the Lorentz force must depend on  $\vec{u}$  in this way. So the Lorentz force is a special case of a charged particle moving along a world line in Minkowskian space-time generated by an infinitesimal Lorentz transformation.

# Infinitesimal Lorentz Transformation: Lorentz Force

- Writing the equations in terms of these

$$\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$$

- This is the same form as the **Lorentz force**
- Make the Identification

$$\vec{P} = \frac{q}{m}\vec{E}, \quad \vec{Q} = \frac{q}{m}\vec{B}, \quad (1)$$

- To be compatible with special relativity the Lorentz force must depend on  $\vec{u}$  in this way. So the Lorentz force is a special case of a charged particle moving along a world line in Minkowskian space-time generated by an infinitesimal Lorentz transformation.



# Infinitesimal Lorentz Transformation: Lorentz Force

- Writing the equations in terms of these

$$\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$$

- This is the same form as the **Lorentz force**
- Make the Identification

$$\vec{P} = \frac{q}{m}\vec{E}, \quad \vec{Q} = \frac{q}{m}\vec{B}, \quad (1)$$

- To be compatible with special relativity the Lorentz force must depend on  $\vec{u}$  in this way. So the Lorentz force is a special case of a charged particle moving along a world line in Minkowskian space-time generated by an infinitesimal Lorentz transformation.

# Infinitesimal Lorentz Transformation: Lorentz Force

- Writing the equations in terms of these

$$\frac{d}{dt}(\gamma(u)\vec{u}) = \vec{P} + \vec{u} \times \vec{Q},$$

- This is the same form as the **Lorentz force**
- Make the Identification

$$\vec{P} = \frac{q}{m}\vec{E}, \quad \vec{Q} = \frac{q}{m}\vec{B}, \quad (1)$$

- **To be compatible with special relativity the Lorentz force must depend on  $\vec{u}$  in this way.** So the Lorentz force is a special case of a charged particle moving along a world line in Minkowskian space-time generated by an infinitesimal Lorentz transformation.

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4  $SL(2, \mathbb{C})$  Matrices of the Lorentz Transformation
- 5 The Fractional Linear Transformation
- 6 Infinitesimal Lorentz Transformation
- 7 Pure Radiation Conditions

# Pure Radiation Conditions

- The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- Fixed points of the system are given by  $\zeta = \zeta'$  and correspond to null directions
- With this condition solve the fractional linear transformation for  $\zeta$

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0.$$

- A quadratic means it has **two roots in general**
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0.$$

refer to this as the **quadratic condition**.

# Pure Radiation Conditions

- The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- Fixed points of the system are given by  $\zeta = \zeta'$  and correspond to null directions
- With this condition solve the fractional linear transformation for  $\zeta$

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0.$$

- A quadratic means it has **two roots in general**
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0.$$

refer to this as the **quadratic condition**.

# Pure Radiation Conditions

- The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- Fixed points of the system are given by  $\zeta = \zeta'$  and correspond to null directions
- With this condition solve the fractional linear transformation for  $\zeta$

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0.$$

- A quadratic means it has **two roots in general**
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0.$$

refer to this as the **quadratic condition**.

# Pure Radiation Conditions

- The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- Fixed points of the system are given by  $\zeta = \zeta'$  and correspond to null directions
- With this condition solve the fractional linear transformation for  $\zeta$

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0.$$

- A quadratic means it has **two roots in general**
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0.$$

refer to this as the **quadratic condition**.

# Pure Radiation Conditions

- The fractional linear transformation of the infinitesimal transformation is

$$\zeta' = \frac{\zeta + \epsilon(\bar{c} - \bar{a}\zeta) + O(\epsilon^2)}{1 + \epsilon(\bar{a} + \bar{b}\zeta) + O(\epsilon^2)}.$$

- Fixed points of the system are given by  $\zeta = \zeta'$  and correspond to null directions
- With this condition solve the fractional linear transformation for  $\zeta$

$$\bar{\beta}\zeta^2 + (\bar{\alpha} - \bar{\delta})\zeta - \bar{\gamma} = 0.$$

- A quadratic means it has **two roots in general**
- Interested in the singular root case so take the discriminant equal to zero to get

$$a^2 + bc = 0.$$

refer to this as the **quadratic condition**.



# Pure Radiation Conditions

- The  $a$  and  $b$  are related to  $\vec{E}$  and  $\vec{B}$  through the Lorentz force as

$$\begin{aligned}a_1 &= -\frac{1}{2}E^3, & b_2 &= \frac{1}{2}(E^2 + B^1), & c_1 &= \frac{1}{2}(E^1 + B^2), \\a_2 &= -\frac{1}{2}B^3, & b_1 &= \frac{1}{2}(E^1 - B^2), & c_2 &= \frac{1}{2}(B^1 - E^2).\end{aligned}$$

- Then the real and imaginary parts of the quadratic condition give us the relations

$$\begin{aligned}|\vec{E}|^2 &= |\vec{B}|^2, \\ \vec{E} \cdot \vec{B} &= 0.\end{aligned}$$

- These are the familiar pure radiation conditions. Thus if the world line of a charged particle is generated by an infinitesimal singular Lorentz transformation then the particle is moving in a pure radiation EM field.

# Pure Radiation Conditions

- The  $a$  and  $b$  are related to  $\vec{E}$  and  $\vec{B}$  through the Lorentz force as

$$\begin{aligned}a_1 &= -\frac{1}{2}E^3, & b_2 &= \frac{1}{2}(E^2 + B^1), & c_1 &= \frac{1}{2}(E^1 + B^2), \\a_2 &= -\frac{1}{2}B^3, & b_1 &= \frac{1}{2}(E^1 - B^2), & c_2 &= \frac{1}{2}(B^1 - E^2).\end{aligned}$$

- Then the real and imaginary parts of the quadratic condition give us the relations

$$\begin{aligned}|\vec{E}|^2 &= |\vec{B}|^2, \\ \vec{E} \cdot \vec{B} &= 0.\end{aligned}$$

- These are the familiar pure radiation conditions. Thus if the world line of a charged particle is generated by an infinitesimal singular Lorentz transformation then the particle is moving in a pure radiation EM field.

# Pure Radiation Conditions

- The  $a$  and  $b$  are related to  $\vec{E}$  and  $\vec{B}$  through the Lorentz force as

$$\begin{aligned} a_1 &= -\frac{1}{2}E^3, & b_2 &= \frac{1}{2}(E^2 + B^1), & c_1 &= \frac{1}{2}(E^1 + B^2), \\ a_2 &= -\frac{1}{2}B^3, & b_1 &= \frac{1}{2}(E^1 - B^2), & c_2 &= \frac{1}{2}(B^1 - E^2). \end{aligned}$$

- Then the real and imaginary parts of the quadratic condition give us the relations

$$\begin{aligned} |\vec{E}|^2 &= |\vec{B}|^2, \\ \vec{E} \cdot \vec{B} &= 0. \end{aligned}$$

- These are the familiar pure radiation conditions. Thus if the world line of a charged particle is generated by an infinitesimal singular Lorentz transformation then the particle is moving in a pure radiation EM field.

# References

- 1 J.L. Synge - "Relativity: The Special Theory" - North Holland Publishing Company (1965)
- 2 D. Finkelstein - "Past-Future Asymmetry of the Gravitational Field of a Point Particle" - Phys.Rev.Vol 110, (1958) -  
<http://journals.aps.org/pr/pdf/10.1103/PhysRev.110.965>
- 3 P.A. Hogan, C.Barrabès - "Advanced General Relativity: Gravity Waves, Spinning Particles and Black Holes" - Oxford University Press (May 2013)
- 4 I. Robinson "Spherical Gravitational Waves" - Phys.Rev.Lett. 4 (1960) 431-432 -  
<http://journals.aps.org/prl/pdf/10.1103/PhysRevLett.4.431>
- 5 P.A Hogan, C. Barrabès - "Singular Null Hypersurfaces" - World Scientific Pub Co Inc (April 2004)
- 6 R. Penrose, W. Rindler - "Spinors and Space-Time: Volume 1, Two-Spinor Calculus and Relativistic Fields" - Cambridge University Press, (Feb 1987)
- 7 Tristan Needham - "Visual Complex Analysis" - Clarendon Press, Oxford (1997)
- 8 Various Authors - "Space-Time and Geometry: The Alfred Schild Lectures" - University of Texas Press (March 21, 2012)