

Singular Lorentz Transformations and Pure Radiation Fields

Kevin Maguire

April 21, 2014

2014-04-21

Layout

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz Transformation
- $SL(2, \mathbb{C})$ Matrices of the Lorentz Transformation

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4 $SL(2, \mathbb{C})$ Matrices of the Lorentz Transformation

2014-04-21

Layout

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- $SL(2, \mathbb{C})$ Matrices of the Lorentz Transformation

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4 $SL(2, \mathbb{C})$ Matrices of the Lorentz Transformation

2014-04-21

Layout

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4 $SL(2, \mathbb{C})$ Matrices of the Lorentz Transformation

2014-04-21

Layout

- Introduction: Lorentz Transformations
- Strange Minkowskian Line Element
- Singular Lorentz transformation
- $SL(2, \mathbb{C})$ Matrices of the Lorentz Transformation

- 1 Introduction: Lorentz Transformations
- 2 Strange Minkowskian Line Element
- 3 Singular Lorentz transformation
- 4 $SL(2, \mathbb{C})$ Matrices of the Lorentz Transformation

Introduction: Lorentz Transformations

- A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2+y'^2+z'^2-t'^2 = x^2+y^2+z^2-t^2,$$

in the transformation
 $(x,y,z,t) \rightarrow (x',y',z',t')$

- Take the Proper Orthochronous Lorentz Transformations(POLTs) which form the restricted Lorentz group $SO^+(1,3)$
- In general lorentz transformations have two invariant null directions

2014-04-21

Introduction: Lorentz Transformations

▼ A Lorentz transformation is defined by the preservation of the quadratic form
 $x'^2+y'^2+z'^2-t'^2 = x^2+y^2+z^2-t^2,$
in the transformation
 $(x,y,z,t) \rightarrow (x',y',z',t')$
• Lorentz Proper Orthochronous Lorentz Transformations(POLTs) form the restricted Lorentz group $SO^+(1,3)$
• In general Lorentz transformations have two invariant null directions

- –Proper is $\det 1$. preserves the orientation of spacial axes, preserves handedness
- –orthochronous means time is always positive and the direction of time is preserved
- –Think of the standard Lorentz transformation, always two null directions at $x \pm t$

Introduction: Lorentz Transformations

- A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2,$$

in the transformation

$$(x, y, z, t) \rightarrow (x', y', z', t')$$

- Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group** $SO^+(1, 3)$
- In general lorentz transformations have two invariant null directions

2014-04-21

Introduction: Lorentz Transformations

▼ A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2,$$

in the transformation

$$(x, y, z, t) \rightarrow (x', y', z', t')$$

• Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group** $SO^+(1, 3)$

- –Proper is $\det 1$. preserves the orientation of spacial axes, preserves handedness
- –orthochronous means time is always positive and the direction of time is preserved
- –Think of the standard Lorentz transformation, always two null directions at $x \pm t$

Introduction: Lorentz Transformations

- A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2,$$

in the transformation

$$(x, y, z, t) \rightarrow (x', y', z', t')$$

- Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group** $SO^+(1, 3)$
- In general lorentz transformations have two invariant null directions

2014-04-21

Introduction: Lorentz Transformations

▼ A Lorentz transformation is defined by the preservation of the quadratic form
 $x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2$,
in the transformation
 $(x, y, z, t) \rightarrow (x', y', z', t')$
• Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group** $SO^+(1, 3)$
▼ In general lorentz transformations have two invariant null directions.

- –Proper is $\det 1$. preserves the orientation of spacial axes, preserves handedness
- –orthochronous means time is always positive and the direction of time is preserved
- –Think of the standard Lorentz transformation, always two null directions at $x \pm t$

Introduction: Lorentz Transformations


- A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2,$$

in the transformation

$$(x, y, z, t) \rightarrow (x', y', z', t')$$

- Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group** $SO^+(1, 3)$
- In general lorentz transformations have two invariant null directions



2014-04-21

Introduction: Lorentz Transformations

• A Lorentz transformation is defined by the preservation of the quadratic form
 $x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2$,
in the transformation
 $(x, y, z, t) \rightarrow (x', y', z', t')$
• Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group** $SO^+(1, 3)$
• In general lorentz transformations have two invariant null directions.



- -Proper is $\det 1$. preserves the orientation of spacial axes, preserves handedness
- -orthochronous means time is always positive and the direction of time is preserved
- -Think of the standard Lorentz transformation, always two null directions at $x \pm t$

Introduction: Lorentz Transformations


- A Lorentz transformation is defined by the preservation of the quadratic form

$$x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2,$$

in the transformation

$$(x, y, z, t) \rightarrow (x', y', z', t')$$

- Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group** $SO^+(1, 3)$
- In general lorentz transformations have two invariant null directions



2014-04-21

Introduction: Lorentz Transformations

• A Lorentz transformation is defined by the preservation of the quadratic form
 $x'^2 + y'^2 + z'^2 - t'^2 = x^2 + y^2 + z^2 - t^2$,
in the transformation
 $(x, y, z, t) \rightarrow (x', y', z', t')$
• Take the **Proper Orthochronous Lorentz Transformations (POLTs)** which form the **restricted Lorentz group** $SO^+(1, 3)$
• In general lorentz transformations have two invariant null directions.



- -Proper is $\det 1$. preserves the orientation of spacial axes, preserves handedness
- -orthochronous means time is always positive and the direction of time is preserved
- -Think of the standard Lorentz transformation, always two null directions at $x \pm t$

2014-04-21

Layout

add in the contents

add in the contents

- derive a strange minkowskian line element
- making a complicated transformation that keeps a single null geodesic fixed look trivial

Strange Minkowskian Line Element

- Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$

- Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

- Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

- Taking the limit as the energy, $\mu \rightarrow 0$ gives The **Kasner Solution**

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

2014-04-21

Strange Minkowskian Line Element

- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coord transforms that this is Kasner, but it won't be done here

- Start with the Schwarzschild solution
$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$
- Make the Eddington-Finkelstein coordinate transformation [2]
$$u = t - r - 2m \ln(r - 2m).$$
- Make further coordinate transformations to obtain
$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$
- Taking the limit as the energy, $\mu \rightarrow 0$ gives The **Kasner Solution**
$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

Strange Minkowskian Line Element

- Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$

- Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

- Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

- Taking the limit as the energy, $\mu \rightarrow 0$ gives The **Kasner Solution**

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

2014-04-21

Strange Minkowskian Line Element

- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coord transforms that this is Kasner, but it won't be done here

Start with the Schwarzschild solution
 $\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$
Make the Eddington-Finkelstein coordinate transformation [2]
 $u = t - r - 2m \ln(r - 2m).$
If you want to see the details of the coordinate transformations, see the notes on the Eddington-Finkelstein coordinate transformation.
Taking the limit as the energy goes to 0 gives the Kasner Solution.

2014-04-21

Strange Minkowskian Line Element

Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dt^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dr^2.$$

Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

It is easily shown with further coord transforms that this is Kasner, but it won't be done here

- Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$

- Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

- Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

- Taking the limit as the energy, $\mu \rightarrow 0$ gives The **Kasner Solution**

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coord transforms that this is Kasner, but it won't be done here

2014-04-21

Strange Minkowskian Line Element

Start with the Schwarzschild solution

$$ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dt^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dr^2.$$

Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

Make further coordinate transformations to obtain

$$ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

Taking the limit as the energy, $\mu \rightarrow 0$ gives The Kasner Solution

$$ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

Strange Minkowskian Line Element

- Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$

- Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

- Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

- Taking the limit as the energy, $\mu \rightarrow 0$ gives The **Kasner Solution**

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coord transforms that this is Kasner, but it won't be done here

2014-04-21

Strange Minkowskian Line Element

Start with the Schwarzschild solution

$$ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dt^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dr^2.$$

Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

Make further coordinate transformations to obtain

$$ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

Taking the limit as the energy, $\mu \rightarrow 0$ gives The **Kasner Solution**

$$ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

Strange Minkowskian Line Element

- Start with the Schwarzschild solution

$$\epsilon ds^2 = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) - \left(1 - \frac{2m}{r}\right) dt^2.$$

- Make the Eddington-Finkelstein coordinate transformation [2]

$$u = t - r - 2m \ln(r - 2m).$$

- Make further coordinate transformations to obtain

$$\epsilon ds^2 = \frac{r^2}{\cosh^2 \mu \xi} (d\xi^2 + d\eta^2) - 2du dr - \left(\mu^2 - \frac{2k}{r}\right) du^2.$$

- Taking the limit as the energy, $\mu \rightarrow 0$ gives The **Kasner Solution**

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2du dr - \frac{2k}{r} du^2.$$

- First we are going to derive a strange form of the Minkowskian line element.. of the vacuum field equations, which will be familiar to most of us
- to remove the coordinate singularity in the Schwarzschild solution
- These transformations put the line element in a form where we can take the limit as the energy goes to 0
- It is easily shown with further coord transforms that this is Kasner, but it won't be done here

2014-04-21

Strange Minkowskian Line Element

Then with $m = 0$ the strange Minkowskian line element is obtained
 $\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2dudr$.

- Then with $m = 0$ the strange Minkowskian line element is obtained

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2dudr.$$

- It is easily shown that $r = 0$ gives

$$\epsilon ds^2 = 0,$$

and thus is a single null geodesic.

- Its easily shown with suitable coordinate transforms that this is minkowskian line element
- This is best shown by calculating the geodesic equations after the Eddington-Finkelstein coord transforms, all zero if u is proper time along the geodesic

2014-04-21

Strange Minkowskian Line Element

Then with $m = 0$ the strange Minkowskian line element is obtained
 $\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2dudr$.
It is easily shown that $r = 0$ gives
 $\epsilon ds^2 = 0$,
and thus is a single null geodesic.

- Then with $m = 0$ the strange Minkowskian line element is obtained

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2dudr.$$

- It is easily shown that $r = 0$ gives

$$\epsilon ds^2 = 0,$$

and thus is a single null geodesic.

- Its easily shown with suitable coordinate transforms that this is minkowskian line element
- This is best shown by calculating the geodesic equations after the Eddington-Finkelstein coord transforms, all zero if u is proper time along the geodesic

2014-04-21

Strange Minkowskian Line Element

Then with $m = 0$ the strange Minkowskian line element is obtained
 $\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2dudr$.
It is easily shown that $r = 0$ gives
 $\epsilon ds^2 = 0$,
and thus is a single null geodesic.

- Then with $m = 0$ the strange Minkowskian line element is obtained

$$\epsilon ds^2 = r^2(d\xi^2 + d\eta^2) - 2dudr.$$

- It is easily shown that $r = 0$ gives

$$\epsilon ds^2 = 0,$$

and thus is a single null geodesic.

- Its easily shown with suitable coordinate transforms that this is minkowskian line element
- This is best shown by calculating the geodesic equations after the Eddington-Finkelstein coord transforms, all zero if u is proper time along the geodesic

2014-04-21

Layout

add in the contents

LTs that leave one null invariant direction are constructed

add in the contents

- Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2du dr.$$

- The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.
- In Cartesian coordinates this transformation becomes

$$\begin{aligned}x' + iy' &= x + iy + w(t - z), \\z' - t' &= -r = z - t, \\z' + t' &= z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Addition of complex numbers is commutative, and w has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

2014-04-21

Singular Lorentz Transformation

- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in cartesians
- Again its clear that $r = 0$ keeps one direction fixed, as then $z=t$
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular lorentz transformations

Singular Lorentz Transformation

- Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]
 $\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2du dr.$
- The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.
- In Cartesian coordinates this transformation becomes

$$\begin{aligned}x' + iy' &= x + iy + w(t - z), \\z' - t' &= -r = z - t, \\z' + t' &= z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).\end{aligned}$$

Addition of complex numbers is commutative, and w has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

- Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2dudr.$$

- The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.
- In Cartesian coordinates this transformation becomes

$$\begin{aligned}x' + iy' &= x + iy + w(t - z), \\z' - t' &= -r = z - t, \\z' + t' &= z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Addition of complex numbers is commutative, and w has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

2014-04-21

Singular Lorentz Transformation

Singular Lorentz Transformation

- Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]
 $\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2dudr$.
- The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.

the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in cartesians
- Again its clear that $r = 0$ keeps one direction fixed, as then $z=t$
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular lorentz transformations

- Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2du dr.$$

- The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.
- In Cartesian coordinates this transformation becomes

$$\begin{aligned}x' + iy' &= x + iy + w(t - z), \\z' - t' &= -r = z - t, \\z' + t' &= z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Addition of complex numbers is commutative, and w has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

2014-04-21

Singular Lorentz Transformation

Singular Lorentz Transformation

- Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]
 $\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2du dr.$
- The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.
- In Cartesian coordinates this transformation becomes
$$\begin{aligned}x' + iy' &= x + iy + w(t - z), \\z' - t' &= -r = z - t, \\z' + t' &= z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).\end{aligned}$$

the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in cartesians
- Again its clear that $r = 0$ keeps one direction fixed, as then $z=t$
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular lorentz transformations

- Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2dudr.$$

- The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.
- In Cartesian coordinates this transformation becomes

$$\begin{aligned}x' + iy' &= x + iy + w(t - z), \\z' - t' &= -r = z - t, \\z' + t' &= z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Addition of complex numbers is commutative, and w has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

2014-04-21

Singular Lorentz Transformation

Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]
 $\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2dudr$.

The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.

In Cartesian coordinates this transformation becomes

$$\begin{aligned}x' + iy' &= x + iy + w(t - z), \\z' - t' &= -r = z - t, \\z' + t' &= z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).\end{aligned}$$

Addition of complex numbers is commutative, and w has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in cartesians
- Again its clear that $r = 0$ keeps one direction fixed, as then $z=t$
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular lorentz transformations

- Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]

$$\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2dudr.$$

- The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.
- In Cartesian coordinates this transformation becomes

$$\begin{aligned}x' + iy' &= x + iy + w(t - z), \\z' - t' &= -r = z - t, \\z' + t' &= z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Addition of complex numbers is commutative, and w has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group

2014-04-21

Singular Lorentz Transformation

Singular Lorentz Transformation

- Define an arbitrary complex parameter $\zeta := \xi + i\eta$, to get the new line element[3]
 $\epsilon ds^2 = r^2 d\zeta d\bar{\zeta} - 2dudr$.
- The transformation $\zeta \rightarrow \zeta + w$, where $w \in \mathbb{C}$ is then trivial and leaves the single null geodesic $r = 0$ invariant.
- In Cartesian coordinates this transformation becomes
$$\begin{aligned}x' + iy' &= x + iy + w(t - z), \\z' - t' &= -r = z - t, \\z' + t' &= z + t + w(x - iy) + w(x + iy) + w\bar{w}(t - z).\end{aligned}$$
- Addition of complex numbers is commutative, and w has two parameters, so the singular Lorentz transformations form a 2-parameter abelian subgroup of the Lorentz group.

- This is what we want, An LT which leaves one null invariant.
- The use in the previous coord transforms was to make this transformation look trivial
- So this is what the seemingly trivial transformation looks like in cartesians
- Again its clear that $r = 0$ keeps one direction fixed, as then $z=t$
- but it doesn't work both ways, not all 2 parameter abelian subgroups are singular lorentz transformations

2014-04-21

Layout

add in the contents

add in the contents

- shown here that there is a 2 to 1 correspondence between $SL(2,C)$ and POLTs
- first show there 1 to 1 correspondence between points in Minkowskian space time and Hermitian matrices

$SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- It is also closely related to spinors

$$A = [t\mathbb{I}_2 - \vec{x} \cdot \vec{\sigma}].$$

- Construct the transformation $A(\vec{x}') = UA(\vec{x})U^\dagger$, where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of $SL(2, \mathbb{C})$

2014-04-21

$SL(2, \mathbb{C})$ Matrices of the POLT

- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- Where σ are the pauli matrices which form a basis for the Lie algebra of $SU(2)$
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

$SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- It is also closely related to spinors

$$A = [t\mathbb{I}_2 - \vec{x} \cdot \vec{\sigma}].$$

- Construct the transformation $A(\vec{x}') = UA(\vec{x})U^\dagger$, where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of $SL(2, \mathbb{C})$

2014-04-21

$SL(2, \mathbb{C})$ Matrices of the POLT

- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- Where σ are the pauli matrices which form a basis for the Lie algebra of $SU(2)$
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix}.$$

$SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- It is also closely related to spinors

$$A = [t\mathbb{I}_2 - \vec{x} \cdot \vec{\sigma}].$$

- Construct the transformation $A(\vec{x}') = UA(\vec{x})U^\dagger$, where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of $SL(2, \mathbb{C})$

$SL(2, \mathbb{C})$ Matrices of the POLT

- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- Where σ are the pauli matrices which form a basis for the Lie algebra of $SU(2)$
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

$SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- It is also closely related to spinors

$$A = [t\mathbb{I}_2 - \vec{x} \cdot \vec{\sigma}].$$

- Construct the transformation $A(\vec{x}') = UA(\vec{x})U^\dagger$, where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of $SL(2, \mathbb{C})$

$SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix
$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix}.$$
- This is useful as its determinant is the Lorentz quadratic form modulo a sign
$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$
- It is also closely related to spinors
$$A = [t\mathbb{I}_2 - \vec{x} \cdot \vec{\sigma}].$$

- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- Where σ are the pauli matrices which form a basis for the Lie algebra of $SU(2)$
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

$SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- It is also closely related to spinors

$$A = [t\mathbb{I}_2 - \vec{x} \cdot \vec{\sigma}].$$

- Construct the transformation $A(\vec{x}') = UA(\vec{x})U^\dagger$, where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of $SL(2, \mathbb{C})$

$SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix
$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix}.$$
- This is useful as its determinant is the Lorentz quadratic form modulo a sign
$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$
- It is also closely related to spinors
$$A = [t\mathbb{I}_2 - \vec{x} \cdot \vec{\sigma}].$$
- Construct the transformation $A(\vec{x}') = UA(\vec{x})U^\dagger$, where
$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$
is an element of $SL(2, \mathbb{C})$

- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- Where σ are the pauli matrices which form a basis for the Lie algebra of $SU(2)$
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

$SL(2, \mathbb{C})$ Matrices of the POLT

- There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices
- Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix},$$

- This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

- It is also closely related to spinors

$$A = [t\mathbb{I}_2 - \vec{x} \cdot \vec{\sigma}].$$

- Construct the transformation $A(\vec{x}') = UA(\vec{x})U^\dagger$, where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of $SL(2, \mathbb{C})$

$SL(2, \mathbb{C})$ Matrices of the POLT

There is a one to one correspondence between points in Minkowskian space-time and Hermitian matrices

Construct the following matrix

$$A = \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix}.$$

This is useful as its determinant is the Lorentz quadratic form modulo a sign

$$\det(A(\vec{x})) = t^2 - x^2 - y^2 - z^2.$$

It is also closely related to spinors

$$A = [t\mathbb{I}_2 - \vec{x} \cdot \vec{\sigma}].$$

Construct the transformation $A(\vec{x}') = UA(\vec{x})U^\dagger$, where

$$U = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

is an element of $SL(2, \mathbb{C})$

- Complex Hermitian matrices have 4 independent components, so the element of such a matrix can be used to represent points in Minkowskian space-time.
- Where σ are the pauli matrices which form a basis for the Lie algebra of $SU(2)$
- where $\alpha, \beta, \gamma, \delta$ are complex its an element of the special linear group. This means it has determinant 1. **write it on the board**

$SL(2, \mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$ and $A(\vec{x})$ have the same determinant so the above transformation preserves the Lorentz quadratic form, thus it is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t' - z' & x' + iy' \\ x' - iy' & t' + z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$

$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t - z)\bar{\alpha} + (x + iy)\bar{\beta} & (t - z)\bar{\gamma} + (x + iy)\bar{\delta} \\ (x - iy)\bar{\alpha} + (t + z)\bar{\beta} & (x - iy)\bar{\gamma} + (t + z)\bar{\delta} \end{pmatrix}.$$

- Thus the **general relations**

$$\begin{aligned} t' - z' &= (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta}, \\ x' + iy' &= (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta}, \\ t' + z' &= (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}. \end{aligned}$$

$SL(2, \mathbb{C})$ Matrices of the POLT

- This is because the determinant of U is 1

- I want to show you an example calculation of U, to do this we write in component form

$SL(2, \mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$ and $A(\vec{x})$ have the same determinant so the above transformation preserves the Lorentz quadratic form, thus it is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t' - z' & x' + iy' \\ x' - iy' & t' + z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$

$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t - z)\bar{\alpha} + (x + iy)\bar{\beta} & (t - z)\bar{\gamma} + (x + iy)\bar{\delta} \\ (x - iy)\bar{\alpha} + (t + z)\bar{\beta} & (x - iy)\bar{\gamma} + (t + z)\bar{\delta} \end{pmatrix}.$$

- Thus the **general relations**

$$\begin{aligned} t' - z' &= (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta}, \\ x' + iy' &= (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta}, \\ t' + z' &= (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}. \end{aligned}$$

$SL(2, \mathbb{C})$ Matrices of the POLT

- This is because the determinant of U is 1
- I want to show you an example calculation of U, to do this we write in component form

$SL(2, \mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$ and $A(\vec{x})$ have the same determinant so the above transformation preserves the Lorentz quadratic form, thus it is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t' - z' & x' + iy' \\ x' - iy' & t' + z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$

$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t - z)\bar{\alpha} + (x + iy)\bar{\beta} & (t - z)\bar{\gamma} + (x + iy)\bar{\delta} \\ (x - iy)\bar{\alpha} + (t + z)\bar{\beta} & (x - iy)\bar{\gamma} + (t + z)\bar{\delta} \end{pmatrix}.$$

• **general relations**

$$\begin{aligned} t' - z' &= (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta}, \\ x' + iy' &= (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta}, \\ t' + z' &= (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}. \end{aligned}$$

$SL(2, \mathbb{C})$ Matrices of the POLT

- $A(\vec{x}')$ and $A(\vec{x})$ have the same determinant so the above transformation preserves the Lorentz quadratic form, thus is a Lorentz transformation.
- Write this transformation component wise

$$\begin{pmatrix} t' - z' & x' + iy' \\ x' - iy' & t' + z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$

$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t - z)\bar{\alpha} + (x + iy)\bar{\beta} & (t - z)\bar{\gamma} + (x + iy)\bar{\delta} \\ (x - iy)\bar{\alpha} + (t + z)\bar{\beta} & (x - iy)\bar{\gamma} + (t + z)\bar{\delta} \end{pmatrix}.$$

- Thus the **general relations**

$$\begin{aligned} t' - z' &= (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta}, \\ x' + iy' &= (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta}, \\ t' + z' &= (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}. \end{aligned}$$

$SL(2, \mathbb{C})$ Matrices of the POLT

• $A(\vec{x}')$ and $A(\vec{x})$ have the same determinant so the above transformation preserves the Lorentz quadratic form, thus is a Lorentz transformation.

• Write this transformation component wise

$$\begin{pmatrix} t' - z' & x' + iy' \\ x' - iy' & t' + z' \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} t - z & x + iy \\ x - iy & t + z \end{pmatrix} \begin{pmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{pmatrix},$$

$$= \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} (t - z)\bar{\alpha} + (x + iy)\bar{\beta} & (t - z)\bar{\gamma} + (x + iy)\bar{\delta} \\ (x - iy)\bar{\alpha} + (t + z)\bar{\beta} & (x - iy)\bar{\gamma} + (t + z)\bar{\delta} \end{pmatrix}.$$

• Thus the **general relations**

$$\begin{aligned} t' - z' &= (t - z)\alpha\bar{\alpha} + (x + iy)\alpha\bar{\beta} + (x - iy)\beta\bar{\alpha} + (t + z)\beta\bar{\beta}, \\ x' + iy' &= (t - z)\alpha\bar{\gamma} + (x + iy)\alpha\bar{\delta} + (x - iy)\beta\bar{\gamma} + (t + z)\beta\bar{\delta}, \\ t' + z' &= (t - z)\gamma\bar{\gamma} + (x + iy)\gamma\bar{\delta} + (x - iy)\delta\bar{\gamma} + (t + z)\delta\bar{\delta}. \end{aligned}$$

- This is because the determinant of U is 1
- I want to show you an example calculation of U, to do this we write in component form

Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$\begin{aligned}t' - z' &= t - z, \\x' + iy' &= x + iy + w(t - z), \\t' + z' &= t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\begin{aligned}\alpha &= \pm 1, & \beta &= 0, \\ \gamma &= \bar{w}a, & \delta &= \alpha.\end{aligned}$$

- So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus **there is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs**

Example: Singular Lorentz transformation

- Where we have also used $\det(U) = 1$
- because the sign doesn't matter is still a solution Eqn(27)
- A, A' are points in Minkowskian space time, and $\pm U$ are POLTs

Take the singular Lorentz transformation from earlier

$$\begin{aligned}t' - z' &= t - z, \\x' + iy' &= x + iy + w(t - z), \\t' + z' &= t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).\end{aligned}$$

Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\begin{aligned}\alpha &= \pm 1, & \beta &= 0, \\ \gamma &= \bar{w}a, & \delta &= \alpha.\end{aligned}$$

So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

Thus there is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs

Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$\begin{aligned}t' - z' &= t - z, \\x' + iy' &= x + iy + w(t - z), \\t' + z' &= t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\begin{aligned}\alpha &= \pm 1, & \beta &= 0, \\ \gamma &= \bar{w}a, & \delta &= \alpha.\end{aligned}$$

- So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus there is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs

Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$\begin{aligned}t' - z' &= t - z, \\x' + iy' &= x + iy + w(t - z), \\t' + z' &= t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\begin{aligned}\alpha &= \pm 1, & \beta &= 0, \\ \gamma &= \bar{w}a, & \delta &= \alpha.\end{aligned}$$

- So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus there is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs

- Where we have also used $\det(U) = 1$
- because the sign doesn't matter is still a solution Eqn(27)
- A, A' are points in Minkowskian space time, and $\pm U$ are POLTs

Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$\begin{aligned}t' - z' &= t - z, \\x' + iy' &= x + iy + w(t - z), \\t' + z' &= t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\begin{aligned}\alpha &= \pm 1, & \beta &= 0, \\ \gamma &= \bar{w}a, & \delta &= \alpha.\end{aligned}$$

- So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus there is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs

Example: Singular Lorentz transformation

Take the singular Lorentz transformation from earlier

$$\begin{aligned}t' - z' &= t - z, \\x' + iy' &= x + iy + w(t - z), \\t' + z' &= t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).\end{aligned}$$

Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\begin{aligned}\alpha &= \pm 1, & \beta &= 0, \\ \gamma &= \bar{w}a, & \delta &= \alpha.\end{aligned}$$

So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

There is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs

- Where we have also used $\det(U) = 1$
- because the sign doesn't matter is still a solution Eqn(27)
- A, A' are points in Minkowskian space time, and $\pm U$ are POLTs

Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$\begin{aligned}t' - z' &= t - z, \\x' + iy' &= x + iy + w(t - z), \\t' + z' &= t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\begin{aligned}\alpha &= \pm 1, & \beta &= 0, \\ \gamma &= \bar{w}a, & \delta &= \alpha.\end{aligned}$$

- So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus **there is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs**

Example: Singular Lorentz transformation

Take the singular Lorentz transformation from earlier

$$\begin{aligned}t' - z' &= t - z, \\x' + iy' &= x + iy + w(t - z), \\t' + z' &= t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).\end{aligned}$$

Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\begin{aligned}\alpha &= \pm 1, & \beta &= 0, \\ \gamma &= \bar{w}a, & \delta &= \alpha.\end{aligned}$$

So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

Thus there is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs

- Where we have also used $\det(U) = 1$
- because the sign doesn't matter is still a solution Eqn(27)
- A, A' are points in Minkowskian space time, and $\pm U$ are POLTs

Example: Singular Lorentz transformation

- Take the singular Lorentz transformation from earlier

$$\begin{aligned}t' - z' &= t - z, \\x' + iy' &= x + iy + w(t - z), \\t' + z' &= t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).\end{aligned}$$

- Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\begin{aligned}\alpha &= \pm 1, & \beta &= 0, \\ \gamma &= \bar{w}a, & \delta &= \alpha.\end{aligned}$$

- So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

- Thus **there is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs**

Example: Singular Lorentz transformation

Take the singular Lorentz transformation from earlier

$$t' - z' = t - z,$$

$$x' + iy' = x + iy + w(t - z),$$

$$t' + z' = t + z + w(x - iy) + \bar{w}(x + iy) + w\bar{w}(t - z).$$

Equate coefficients on the RHS of this equation with the RHS of the general relations on the previous slide to obtain

$$\alpha = \pm 1, \quad \beta = 0,$$

$$\gamma = \bar{w}a, \quad \delta = \alpha.$$

So there are always **two** possible choices of U

$$U = \pm \begin{pmatrix} 1 & 0 \\ \bar{w} & 1 \end{pmatrix}$$

Thus there is a 2 to 1 correspondence between elements of $SL(2, \mathbb{C})$ and POLTs

- Where we have also used $\det(U) = 1$
- because the sign doesn't matter is still a solution Eqn(27)
- A, A' are points in Minkowskian space time, and $\pm U$ are POLTs

2014-04-21

temp

temp

temp

- temp

- temp

- 1 J.L. Synge - “Relativity: The Special Theory” - North Holland Publishing Company (1965)
- 2 D. Finkelstein - “Past-Future Asymmetry of the Gravitational Field of a Point Particle” - Phys.Rev.Vol 110, (1958) -
<http://journals.aps.org/pr/pdf/10.1103/PhysRev.110.965>
- 3 P.A. Hogan, C.Barrabès - “Advanced General Relativity: Gravity Waves, Spinning Particles and Black Holes” - Oxford University Press (May 2013)
- 4 I. Robinson “Spherical Gravitational Waves” - Phys.Rev.Lett. 4 (1960) 431-432 -
<http://journals.aps.org/prl/pdf/10.1103/PhysRevLett.4.431>
- 5 P.A Hogan, C. Barrabès - “Singular Null Hypersurfaces” - World Scientific Pub Co Inc (April 2004)
- 6 R. Penrose, W. Rindler - “Spinors and Space-Time: Volume 1, Two-Spinor Calculus and Relativistic Fields” - Cambridge University Press, (Feb 1987)
- 7 Tristan Needham - “Visual Complex Analysis” - Clarendon Press, Oxford (1997)
- 8 Various Authors - “Space-Time and Geometry: The Alfred Schild Lectures” - University of Texas Press (March 21, 2012)

2014-04-21

References

References

- 1 J.L. Synge - “Relativity: The Special Theory” - North Holland Publishing Company (1965)
- 2 D. Finkelstein - “Past-Future Asymmetry of the Gravitational Field of a Point Particle” - Phys.Rev.Vol 110, (1958) -
<http://journals.aps.org/pr/pdf/10.1103/PhysRev.110.965>
- 3 P.A. Hogan, C.Barrabès - “Advanced General Relativity: Gravity Waves, Spinning Particles and Black Holes” - Oxford University Press (May 2013)
- 4 I. Robinson “Spherical Gravitational Waves” - Phys.Rev.Lett. 4 (1960) 431-432 -
<http://journals.aps.org/prl/pdf/10.1103/PhysRevLett.4.431>
- 5 P.A Hogan, C. Barrabès - “Singular Null Hypersurfaces” - World Scientific Pub Co Inc (April 2004)
- 6 R. Penrose, W. Rindler - “Spinors and Space-Time: Volume 1, Two-Spinor Calculus and Relativistic Fields” - Cambridge University Press, (Feb 1987)
- 7 Tristan Needham - “Visual Complex Analysis” - Clarendon Press, Oxford (1997)
- 8 Various Authors - “Space-Time and Geometry: The Alfred Schild Lectures” - University of Texas Press (March 21, 2012)