

Table 14.2.3.5
Demographic Characteristics
Safety Population

Characteristic Category/Statistic	Control (N=#)	Study Drug 1 (N=#)	Study Drug 2 (N=#)
Age at Informed Consent			
N	##	##	##
Mean	##.##	##.##	##.##
Standard Deviation	##.##	##.##	##.##
Median	##.##	##.##	##.##
Min, Max	##, ##	##, ##	##, ##
Sex, n (%)			
Female	## (###.%)	## (###.%)	## (###.%)
Male	## (###.%)	## (###.%)	## (###.%)
Ethnicity, n (%)			
Hispanic or Latino	## (###.%)	## (###.%)	## (###.%)
Not Hispanic or Latino	## (###.%)	## (###.%)	## (###.%)
Race, n (%)			
White	## (###.%)	## (###.%)	## (###.%)
Black or African American	## (###.%)	## (###.%)	## (###.%)
Asian	## (###.%)	## (###.%)	## (###.%)
American Indian or Alaskan Native	## (###.%)	## (###.%)	## (###.%)
Native Hawaiian or Other Pacific Islander	## (###.%)	## (###.%)	## (###.%)
Multiple	## (###.%)	## (###.%)	## (###.%)
Age categories, n(%)			
< 65 years	## (###.%)	## (###.%)	## (###.%)
>= 65 years	## (###.%)	## (###.%)	## (###.%)
65 - < 75 years	## (###.%)	## (###.%)	## (###.%)
75 - < 85 years	## (###.%)	## (###.%)	## (###.%)
>= 85 years	## (###.%)	## (###.%)	## (###.%)
Missing	## (###.%)	## (###.%)	## (###.%)

Baseline is defined as the last measurement prior to the first dose of study drug.

%=100*n/N where n is the number of subjects in the specified category and N is the number of dosed subjects.

CONFIDENTIAL

Table 14.2.3.6
Disposition Table
Safety Population

Category	Control (N=#)	Study Drug 1 (N=#)	Study Drug 2 (N=#)	Total (N=#)
Screened	##	##	##	##
Randomized	##	##	##	##
Completed	## (##. #%)	## (##. #%)	## (##. #%)	## (##. #%)
Early Termination	## (##. #%)	## (##. #%)	## (##. #%)	## (##. #%)
Primary Reason for Early Termination				
Reason 1	## (##. #%)	## (##. #%)	## (##. #%)	## (##. #%)
Reason 2	## (##. #%)	## (##. #%)	## (##. #%)	## (##. #%)

Percentages are calculated with the number of Population subjects.

CONFIDENTIAL

Table 14.3.1.1
Overview of Adverse Events
Safety Population

Category	Control (N=#)	Study Drug 1 (N=#)		Study Drug 2 (N=#)	Total (N=#)
	n (%)	n (%)	n (%)	n (%)	n (%)
Subjects with any adverse event (AE)					# (###.%)
Subjects with any treatment-emergent AE (TEAE)	# (###.%)	# (###.%)	# (###.%)		# (###.%)
Maximum severity of TEAE					
Mild	# (###.%)	# (###.%)	# (###.%)		# (###.%)
Moderate	# (###.%)	# (###.%)	# (###.%)		# (###.%)
Severe	# (###.%)	# (###.%)	# (###.%)		# (###.%)
Subjects with any study drug related TEAE	# (###.%)	# (###.%)	# (###.%)		# (###.%)
Maximum severity of study drug related TEAE					
Mild	# (###.%)	# (###.%)	# (###.%)		# (###.%)
Moderate	# (###.%)	# (###.%)	# (###.%)		# (###.%)
Severe	# (###.%)	# (###.%)	# (###.%)		# (###.%)
Subjects with any serious AE (SAE)	# (###.%)	# (###.%)	# (###.%)		# (###.%)
Subjects with any AE leading to discontinuation of study drug	# (###.%)	# (###.%)	# (###.%)		# (###.%)

%=100*n/N, where n is the number of subjects in the specified category and N is the number of dosed subjects per column.
TEAEs are defined as any AE beginning on or after the first dose date of study drug.
Study drug related AEs include those that are possibly, probably, or definitely related.

<< PROGRAMMING NOTE: For TEAE, use TRTEMFL='Y'. >>

CONFIDENTIAL

Table 2.1.3
Summary of Treatment-Emergent Adverse Events by System Organ Class and Preferred Term
Safety Population

System Organ Class Preferred Term	Control (N=x)	Study Drug 1 (N=x)		Study Drug 2 (N=x)		Total (N=x)
	n (%)	n	(%)	n	(%)	n (%)
Subjects with any TEAE	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)

%=100*n/N, where n is the number of subjects in the specified category and N is the number of dosed subjects per column.
Treatment-emergent adverse events (TEAEs) are defined as any AE beginning on or after the first dose date of study drug.
AEs are summarized by the dose level at the onset of the event. At each level of summation (overall, system organ class, preferred term), subjects reporting more than one AE are counted only once. A subject may contribute to more than one preferred term. AEs are coded using MedDRA dictionary version 18.0.

<< Programming Note: Table will be sorted by descending frequency (n) in the total column for system organ class and preferred term within system organ class. Display all missing values as 0 using the mock shell's formatting. >>


Table 14.2.5.3
Change from Baseline for Vital Signs
Safety Population

Parameter	Visit	Statistics	Placebo (N=#)	Study Drug 1 (N=#)	Study Drug 2 (N=#)
Systolic Blood Pressure (SBP)	Week 2	N	##	##	##
		Mean (SD)	##.#(##.##)	##.#(##.##)	##.#(##.##)
		Median	##.##	##.##	##.##
		Q1, Q3	##.##,##.##	##.##,##.##	##.##,##.##
		Min, Max	##.##,##.##	##.##,##.##	##.##,##.##
	Week 4	N	##	##	##
		Mean (SD)	##.#(##.##)	##.#(##.##)	##.#(##.##)
		Median	##.##	##.##	##.##
		Q1, Q3	##.##,##.##	##.##,##.##	##.##,##.##
		Min, Max	##.##,##.##	##.##,##.##	##.##,##.##
	• • •				
	Week 24	N	##	##	##
		Mean (SD)	##.#(##.##)	##.#(##.##)	##.#(##.##)
		Median	##.##	##.##	##.##
		Q1, Q3	##.##,##.##	##.##,##.##	##.##,##.##
		Min, Max	##.##,##.##	##.##,##.##	##.##,##.##

N = number of subjects per treatment group; SD = Standard Deviation; Change from Baseline = [post-baseline value] - [Baseline value]

<< Programming Note: Consider all “PARAM” values in “Parameter” column; Inside “ATPT” variable containing 3 different measurements per visit consider only “After Standing for 3 Minutes”; Don’t consider “Height” in “Parameter” column; For treatment related variable consider “TRTAN”, “TRTA”. >>


Figure 14.2.1
Kaplan Meier Curve: Overall Survival by Treatment Group
ITT Population



Subjects without a death event are censored at the last date known to be alive.
Kaplan-Meier estimates are calculated using the product-limit method.

<<Programming Note: Use for input data, use adtteos. For Overall Survival, Use PARAMCD="OS" (or OS dataset).
Use USUBJID. CNSR=0 for EVNTDESC in ("DEATH"), else 1. >>


Figure 14.2.4.2
Kaplan Meier Curve: Progression Free Survival by Treatment Group
ITT Population



Progression-Free Survival (PFS) is defined as the time from randomization to disease progression or death from any cause, whichever occurs first.

<< Programming Note: Use for input data, use adttepfs. For Progression Free Survival, Use PARAMCD="PFS" (or PFS dataset). Use SUBJID, not USUBJID. CNSR=0 for EVNTDESC in ("DEATH","PROGRESSIVE DISEASE"), else 1. >>


Figure 14.1.3.2
Change in Diastolic Blood Pressure from Baseline Over Visits
Safety Population



Baseline diastolic blood pressure is defined as the last non-missing measurement prior to the first dose.
Visits with insufficient data may not be displayed on the figure.

<< Programming Note: Use for input data, use advs. Keep only analysis flag 1 = Y. >>

Figure 14.1.6.3
Change in Systolic Blood Pressure from Baseline Over Visits
Safety Population



Baseline systolic blood pressure is defined as the last non-missing measurement prior to the first dose.
Change from baseline is calculated as post-baseline systolic blood pressure minus the baseline value.

<< Programming Note: Use for input data, use advs. Keep only analysis flag 1 = Y. >>