

Table 14.2.3.5
Demographic Characteristics
Safety Population

Characteristic Category/Statistic	Placebo (N=#)	Study Drug 1 (N=#)	Study Drug 2 (N=#)
Age at Informed Consent			
N	##	##	##
Mean	##.##	##.##	##.##
Standard Deviation	##.##	##.##	##.##
Median	##.##	##.##	##.##
Min, Max	##, ##	##, ##	##, ##
Sex, n (%)			
Female	## (###.%)	## (###.%)	## (###.%)
Male	## (###.%)	## (###.%)	## (###.%)
Ethnicity, n (%)			
Hispanic or Latino	## (###.%)	## (###.%)	## (###.%)
Not Hispanic or Latino	## (###.%)	## (###.%)	## (###.%)
Race, n (%)			
White	## (###.%)	## (###.%)	## (###.%)
Black or African American	## (###.%)	## (###.%)	## (###.%)
Asian	## (###.%)	## (###.%)	## (###.%)
American Indian or Alaskan Native	## (###.%)	## (###.%)	## (###.%)
Native Hawaiian or Other Pacific Islander	## (###.%)	## (###.%)	## (###.%)
Multiple	## (###.%)	## (###.%)	## (###.%)
Age categories, n(%)			
< 65 years	## (###.%)	## (###.%)	## (###.%)
>= 65 years	## (###.%)	## (###.%)	## (###.%)
65 - < 75 years	## (###.%)	## (###.%)	## (###.%)
75 - < 85 years	## (###.%)	## (###.%)	## (###.%)
>= 85 years	## (###.%)	## (###.%)	## (###.%)
Missing	## (###.%)	## (###.%)	## (###.%)

Baseline is defined as the last measurement prior to the first dose of study drug.

%=100*n/N where n is the number of subjects in the specified category and N is the number of dosed subjects.

CONFIDENTIAL

Table 14.2.3.6
Disposition Table
Safety Population

Category	Placebo (N=)	Study Drug 1 (N=)	Study Drug 2 (N=)	Total (N=)
Screened	##	##	##	##
Randomized	##	##	##	##
Completed	## (##. #%)	## (##. #%)	## (##. #%)	## (##. #%)
Early Termination	## (##. #%)	## (##. #%)	## (##. #%)	## (##. #%)
Primary Reason for Early Termination				
Reason 1	## (##. #%)	## (##. #%)	## (##. #%)	## (##. #%)
Reason 2	## (##. #%)	## (##. #%)	## (##. #%)	## (##. #%)

Percentages are calculated with the number of Population subjects.

Table 14.3.1.1
Overview of Adverse Events
Safety Population

Category	Placebo (N=#)	Study Drug 1 (N=#)	Study Drug 2 (N=#)	Total (N=#)
	n (%)	n (%)	n (%)	n (%)
Subjects with any adverse event (AE)				# (###.%)
Subjects with any treatment-emergent AE (TEAE)	# (###.%)	# (###.%)	# (###.%)	# (###.%)
Maximum severity of TEAE				
Mild	# (###.%)	# (###.%)	# (###.%)	# (###.%)
Moderate	# (###.%)	# (###.%)	# (###.%)	# (###.%)
Severe	# (###.%)	# (###.%)	# (###.%)	# (###.%)
Subjects with any study drug related TEAE	# (###.%)	# (###.%)	# (###.%)	# (###.%)
Maximum severity of study drug related TEAE				
Mild	# (###.%)	# (###.%)	# (###.%)	# (###.%)
Moderate	# (###.%)	# (###.%)	# (###.%)	# (###.%)
Severe	# (###.%)	# (###.%)	# (###.%)	# (###.%)
Subjects with any serious AE (SAE)	# (###.%)	# (###.%)	# (###.%)	# (###.%)
Subjects with any AE leading to discontinuation of study drug	# (###.%)	# (###.%)	# (###.%)	# (###.%)

%=100*n/N, where n is the number of subjects in the specified category and N is the number of dosed subjects per column.

TEAEs are defined as any AE beginning on or after the first dose date of study drug.

Study drug related AEs include those that are possibly, probably, or definitely related.

<< PROGRAMMING NOTE: For TEAE, use TRTEMFL='Y'. >>

Table 2.1.3
Summary of Treatment-Emergent Adverse Events by System Organ Class and Preferred Term
Safety Population

System Organ Class Preferred Term	Placebo (N=x)	Study Drug 1 (N=x)		Study Drug 2 (N=x)		Total (N=x)
	n (%)	n	(%)	n	(%)	n (%)
Subjects with any TEAE	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	x (xxx.x)	x	(xxx.x)	x	(xxx.x)	x (xxx.x)

%=100*n/N, where n is the number of subjects in the specified category and N is the number of dosed subjects per column.
Treatment-emergent adverse events (TEAEs) are defined as any AE beginning on or after the first dose date of study drug.
AEs are summarized by the dose level at the onset of the event. At each level of summation (overall, system organ class, preferred term), subjects reporting more than one AE are counted only once. A subject may contribute to more than one preferred term. AEs are coded using MedDRA dictionary version 18.0.

<< Programming Note: Table will be sorted by descending frequency (n) in the total column for system organ class and preferred term within system organ class; Display all missing values as 0 using the mock shell's formatting. >>

Table 14.2.5.3
Change from Baseline for Vital Signs
Safety Population

Parameter	Visit	Statistics	Placebo (N=#)	Study Drug 1 (N=#)	Study Drug 2 (N=#)
Systolic Blood Pressure (SBP)	Week 2	N	##	##	##
		Mean (SD)	##.#(##.#)	##.#(##.#)	##.#(##.#)
		Median	##.#	##.#	##.#
		Q1, Q3	##.#,##.#	##.#,##.#	##.#,##.#
		Min, Max	##.#,##.#	##.#,##.#	##.#,##.#
	Week 4	N	##	##	##
		Mean (SD)	##.#(##.#)	##.#(##.#)	##.#(##.#)
		Median	##.#	##.#	##.#
		Q1, Q3	##.#,##.#	##.#,##.#	##.#,##.#
		Min, Max	##.#,##.#	##.#,##.#	##.#,##.#
	Week 24	N	##	##	##
		Mean (SD)	##.#(##.#)	##.#(##.#)	##.#(##.#)
		Median	##.#	##.#	##.#
		Q1, Q3	##.#,##.#	##.#,##.#	##.#,##.#
		Min, Max	##.#,##.#	##.#,##.#	##.#,##.#

N = number of subjects per treatment group; SD = Standard Deviation; Change from Baseline = [post-baseline value] - [Baseline value]

<< Programming Note: Consider all "PARAM" values in "Parameter" column; Inside "ATPT" variable containing 3 different measurements per visit consider only "After Standing for 3 Minutes"; Don't consider "Height" in "Parameter" column; For treatment related variable consider "TRTAN"," TRTA". >>

Table 14.2.2
Progression Free Survival
ITT Population

	Placebo (N=#)	Study Drug 1 (N=#)	Study Drug 2 (N=#)
Number of Patients	##	##	##
Number of Death Observed	##	##	##
Number of Censoring	##	##	##
Number of Censoring (%)	##.##	##.##	##.##
Median Survival (Days)	##.##	##.##	##.##
95% of Median Survival (Days)	(##.##, ##.##)	(##.##, ##.##)	(##.##, ##.##)
Log-Rank Test P Value		##.##	##.##
Hazard Ratio		##.##	##.##

<< Programming Note: Use adttepfs.xpt datasets; Use "TRTP" for treatment group labels and "TRTPN" for treatment group numeric codes. >>

Table 3.1.2
Summary of Concomitant Medications by Preferred Terms
Safety Population

Preferred Term	Placebo (N=x) n (%)	Study Drug 1 (N=x) n (%)	Study Drug 2 (N=x) n (%)	Total (N=x) n (%)
Subjects with any Concomitant Medication	x (xxx.x)	x (xxx.x)	x (xxx.x)	x (xxx.x)
xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx	x (xxx.x)	x (xxx.x)	x (xxx.x)	x (xxx.x)
xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx	x (xxx.x)	x (xxx.x)	x (xxx.x)	x (xxx.x)
xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx	x (xxx.x)	x (xxx.x)	x (xxx.x)	x (xxx.x)
xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx	x (xxx.x)	x (xxx.x)	x (xxx.x)	x (xxx.x)
xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx	x (xxx.x)	x (xxx.x)	x (xxx.x)	x (xxx.x)
xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx	x (xxx.x)	x (xxx.x)	x (xxx.x)	x (xxx.x)

%=100*n/N, where n is the number of subjects in the specified category and N is the number of dosed subjects per column.

<< Programming Note: Table will be sorted by descending frequency (n) in the total column for preferred term. >>

Table 4.1.3.1
ANCOVA of Change from Baseline Composite Score at Week 24 (LOCF)
ITT Population

	Baseline		Week 24 (LOCF)		Change from baseline		
Treatment Group	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	LS Mean (95% CI)
Placebo	xxx	x.x(x.xx)	xxx	x.x(x.xx)	xxx	x.x(x.xx)	x.x(x.xx, x.xx)
Xanomeline Low Dose	xxx	x.x(x.xx)	xxx	x.x(x.xx)	xxx	x.x(x.xx)	x.x(x.xx, x.xx)
Xanomeline High Dose	xxx	x.x(x.xx)	xxx	x.x(x.xx)	xxx	x.x(x.xx)	x.x(x.xx, x.xx)

ANCOVA = Analysis of Covariance; CI = Confidence Interval; LS = Least Squares; SD = Standard Deviation

Source: ADQS

<< Programming Notes: LS means and differences in LS means are based on an ANCOVA model with treatment and baseline glucose as covariates. LOCF (Last Observation Carried Forward) approach is used for missing Week 24 values; For treatment related variables consider “TRT01P” and “TRT01PN”. >>

Table 4.1.3.2
ANCOVA of Change from Baseline Composite Score at Week 24 (LOCF)-Pairwise Comparison
ITT Population

Pairwise Comparison	Difference in LS Mean (95% CI)	p-Value
Xanomeline Low Dose - Placebo	xx.xx (x.xx, x.xx)	x.xxx
Xanomeline High Dose - Placebo	xx.xx (x.xx, x.xx)	x.xxx


Source: ADQS

ANCOVA = Analysis of Covariance; CI = Confidence Interval; LS = Least Squares

<< Programming Notes:

1. LS means and differences in LS means are based on an ANCOVA model with treatment and baseline glucose as covariates. p-values are from the ANCOVA model testing treatment effects LOCF (Last Observation Carried Forward) approach is used for missing Week 24 values.
 2. Least Squares (LS) means, differences in LS means, 95% confidence intervals, and p-values were derived from PROC Mixed.
 3. *Pairwise comparisons were performed versus placebo using the DIFF=CONTROL option.*
 4. *Use Treatment variables "TRT01A" and "TRT01AN"*
- >>


Figure 14.2.1
Kaplan Meier Curve: Overall Survival by Treatment Group
ITT Population



Subjects without a death event are censored at the last date known to be alive.
Kaplan-Meier estimates are calculated using the product-limit method.

<<Programming Note: Use for input data, use adtteos. For Overall Survival, Use PARAMCD="OS" (or OS dataset).
Use USUBJID. CNSR=0 for EVNTDESC in ("DEATH"), else 1. >>


Figure 14.2.4.2
Kaplan Meier Curve: Progression Free Survival by Treatment Group
ITT Population



Progression-Free Survival (PFS) is defined as the time from randomization to disease progression or death from any cause, whichever occurs first.

<< Programming Note: Use for input data, use adttepfs. For Progression Free Survival, Use PARAMCD="PFS" (or PFS dataset). Use SUBJID, not USUBJID. CNSR=0 for EVNTDESC in ("DEATH","PROGRESSIVE DISEASE"), else 1. >>


Figure 14.1.3.2
Change in Diastolic Blood Pressure from Baseline Over Visits
Safety Population



Baseline diastolic blood pressure is defined as the last non-missing measurement prior to the first dose.
Visits with insufficient data may not be displayed on the figure.

<< Programming Note: Use for input data, use advs. Keep only analysis flag 1 = Y. >>


Figure 14.1.6.3
Change in Systolic Blood Pressure from Baseline Over Visits
Safety Population



Baseline systolic blood pressure is defined as the last non-missing measurement prior to the first dose.
Change from baseline is calculated as post-baseline systolic blood pressure minus the baseline value.

<< Programming Note: Use for input data, use advs. Keep only analysis flag 1 = Y. >>

Figure 14.1.3.6
Lymphocyte Count Plotting Over Visits for Subject
Safety Population



<< Programming Note: For Lymphocyte Count, use adlbh.xpt dataset, where PARAMCD="LYM". >>

Listing 16.2.9.3
Demographic
Safety Population

Subject	Age (Years)	Sex	Ethnicity	Race
###-##	##	XXXX	XXXXXXXXXXXXX	XXXXXXXXXXXXX
###-##	##	XXXX	XXXXXXXXXXXXX	XXXXXXXXXXXXX
###-##	##	XXXX	XXXXXXXXXXXXX	XXXXXXXXXXXXX
###-##	##	XXXX	XXXXXXXXXXXXX	XXXXXXXXXXXXX
###-##	##	XXXX	XXXXXXXXXXXXX	XXXXXXXXXXXXX
###-##	##	XXXX	XXXXXXXXXXXXX	XXXXXXXXXXXXX
###-##	##	XXXX	XXXXXXXXXXXXX	XXXXXXXXXXXXX
###-##	##	XXXX	XXXXXXXXXXXXX	XXXXXXXXXXXXX

Baseline values (e.g., height, weight, BMI) are defined as the last non-missing assessments obtained prior to the first vaccination (Day 1).

<< Programming note: Use dm (XPT dataset) dataset. >>

Listing 16.2.9.3
Disposition Data
Safety Population

Subject	Date of Discontinuation / Completion	Reason for Discontinuation
###-### ###-### ###-### ###-### ###-### ###-### ###-### ###-###	DDMMYYYY DDMMYYYY DDMMYYYY DDMMYYYY DDMMYYYY	XXXXXXXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXXXXXXXXXXX

Disposition information is presented as collected in the study database.
Subjects are listed once based on their unique subject identifier.

Program Name: XXXXXXXX.sas Run Date: DDMMYYYY HH:

<< Programming Note: Use ADSL(.xpt) dataset. ADSL.RFENDT = "Date of Discontinuation/Completion"; ADSL.DCREASCD = "Reason for Discontinuation".
>>

Listing 16.2.8.1
Safety Laboratory Values: Chemistry
Safety Population

Subject	Parameter (Unit)	Visit	Date/Time of Collection	Result	Normal Range	Flag [1]
###-### (SF)	XXXXXXXXXX (XXXXX)	XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
	XXXXXXXXXX (XXXXX)	XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
###-###	XXXXXXXXXX (XXXXX)	XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
###-###	XXXXXXXXXX (XXXXX)	XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX
		XXXXXXXXXXXX	DDMMYYYY/HH:MM	XXXXXX	XX - XX	XX

SF = Screen failure subject.

[1] Laboratory Flag: RH=Reference High, RL=Reference Low, NH=Notable High, NL=Notable Low, CH=Critical High, CL=Critical Low

<< Programming Note: Including all non-missing LBORRES where LBCAT = "CHEMISTRY"; Exclude LBTESTCD = "HCG" since that is a pregnancy test which is included in another listing. >>

Listing 16.2.9.3
Vital Signs
Safety Population

Subject	Visit	Date Performed	Time Point	SBP (mmHg)	DBP (mmHg)	HR (bpm)	Temperature (°C)	RR (breaths/min)	Weight (kg)	Height (cm)
###-###	XXXXXXXXXXXX	DDMMYYYY	hh:mm:ss	###	###	##	###	###	###.#	###.#
	XXXXXXXXXXXX	DDMMYYYY	hh:mm:ss	###	###	##	###	###	###.#	###.#
	XXXXXXXXXXXX	DDMMYYYY	hh:mm:ss	###	###	##	###	###	###.#	###.#
###-###	XXXXXXXXXXXX	DDMMYYYY	hh:mm:ss	###	###	##	###	###	###.#	###.#
	XXXXXXXXXXXX	DDMMYYYY	hh:mm:ss	###	###	##	###	###	###.#	###.#
	XXXXXXXXXXXX	DDMMYYYY	hh:mm:ss	###	###	##	###	###	###.#	###.#
###-###	XXXXXXXXXXXX	DDMMYYYY	hh:mm:ss	###	###	##	###	###	###.#	###.#
	XXXXXXXXXXXX	DDMMYYYY	hh:mm:ss	###	###	##	###	###	###.#	###.#
	XXXXXXXXXXXX	DDMMYYYY	hh:mm:ss	###	###	##	###	###	###.#	###.#

SBP = systolic blood pressure. DBP = diastolic blood pressure. RR = respiratory rate (breaths per minute). HR = Heart Rate (Beats per minute)

<< Programming Note: Use VS(.xpt) dataset; Respiratory Rate (RR) - RESP variable. >>