








D ranked contributions

rank	residue (ligand->protein)	share	atom (protein->ligand)	share
1	Q92	0.1056	C24	0.2458
2	W191	0.0051	C11	0.0529
3	S172	0.0049	C18	0.0528
4	V162	0.0048	N13	0.0515
5	W208	0.0046	N1	0.0474
6	G81	0.0045	C2	0.0454
7	N67	0.0045	O15	0.0389
8	G25	0.0045	O14	0.0389
9	D75	0.0043	C30	0.0351
10	K80	0.0042	C22	0.0351

predicted pKd 9.09 / pKi 7.81 / pIC50 7.45 | measured pKi 7.22, error +0.59 log units selection: the median-potency observation of each of the most-measured distinct targets; one pair per distinct target; sequences of at most 500 residues; drawn from 273,992 observations in the val partition
non-physical token mass: 0.0000 (0.00% of the total) - knowledge and virtual nodes map to no atom and are excluded from panels B-C
shares are fractions of the summed physical attention (260.0000)

N#CCCN(CCc1ccc(S(N)(=O)=O)cc1)C(=O)c1ccc(-c2cssc2=S)cc1 | ligand_gat + esmc | checkpoint runs\random_pair-gat-frozen-v1\best.pt
aggregation: last layer, mean over heads, bond contributions split equally between endpoint atoms (derived aggregation)
[a69d0443fbd69c13]
Attention-based explanation. NOT an experimentally established contact map, binding pose, or mechanism. The residue x atom view is a derived aggregation.