


A ligand-to-protein attention per residue over all 460 residues (shaded: panel B window 1-200)



B residue x atom - DERIVED AGGREGATION



C per-atom contribution (dark = low, yellow = high)



D ranked contributions

rank	residue (ligand->protein)	share	atom (protein->ligand)	share
1	I53	0.0026	C12	0.1219
2	Y413	0.0025	C1	0.0774
3	I380	0.0025	C30	0.0512
4	C289	0.0025	C8	0.0396
5	T223	0.0025	C17	0.0347
6	S434	0.0025	C24	0.0340
7	S423	0.0025	C18	0.0329
8	I299	0.0025	O2	0.0325
9	N419	0.0025	N32	0.0316
10	I410	0.0025	C19	0.0316

predicted pKd 7.47 / pKi 8.08 / pIC50 7.89 | measured pIC50 7.92, error -0.04 log units selection: the median-potency observation of each of the most-measured distinct targets; one pair per distinct target; sequences of at most 500 residues; drawn from 273,992 observations in the val partition
non-physical token mass: 0.0000 (0.00% of the total) - knowledge and virtual nodes map to no atom and are excluded from panels B-C
shares are fractions of the summed physical attention (460.0000)

COC(=O)N1CCC(C#Cc2cnc(OC[C@@H]3CCC(=O)N3)c3cc(OC(C)C)c(C(N)=O)cc23)CC1 | ligand_gat + esmc | checkpoint runs\random_pair-gat-frozen-v1\best.pt
aggregation: last layer, mean over heads, bond contributions split equally between endpoint atoms (derived aggregation)
[a69d0443fbd69c13]
Attention-based explanation. NOT an experimentally established contact map, binding pose, or mechanism. The residue x atom view is a derived aggregation.