






C per-atom contribution (dark = low, yellow = high)



D ranked contributions

rank	residue (ligand->protein)	share	atom (protein->ligand)	share
1	K58	0.0024	C27	0.1290
2	I384	0.0024	N8	0.0773
3	K50	0.0024	C26	0.0766
4	V157	0.0024	C9	0.0729
5	V122	0.0024	O10	0.0717
6	G97	0.0024	N16	0.0626
7	L320	0.0023	N28	0.0479
8	R93	0.0023	C7	0.0443
9	K165	0.0023	C6	0.0392
10	V11	0.0023	O18	0.0381

predicted pKd 6.72 / pKi 6.94 / pIC50 6.81 | measured pIC50 6.58, error +0.23 log units selection: the median-potency observation of each of the most-measured distinct targets; one pair per distinct target; sequences of at most 500 residues; drawn from 273,992 observations in the val partition
non-physical token mass: 0.0000 (0.00% of the total) - knowledge and virtual nodes map to no atom and are excluded from panels B-C
shares are fractions of the summed physical attention (482.0000)

Nc1ccccc1NC(=O)CCCCCNC(=O)c1cc2ccccc2cn1 | ligand_gat + esmc | checkpoint runs\random_pair-gat-frozen-v1\best.pt
aggregation: last layer, mean over heads, bond contributions split equally between endpoint atoms (derived aggregation)
[a69d0443fbd69c13]
Attention-based explanation. NOT an experimentally established contact map, binding pose, or mechanism. The residue x atom view is a derived aggregation.