

NRES 746 – Workout of the Day #4: Limits and a Derivative

Name: _____

Instructions: Work alone for about 5-8 minutes. Then compare with a neighbor.

Part 1: What happens at the edges?

For each function below, use the “throw out the lower-order terms” trick from Monday’s lecture to state what happens as $x \rightarrow 0$ and as $x \rightarrow \infty$. You don’t need to show work – just the two limiting behaviors for each.

1a. Michaelis-Menten: $f(x) = \frac{ax}{b+x}$

As $x \rightarrow 0$: _____

As $x \rightarrow \infty$: _____

1b. Ricker: $f(x) = ax e^{-bx}$

As $x \rightarrow 0$: _____

As $x \rightarrow \infty$: _____

1c. Logistic: $f(x) = \frac{1}{1+e^{-bx}}$

As $x \rightarrow 0$: _____

As $x \rightarrow \infty$: _____

1d. Holling type III: $f(x) = \frac{ax^2}{b^2+x^2}$

As $x \rightarrow 0$: _____

As $x \rightarrow \infty$: _____

Part 2: What kind of point is $x = 0$?

The logistic function, $f(x) = \frac{1}{1+e^{-bx}}$, is the simplest sigmoid in the bestiary – it’s symmetric around $x = 0$.

2a. Rewrite $f(x)$ as $(1+e^{-bx})^{-1}$, then find $f'(x)$ using the chain rule (twice: once for the outer power, once for the inner exponential).

Reminders: $\frac{d}{dx} u^n = n u^{n-1} u'$ $\frac{d}{dx} e^{g(x)} = g'(x) e^{g(x)}$

Show your work:

2b. Evaluate $f'(x)$ at $x = 0$.

$f'(0) =$ _____

2c. In Lab 1 you found that the Ricker function's derivative is zero at its peak – a critical point. Is $f'(0)$ also zero here, at the logistic's midpoint? If not, what kind of special point is $x = 0$ instead, and what does the actual value of $f'(0)$ tell you about the curve there? (Hint: think about the curve's concavity just before and just after $x = 0$ – what would that imply about the *second* derivative at $x = 0$?)
