Week 10, Lecture 2: Bayesian Model Comparison and Shrinkage Priors

Instructor notes. Part 1: prep reading. Part 2: podium cheat sheet

How to prepare (for students)

- **Big picture.** Monday compared models with likelihood ratio tests, AIC, and penalties on coefficient size (ridge, lasso). Today: the Bayesian versions. How do we compare models fit by MCMC, where there is no single best-fit parameter set, and how do priors do the job of a penalty?
- **Read:** Hooten and Hobbs (2015), *A guide to Bayesian model selection for ecologists* (Ecological Monographs 85: 3-28), the sections "Model validation," "Statistical regularization and information criteria," and "Guidance"; skim the rest. Bolker 6.6.3 (printed p. 280-284) on Bayes factors and DIC.
- **Review:** Bayes' rule for parameters (posterior $\propto$ likelihood $\times$ prior); Monday's ridge and lasso penalties; fitting a model with `cmdstanr` (last week's take-home).

Overview (instructor only)

Beat	Mode	Time
1. Big picture	BOARD	~3 min
2. WOD: *Ridge Is a Prior*	handout	~10 min
3. WAIC and LOO	BOARD $\rightarrow$ R	~12 min
4. Shrinkage priors	BOARD $\rightarrow$ R	~12 min
5. Discussion	BOARD	~5 min
If time: Bayes factors, RJMCMC	BOARD	

- R = Wednesday section of `CODE_Week10_ModelComparison.R` (needs Monday's section run first). Hand out the take-home WOD `WOD_Week10_Lecture2_PondModelsWithLOO.Rmd` at the end (due Monday).
- **Behind?** Skip the DIC row; in section 4 show only the interval plot and its pause.
- **Built before Monday was taught;** check what Monday and Tuesday covered and adjust.

Part 1: Lecture content

1. Big picture (~3 min) [BOARD]

- COMPARE (2), Bayesian side. Monday: penalties on count and size. Today: the same two ideas for MCMC fits
- No single $\hat{\theta} \Rightarrow$ no AIC; priors blur "number of parameters"

2. WOD reveal (~10 min) [handout]

- $-\log P(\mu \mid y) = \sum(y_i - \mu)^2/(2\sigma^2) + \mu^2/(2\tau^2)$, so $\lambda = 1/(2\tau^2)$
- Ridge estimate = posterior mode; vague prior ($\tau \to \infty$) = MLE
- Lasso $\lambda|\mu|$ = Laplace prior: spike at 0, heavier tails

3. WAIC and LOO (~12 min) [BOARD → R]


- Score = **elpd**: for each data point, log of its likelihood averaged over the posterior; sum. Higher is better; IC = $-2\times$ elpd
- Penalty = effective parameters, estimated from the posterior:
- DIC (Bolker): p_D = mean deviance − deviance at posterior means
- WAIC: p_{waic} = summed posterior variance of each point's log lik.
- LOO: approximates refitting without each point (PSIS); Pareto $k > 0.7$ flags points where the shortcut fails
- Stan must save `log_lik[i]` per observation (generated quantities)

	no wave	full	effective parameters
AIC	2278.0	2282.0	k = 3 vs. 6
DIC	2278.0	2281.9	pD = 3.0 vs. 5.9
WAIC	2278.6	2282.7	p_waic = 3.5 vs. 6.5
LOOIC	2278.6	2282.7	p_loo = 3.5 vs. 6.5


- Fir: elpd difference -2.1 (SE 1.8); all Pareto k good
- **Pause:** full model 2.1 behind, SE 1.8. Conclusion? *Land:* about one SE; can't tell the models apart (not "no wave effect")
- **Sticking point:** AIC gives no SE for a difference; LOO does

4. Shrinkage priors (~12 min) [BOARD → R]

- Prior scale plays λ's role: Normal = ridge, Laplace = lasso
- Creepers (Monday), three priors on the 10 slopes:



- Northness (truth 0): vague 0.23 (-0.02 to 0.49); Laplace 0.08 (-0.07 to 0.28)
- p_{loo}: vague 9.1, Normal 7.8, Laplace 7.1 (10 slopes + intercept): tighter priors, fewer effective parameters
- LOO best: laplace; others within 1.5 elpd (SE up to 1.6). 3 stations have $k > 0.7$ (mention only)
- **Pause:** the lasso set northness to exactly 0; the Laplace posterior says 0.08 (-0.07 to 0.28). Which is more honest? *Land:* the posterior keeps the uncertainty; a Bayesian lasso never gives exact zeros
- Horseshoe: tall spike plus heavy tails (small effects shrink hard, big ones barely). Sketch only:



- **Sticking point:** choosing the prior scale = choosing λ; compare by LOO, or estimate

it (a hierarchical prior, later)

5. Discussion (~5 min) [BOARD]

Q1. LOO prefers the no-wave fir model by 2.1 elpd (SE 1.8). A colleague writes "wave regeneration has no effect on cone production." Agree?

Q2. With a Laplace prior, northness gets a posterior median of 0.08 (90% interval -0.07 to 0.28); Monday's lasso said exactly 0. Which would you report, and why?

- Q1 lands on: a difference within ~1 SE is inconclusive; "not detected" $\neq$ "no effect"; corroborated, not proven
- Q2 lands on: the posterior shows the uncertainty a single zero hides; for prediction either works, for inference report the interval

If time: Bayes factors and selection inside the sampler [BOARD]

- Bayes factor = ratio of marginal likelihoods (averaged over the prior); sensitive to vague priors, hard to compute (Bolker 6.6.3)
- RJMCMC / spike-and-slab: the sampler jumps between models; posterior model probabilities. Needs discrete parameters, so not Stan (JAGS, NIMBLE); legacy LECTURE8 has BUGS code

PART 2: PODIUM CHEAT SHEET

Terse on purpose. Detail: Part 1. R = Wednesday section of CODE_Week10_ModelComparison.R.

Big picture → WOD → lecture. Take-home WOD out at end. Behind? Skip DIC; sec. 4 plot + pause only.

1. Big picture (~3 min) [BOARD]

- COMPARE (2), Bayesian side
- Monday: count and size penalties. Today: both for MCMC fits
- No $\hat{\theta}$, fuzzy $k \Rightarrow$ no AIC

2. WOD reveal (~10 min)

- $-\log$ post. $= \sum(y_i - \mu)^2/(2\sigma^2) + \mu^2/(2\tau^2)$
- $\lambda = 1/(2\tau^2)$; ridge = posterior mode; $\tau \to \infty$ = MLE
- Lasso = Laplace prior


3. WAIC and LOO (~12 min) [BOARD → R]

- elpd: per point, log of posterior-averaged lik.; sum
- p_D, p_{waic}, p_{loo}: effective parameters
- LOO = PSIS refit shortcut; Pareto $k > 0.7$ = unreliable
- Stan saves `log_lik[i]`

	no wave	full	effective parameters
AIC	2278.0	2282.0	k = 3 vs. 6
DIC	2278.0	2281.9	pD = 3.0 vs. 5.9
WAIC	2278.6	2282.7	p_waic = 3.5 vs. 6.5
LOOIC	2278.6	2282.7	p_loo = 3.5 vs. 6.5

- **Pause:** full 2.1 behind, SE 1.8? About 1 SE: can't tell apart
- AIC: no SE. LOO: SE

4. Shrinkage priors (~12 min) [BOARD → R]



- Prior scale = λ; Normal = ridge, Laplace = lasso
- North: vague 0.23, Laplace 0.08
- p_{loo}: 9.1 → 7.8 → 7.1
- **Pause:** lasso 0 vs. Laplace 0.08 (-0.07, 0.28)? Posterior keeps uncertainty
- Horseshoe: spike + heavy tails (sketch)
- Prior scale = λ: LOO, or hierarchical later

5. Discussion (~5 min)

Q1. No-wave fir model ahead by 2.1 elpd (SE 1.8). "Wave has no effect"?

- Inconclusive; not detected $\neq$ no effect

Q2. Northness: Laplace 0.08 (-0.07 to 0.28) vs. lasso 0. Report which?

- Posterior shows the uncertainty a zero hides

If time: Bayes factors (marginal lik., prior-sensitive); RJMCMC / spike-and-slab (not Stan)