Power of a point and cyclic quadrilaterals

Kevin Hu
November 2026

L3-GE1 Level 3 Senior · Geometry

Turn circle diagrams into equations using equal tangents, the power of a point and the tests for four points being concyclic.
By the end you can use equal tangent lengths and the power of a point to turn a circle diagram into equations; prove that four points are concyclic, including with the converse of power of a point; combine power of a point with Pythagoras and similar triangles to find lengths at SMC, AMC 12 and early AIME level.
Prerequisites: Angle chasing and polygons (L2-GE1), Area, Pythagoras and similar shapes (L2-GE2).
Contests: UKMT SMC, Senior Kangaroo, Maclaurin, AMC 12, AIME. **Time:** about 90 minutes.

1 Lesson

Two chords AB and CD of a circle cross at P, with $AP = 4$, $PB = 6$ and $CP = 3$. Neither the radius nor any angle is given, but PD is determined: $PD = 8$. This is a special case of the *power of a point*, a single theorem which also covers tangents, lines through a point outside the circle, and many questions of the form "is this quadrilateral cyclic?". At SMC and AMC 12 level, many circle questions that seem to need trigonometry can be done in two or three lines with it.

1.1 Tangent lengths

Equal tangents.
The two tangents from an external point P to a circle have equal length, and the line from P to the centre bisects the angle between them.


Proof. let the tangents touch the circle at S and T, and let O be the centre. A tangent is perpendicular to the radius at the point of contact, so $\angle OSP = \angle OTP = 90^\circ$. Also $OS = OT$ and OP is shared, so triangles OSP and OTP are congruent (RHS). Hence $PS = PT$ and $\angle SPO = \angle TPO$.

The most common application is to the incircle. If the incircle of triangle ABC touches BC, CA, AB at D, E, F, the six tangent segments come in three equal pairs: $AE = AF = x$, $BF = BD = y$ and $CD = CE = z$. Adding all six gives $2(x + y + z) = a + b + c$, so $x + y + z = s$, the semiperimeter, and therefore

$$AE = AF = s - a, \qquad BF = BD = s - b, \qquad CD = CE = s - c,$$

where $a = BC$, $b = CA$ and $c = AB$. In the same way, if a circle touches all four sides of a quadrilateral $ABCD$, then $AB + CD = BC + DA$, because each side of that equation contains each of the four tangent lengths exactly once.

Kevinvariant
kevinvariant08.github.io



Example 1.
A right-angled triangle has an incircle of radius 4 and a hypotenuse of length 26. Find its perimeter.

Solution.
Calling the legs a and b gives $a^2 + b^2 = 676$, which is one equation in two unknowns. The inradius has not been used yet, and tangent lengths are how it enters.
Note that at the right angle, the incentre and the two points of contact on the legs form a square of side 4, since both radii are perpendicular to the legs. So the two tangents from the right-angled vertex have length 4. Call the tangent lengths from the other two vertices x and y. The hypotenuse is made of one tangent from each of those vertices, so $x + y = 26$. The perimeter consists of every tangent length twice, so it equals

$$2(4 + x + y) = 2(4 + 26) = 60.$$

The legs were not needed. For the record, $a + b = 34$ and $a^2 + b^2 = 676$ give $ab = 240$, so the triangle is 10, 24, 26, and indeed $\frac{1}{2}(10 + 24 - 26) = 4$.

1.2 The power of a point

Power of a point.
Let ω be a circle with centre O and radius r, and let P be any point. If a line through P meets ω at A and B, then


$$PA \cdot PB = \left|OP^2 - r^2\right|,$$

whichever line through P you choose. In particular:

- if chords AB and CD meet at a point P inside the circle, then $PA \cdot PB = PC \cdot PD$;
- if two lines through an outside point P meet the circle at A, B and at C, D, then $PA \cdot PB = PC \cdot PD$;
- if P is outside and PT is a tangent, then $PT^2 = PA \cdot PB$.

The number $OP^2 - r^2$ is called the power of P. It is negative inside the circle, zero on it and positive outside.

Kevinvariant
kevinvariant08.github.io




Proof. Drop the perpendicular from O to the line, meeting it at M. The perpendicular from the centre to a chord bisects it, so $MA = MB$. Suppose first that P is outside, as in the diagram, and label so that A is the nearer point. Then $PA = PM - MA$ and $PB = PM + MA$, so

$$PA \cdot PB = PM^2 - MA^2 = (OP^2 - OM^2) - (r^2 - OM^2) = OP^2 - r^2,$$

using Pythagoras in the right-angled triangles OMP and OMA. If P is inside, then with A again the nearer point, $PA = MA - PM$ and $PB = MA + PM$, and the same calculation gives $r^2 - OP^2$. For the tangent, Pythagoras in triangle OTP gives $PT^2 = OP^2 - r^2$ directly. The answer involves only OP and r, so every line through P gives the same product.

I include a second proof by similar triangles, because the same triangles turn up in harder problems. For chords AB and CD meeting at P, we have $\angle PAC = \angle BAC = \angle BDC = \angle PDB$ (angles in the same segment) and $\angle APC = \angle DPB$ (vertically opposite), so triangles PAC and PDB are similar and $PA/PD = PC/PB$. For a tangent, the alternate segment theorem gives $\angle PTA = \angle PBT$, so triangles PTA and PBT are similar and $PT/PA = PB/PT$.




Example 2.
Point P lies outside a circle. One line through P meets the circle at A and B, with $PA = 4$ and $AB = 5$. A second line through P meets the circle at C and D, with C between P and D and $PC = CD$. Find PC, and the length of a tangent from P to the circle.

Solution.
A common slip is to multiply PA by AB. Power of a point always measures both lengths from P: here $PB = PA + AB = 9$, so the power of P is $PA \cdot PB = 36$.
On the second line $PD = 2\,PC$, so $PC \cdot 2PC = 36$, giving $PC^2 = 18$ and $PC = 3\sqrt{2}$. A

Kevinvariant
kevinvariant08.github.io

tangent PT satisfies $PT^2 = 36$, so $PT = 6$.

Power of a point gives products of lengths. To get at the radius you usually need the centre as well, and the centre is easy to locate: it lies on the perpendicular bisector of every chord.



Example 3.
Chords AB and CD of a circle are perpendicular and meet at P, with $AP = 3$, $PB = 8$ and $CP = 4$. Find the radius of the circle.

Solution.
First, power of a point gives the missing length: $PD = \dfrac{AP \cdot PB}{CP} = \dfrac{24}{4} = 6$.
To find the radius we locate the centre. The perpendicular bisector of AB is perpendicular to AB, so it is parallel to CD, and the perpendicular bisector of CD is parallel to AB. So the position of the centre O relative to P can be read off without any angles.
Let M and N be the midpoints of AB and CD. Since $AB = 11$, $PM = \frac{11}{2} - 3 = \frac{5}{2}$, and since $CD = 10$, $PN = 5 - 4 = 1$. As $OM \perp AB$ and $ON \perp CD$, the quadrilateral $PMON$ is a rectangle, so $OM = PN = 1$. In the right-angled triangle OMB,

$$r^2 = OM^2 + MB^2 = 1^2 + \left(\tfrac{11}{2}\right)^2 = \tfrac{125}{4},$$

so $r = \frac{5\sqrt{5}}{2}$.
More generally, whenever perpendicular chords meet at P, we get $PA^2 + PB^2 + PC^2 + PD^2 = 4r^2$, and indeed $9 + 64 + 16 + 36 = 125$. Proving it is a good exercise: put P at the origin with the chords along the axes.

1.3 Cyclic quadrilaterals

Concyclic criteria.
Four points A, B, C, D lie on a circle if any one of these holds:


1. $ABCD$ is a convex quadrilateral with a pair of opposite angles adding to $180°$;
2. C and D are on the same side of line AB and $\angle ACB = \angle ADB$;

Kevinvariant
kevinvariant08.github.io

3. (converse of power of a point) lines AB and CD meet at P, the point P lies either inside both segments AB, CD or outside both, and $PA \cdot PB = PC \cdot PD$.

You met the first two in L2-GE1. The third is the converse of power of a point, and it is often overlooked. Proof: let the circle through A, B, C meet line PC again at D'. Power of a point gives $PA \cdot PB = PC \cdot PD'$, so $PD' = PD$. If P is inside segment AB, it is inside the circle, so D' is on the opposite side of P from C, just as D is. If P is outside segment AB, it is outside the circle, so D' and D are both on the same side of P as C. Either way $D' = D$.

There are two special cases you must know. If $\angle AXB = \angle AYB = 90°$, then X and Y lie on the circle with diameter AB. In particular, if BE and CF are altitudes of triangle ABC, then B, C, E, F are concyclic; the next example uses this.



Example 4.
In an acute triangle ABC the altitudes BE and CF meet at H, with E on AC and F on AB. Given $AF = 3$, $FB = 5$ and $AE = 4$, find AC and BC.

Solution.
Pythagoras in triangle AEB gives $BE^2 = 8^2 - 4^2 = 48$, but in triangle AFC we know only AF, and nothing so far determines C. We need a relation involving AC, and finding it is the hard step.
Note that $\angle BEC = \angle BFC = 90°$, so E and F both lie on the circle with diameter BC (dashed). Since angle A is acute, A lies outside that circle, and it has two lines through it: line AB meets the circle at F and B, and line AC meets it at E and C. Power of the point A gives

$$AF \cdot AB = AE \cdot AC, \qquad 3 \cdot 8 = 4 \cdot AC,$$

so $AC = 6$ and $EC = 2$.
Pythagoras in triangle BEC now gives $BC^2 = BE^2 + EC^2 = 48 + 4 = 52$, so $BC = 2\sqrt{13}$. As a check, $CF^2 = 6^2 - 3^2 = 27$ and $CF^2 + FB^2 = 27 + 25 = 52$ as well.
There is a second route. In right-angled triangle AFC, $\cos A = AF/AC$, and in triangle AEB, $\cos A = AE/AB$. Equating the two gives $AF \cdot AB = AE \cdot AC$ again.

Common traps.

- Measure from P. For a line through P meeting the circle at A then B, the product is

Kevinvariant
kevinvariant08.github.io

$PA \cdot PB$, never $PA \cdot AB$.

- Check the order of the points before writing $PB = PA + AB$. A diagram that is not to scale can put the nearer point on the wrong side.
- Do not assume points are concyclic because the picture looks round. Prove it with one of the criteria: right angles, equal angles on the same side, or the converse of power of a point.
- In a quadrilateral with an incircle, the equal sums are of *opposite* sides: $AB + CD = BC + DA$.
- A chord through an interior point P is at least as long as the chord perpendicular to OP and at most a diameter. If a question asks for all possible lengths, check both ends of that range.

1.4 How to spot this in a contest

For lengths around a circle, the strategies in short are:

1. If two lines pass through a common point and both meet the same circle, use power of a point, especially when the question gives lengths along those lines and no angles. A tangent from an external point is the case where the two intersection points coincide.
2. If no circle is drawn, look for one: two right angles on the same segment, or two equal angles subtending the same segment on the same side, mean four points are concyclic whether or not the setter says so.
3. If the lengths must be integers (common in the AMC), factorise the power and check which factor pairs give a chord of possible length.
4. If a circle touches several sides of a polygon, label the tangent lengths before doing anything else; the number of unknowns often drops from six to three or fewer.

2 Practice problems

The problems are roughly in order of difficulty, from ⋆ to ⋆⋆⋆⋆. Hints are in Section 4 and full solutions in Section 5.

Problem 1 ⋆ Chords AB and CD of a circle meet at P. Given $AP = 6$, $PB = 4$ and $CD = 11$, find the length of the longer of the two segments CP and PD.

Problem 2 ⋆⋆ Point P is at distance 13 from the centre of a circle of radius 5. A line through P meets the circle at A and B, with A between P and B, and $PA = 2\,AB$. Find AB.

Problem 3 ⋆⋆ In a convex quadrilateral $ABCD$ the diagonals meet at P, with $AP = 4$, $PC = 9$, $BP = 6$ and $PD = 6$. Given that $AB = 5$, find CD.

Problem 4 ⋆⋆⋆ A circle has centre O and radius 7, and P is a point with $OP = 3$. A chord AB passes through P, and both PA and PB are whole numbers. Find the sum of all possible lengths of AB.

Kevinvariant
kevinvariant08.github.io

Problem 5 ⋆⋆⋆ Triangle ABC has $AB = 13$, $BC = 14$ and $CA = 15$. The circle through A that touches the line BC at B meets CA again at D. Find BD.

Problem 6 ⋆⋆⋆⋆ Triangle ABC has $AB = 13$, $BC = 14$ and $CA = 15$. Its incircle touches BC at D, and the segment AD meets the incircle again at X. Find $\dfrac{XD}{AX}$.

3 Contest ladder

Real past problems on the ideas of this unit, easiest first. Each source is a link to the official paper or the AoPS wiki; the note under it says where to start, not how to finish.

1. **UKMT SMC 2020 Question 15** ⋆
   Two tangents from one point: split the region along the line of symmetry
2. **UKMT SMC 2015 Question 12** ⋆⋆
   Equal tangents make isosceles triangles; chase angles
3. **UKMT SMC 2016 Question 24** ⋆⋆
   Label the tangent lengths from each external point, then Pythagoras
4. **AMC 12A 2013 Problem 19** ⋆⋆⋆
   The power of C with respect to the circle, then a size check
5. **UKMT SMC 2017 Question 19** ⋆⋆⋆
   The corner of the rectangle has a known power; or drop a perpendicular from the centre
6. **AMC 12B 2019 Problem 20** ⋆⋆⋆
   Equal tangents put the meeting point on the perpendicular bisector of the chord
7. **AIME 1999 Problem 12** ⋆⋆⋆
   Tangent lengths give every side in terms of one unknown; compare two area formulas
8. **UKMT Maclaurin 2022 Question M6** ⋆⋆⋆
   Angles in the same segment give a similar pair sharing the angle at C
9. **AIME I 2024 Problem 10** ⋆⋆⋆⋆
   Tangent-chord angles, then the power of the point where the tangents meet
10. **UKMT Maclaurin 2021 Question M3** ⋆⋆⋆⋆
    Write down the power of A with respect to each circle
11. **AIME I 2019 Problem 13** ⋆⋆⋆⋆
    Find a cyclic angle equality, then meet line CF with line AB and use power of a point
12. **AIME I 2016 Problem 15** ⋆⋆⋆⋆
    The line XY bisects the common tangent; compute powers along it

Kevinvariant
kevinvariant08.github.io


4 Hints

1. Write $CP = x$ and $PD = 11 - x$. You already know the product $CP \cdot PD$.
2. Find the power of P from the distance to the centre and the radius, then write everything in terms of AB.
3. Compare $PA \cdot PC$ with $PB \cdot PD$. What does that tell you about the four vertices?
4. Power of a point gives the product $PA \cdot PB$. Then ask: what are the shortest and longest chords through P?
5. Look at the point C: one of the lines through it is a tangent. The similar triangles behind the tangent case give more than the power alone.
6. The tangent length from A to the incircle is easy to find, and then power of a point needs only one more length.

5 Solutions

Problem 1

Chords AB and CD of a circle meet at P. Given $AP = 6$, $PB = 4$ and $CD = 11$, find the length of the longer of the two segments CP and PD.




By the intersecting chords theorem, $CP \cdot PD = AP \cdot PB = 24$. With $CP = x$ and $PD = 11 - x$ this reads $x(11 - x) = 24$, that is $x^2 - 11x + 24 = 0$, which factorises as $(x - 3)(x - 8) = 0$. So the two segments have lengths 3 and 8, and the longer one is 8.

Answer: 8

Problem 2

Point P is at distance 13 from the centre of a circle of radius 5. A line through P meets the circle at A and B, with A between P and B, and $PA = 2\,AB$. Find AB.

Kevinvariant
kevinvariant08.github.io




The power of P is $OP^2 - r^2 = 169 - 25 = 144$; equivalently, a tangent from P has length 12. Write $AB = x$. Then $PA = 2x$ and $PB = PA + AB = 3x$, so $PA \cdot PB = 6x^2 = 144$. This gives $x^2 = 24$, so $AB = 2\sqrt{6}$.

We should check that such a line exists. The nearer intersection point is at distance between $OP - r = 8$ (the line through the centre) and 12 (the tangent), and $PA = 4\sqrt{6} \approx 9.8$ lies in that range.

Answer: $2\sqrt{6}$

Problem 3

In a convex quadrilateral $ABCD$ the diagonals meet at P, with $AP = 4$, $PC = 9$, $BP = 6$ and $PD = 6$. Given that $AB = 5$, find CD.



We have $PA \cdot PC = 36 = PB \cdot PD$, and P lies inside both diagonals, so by the converse of the intersecting chords theorem $ABCD$ is cyclic. Then $\angle BAC = \angle BDC$ (angles in the same segment), that is $\angle PAB = \angle PDC$, and $\angle APB = \angle DPC$ (vertically opposite). So triangles APB and DPC are similar, with A, P, B corresponding to D, P, C. The ratio of similarity is $DP/AP = 6/4 = \frac{3}{2}$, so $CD = \frac{3}{2} \cdot AB = \frac{15}{2}$.


As a check, CP/BP should also equal $\frac{3}{2}$, and indeed $9/6 = \frac{3}{2}$. In fact that check is a proof on its own: $AP/DP = BP/CP$ with the vertically opposite angles gives the similarity directly (SAS). So the converse of power of a point is equivalent to this SAS similarity.

Kevinvariant
kevinvariant08.github.io

Answer: $\frac{15}{2}$

Problem 4

A circle has centre O and radius 7, and P is a point with $OP = 3$. A chord AB passes through P, and both PA and PB are whole numbers. Find the sum of all possible lengths of AB.



The power of P gives $PA \cdot PB = r^2 - OP^2 = 49 - 9 = 40$.

Next we find the possible lengths of a chord through P. A chord at distance d from O has length $2\sqrt{49 - d^2}$, and a chord through P has $0 \le d \le OP = 3$, with every such d occurring as the chord turns about P. So the possible lengths are exactly those from $2\sqrt{40} \approx 12.6$ (the chord perpendicular to OP) up to 14 (the diameter).


The factor pairs of 40 are 1×40, 2×20, 4×10 and 5×8, with sums 41, 22, 14 and 13. Only 14 and 13 lie in the allowed range. The sum 14 is the diameter through P, with $PA = 7 - 3 = 4$ and $PB = 7 + 3 = 10$; the sum 13 is a chord with $PA = 5$ and $PB = 8$. The answer is $13 + 14 = 27$. Forgetting the diameter is the trap here.

Answer: 27

Problem 5

Triangle ABC has $AB = 13$, $BC = 14$ and $CA = 15$. The circle through A that touches the line BC at B meets CA again at D. Find BD.

Kevinvariant
kevinvariant08.github.io



From C, the line CB is tangent at B and the line CA meets the circle at D and A, so power of a point gives $CB^2 = CD \cdot CA$, that is $CD = \frac{196}{15}$. This is less than 15, so D lies between A and C.

For BD, use the similar triangles behind the tangent case. The line BD separates A from C, so by the alternate segment theorem the angle between the tangent BC and the chord BD equals the angle that BD subtends at A: $\angle CBD = \angle BAD = \angle BAC$. Triangles CBD and CAB also share the angle at C, so they are similar, with C, B, D corresponding to C, A, B. Hence $BD/AB = CB/CA$, and


$$BD = 13 \cdot \frac{14}{15} = \frac{182}{15}.$$

The same similarity gives $CD/CB = CB/CA$, which is the power of a point statement again.

Answer: $\frac{182}{15}$

Problem 6

Triangle ABC has $AB = 13$, $BC = 14$ and $CA = 15$. Its incircle touches BC at D, and the segment AD meets the incircle again at X. Find $\dfrac{XD}{AX}$.



We start with tangent lengths. The semiperimeter is $s = 21$, so the tangents from A to the incircle (to the points Y and Z in the diagram) have length $s - a = 21 - 14 = 7$, and $BD = s - b =$

Kevinvariant
kevinvariant08.github.io

$21 - 15 = 6$. Power of the point A with respect to the incircle gives

$$AX \cdot AD = 7^2 = 49.$$

It remains to find AD, which is fairly routine with coordinates. Put $B = (0,0)$ and $C = (14,0)$, and let $A = (x,y)$. From $x^2 + y^2 = 169$ and $(x-14)^2 + y^2 = 225$, subtracting gives $28x - 196 = -56$, so $x = 5$ and then $y = 12$. Since $D = (6,0)$, we get $AD^2 = 1^2 + 12^2 = 145$.

Therefore $AX = \dfrac{49}{\sqrt{145}}$ and $XD = AD - AX = \dfrac{145 - 49}{\sqrt{145}} = \dfrac{96}{\sqrt{145}}$, so

$$\frac{XD}{AX} = \frac{96}{49}.$$

In general the ratio is $\dfrac{AD^2 - t^2}{t^2}$, where $t = 7$ is the tangent length, so $\sqrt{145}$ need not appear at all.

Answer: $\frac{96}{49}$