HU Collection · Kevinvariant Pathways

Writing olympiad solutions

Level 4 Olympiad · Strategy

How BMO Round 1 is marked and how to write solutions that earn full credit: structure, complete proofs, case analysis, both directions of a "find all" answer, and the classic ways to lose marks.

By the end you can: You will be able to explain how BMO1 scripts are marked and what earns credit; You will be able to write "find all" solutions that prove both directions with complete case analysis; You will be able to find and repair the gaps in a flawed proof.

Lesson

Consider the following question and two answers to it.

Find all primes p such that $p^2 + 2$ is also prime.

Script A. "If $p = 2$ we get 6, not prime. If $p = 3$ we get 11, prime. If $p = 5$ we get 27, and if $p = 7$ we get $51 = 3 \times 17$. So the answer is $p = 3$."


Script B. "The answer is $p = 3$. Indeed $3^2 + 2 = 11$ is prime. Conversely, suppose $p \neq 3$. Then $3 \nmid p$, so $p \equiv \pm 1 \pmod 3$ and $p^2 \equiv 1 \pmod 3$. Hence $3 \mid p^2 + 2$, and since $p^2 + 2 > 3$ it is not prime. So $p = 3$ is the only such prime."

Both have the right answer. On a BMO paper, Script A scores 0 or 1 out of 10 and Script B scores 10. Script A has found the answer; Script B has proved it. This unit is about writing solutions of the second kind.

How BMO Round 1 is marked

From UKMT's BMO1 pages and the paper's instructions: six problems, three and a half hours, 10 marks each, 60 in total. You must give full written solutions, not just answers, with complete proofs of any assertions; marks depend on the clarity of your presentation, and one complete solution gains more credit than several unfinished attempts.

UKMT's markers' reports explain what happens in practice. Marking is holistic. A script is first judged on whether it is essentially a complete solution. If it is, it is marked down from 10 ("10 minus") for slips, gaps, omitted cases or technical faults. If it is not, it is marked up from 0 ("0 plus"), and partial progress such as particular cases or a significant idea earns at most about 4 marks. On many questions a 5 is not available. The reports also say that correct arguments are rewarded more generously than correct calculations, and that you should aim to finish whole questions rather than collect disconnected parts.



Compare this with typical scores. On the November 2025 paper the mean was 12.6 out of 60, and the Distinction threshold was 18. Two complete solutions put you in the Distinction band; six half-solutions, each marked "0 plus", might not. Hence the priority is to finish questions.

Kevinvariant
kevinvariant08.github.io

The shape of a good solution

1. **State the answer first** in any "find" or "determine" question, in one line.
2. **Say what you will prove**: "We show that n works if and only if $n \equiv 0$ or $3 \pmod 4$." Introduce notation before you use it.
3. **Break the proof into claims**, prove each one and say how they combine.
4. **End with a conclusion** that answers the question as asked.

Rough work stays on rough paper. In geometry, draw a large labelled diagram and make the order of deductions clear: a diagram covered in marked angles with no explanation earns nothing.

What counts as a complete proof

Every step must follow from earlier steps or from a standard result that you name. "Clearly" is fine only if a reader could fill the gap in five seconds. Reread your solution as a hostile marker, asking "why?" at every line; if the answer is not on the page and is not obvious, put it there. Three failures come up year after year.

Dividing by something that might be zero. Cancelling $x - y$ assumes $x \neq y$. Split into cases instead: either $x = y$, or you may divide.

"Eventually" claims. "Exponentials beat polynomials, so $2^n > n^3$ for large n" is not a proof. Say from which n it holds and prove it, usually by induction.

Careless coprimality. If $ab = 2^k m$ with m odd and $\gcd(a, b) = 1$, then 2^k divides one of a, b. It does not follow that $\{a, b\} = \{2^k, m\}$, because the prime factors of m can be split between a and b.

Example 1. Find all integers n such that $n^2 + 2$ divides $n^4 + 7$.

Solution.
Small values first: $n = 0, \pm 1, \pm 2$ fail, and $n = \pm 3$ gives $11 \mid 88$, which works. To see whether that is all, remove the leading term by division: $n^4 + 7 = (n^2 + 2)(n^2 - 2) + 11$. The rest follows from this identity.
A good write-up: "The answer is $n = \pm 3$. Since $n^4 + 7 = (n^2 + 2)(n^2 - 2) + 11$, the number $n^2 + 2$ divides $n^4 + 7$ if and only if it divides 11. As $n^2 + 2$ is a positive integer that is at least 2, this happens if and only if $n^2 + 2 = 11$, that is, $n = \pm 3$."
Because each step is an "if and only if", both directions are proved at once; one-way implications would only show that ± 3 are the only candidates. And the question says integers, so a script giving only $n = 3$ has lost a solution.

Case analysis

Cases must cover every possibility, and you must say why they do. Announce the split before you start ("either $p = 2$ or p is odd") and treat each case under its own heading. Small cases matter: if your general argument needs $n \geq 4$, then $n = 1, 2, 3$ must be checked with the working shown. One recent report notes that candidates merely stating they had checked the cases was not enough. You may assume an order such as $a \leq b \leq c$ only when the problem is symmetric, and you must then restore every ordering in the final answer.

Example 2. Find all triples of positive integers (a, b, c) with $abc = 2(a + b + c)$.

Solution.

Kevinvariant
kevinvariant08.github.io

The equation is symmetric, so we may assume $a \le b \le c$ and permute at the end. The first step is a bound. The right side is at most $6c$, so $abc \le 6c$ and $ab \le 6$. That leaves finitely many cases. Rewrite the equation as $c(ab-2) = 2(a+b)$.
The pairs with $a \le b$ and $ab \le 6$ are $(1,1)$, $(1,2)$, $(1,3)$, $(1,4)$, $(1,5)$, $(1,6)$, $(2,2)$ and $(2,3)$. For $(1,1)$ the equation reads $-c = 4$, and for $(1,2)$ it reads $0 = 6$; both are impossible. For $(1,3)$ we get $c = 8$, for $(1,4)$ we get $2c = 10$ so $c = 5$, and for $(2,2)$ we get $2c = 8$ so $c = 4$. For $(1,5)$ we get $3c = 12$, so $c = 4$, which breaks $b \le c$; it is $(1,4,5)$ again in another order. For $(1,6)$ we get $4c = 14$ and for $(2,3)$ we get $4c = 10$, neither of which has an integer solution.
So the solutions are the permutations of $(1,3,8)$, $(1,4,5)$ and $(2,2,4)$, and each works: $24 = 2 \times 12$, $20 = 2 \times 10$ and $16 = 2 \times 8$. In the write-up, justify the ordering by symmetry, show the bound and list every case with its outcome, so that a marker can tick each one without doing any arithmetic.

Writing up "find all"

Two directions. To answer "find all x with property P" with a set S, you must prove both of the following.

1. Every element of S has property P. This is the checking direction, done by substitution or by construction.
2. Every x with property P lies in S. This is the necessity direction, usually the hard part.

Missing the first direction costs marks. Missing the second means the problem is not solved. The same applies to "for which n is it possible?": you need a construction for the good n and an impossibility proof for the rest.

The checking direction is not a formality. Squaring, multiplying or combining equations can let in false solutions; the 2025 report notes candidates whose derived equations admitted sign combinations that failed the original system.

Example 3. For which positive integers n can $\{1, 2, \dots, n\}$ be split into two sets with equal sums?

Solution.
Necessity is a parity argument. The total is $\frac{n(n+1)}{2}$, and it must be even, so $4 \mid n(n+1)$. Since n and $n+1$ are coprime and only one of them is even, this means $4 \mid n$ or $4 \mid n+1$, that is, $n \equiv 0$ or $3 \pmod 4$.
Sufficiency needs a construction. The block $\{k, k+1, k+2, k+3\}$ splits evenly as $\{k, k+3\}$ and $\{k+1, k+2\}$. If $n \equiv 0 \pmod 4$, cut $1, \dots, n$ into such blocks and split each. If $n \equiv 3 \pmod 4$, put $\{1,2\}$ against $\{3\}$ and cut $4, \dots, n$ into blocks.
The write-up should state the answer, then prove necessity and sufficiency under separate headings. A script with only one of the two is marked "0 plus".

An annotated model solution

Example 4. Find all pairs (m, n) of positive integers such that $m+n$ divides both m^2+n and n^2+m.

Solution.
In rough: $(1,1)$ and $(3,3)$ work, while $(2,2)$ and $(1,2)$ fail. A guess: $m = n$ with m odd. Note that modulo $m+n$ we can replace n by $-m$. I have annotated the write-up below with remarks

Kevinvariant
kevinvariant08.github.io

on how a marker reads it.
Answer. The pairs (m, m) with m odd.
Marker's eye: the answer comes first, so the marker knows what is being proved.
These pairs work. If $m = n$ is odd, then $m + n = 2m$ and $m^2 + n = m(m + 1)$. As $m + 1$ is even, $m(m + 1) = 2m \cdot \frac{m+1}{2}$ is a multiple of $2m$. The second condition is identical, since $m = n$.
Marker's eye: the checking direction is short, so do it first and secure it.
No other pairs work. Let $s = m + n$ and suppose s divides $m^2 + n$ and $n^2 + m$. Since $n \equiv -m \pmod{s}$,

$$m^2 + n \equiv m^2 - m = m(m - 1) \quad \text{and} \quad n^2 + m \equiv m^2 + m = m(m + 1) \pmod{s}.$$

So s divides both $m(m - 1)$ and $m(m + 1)$, and therefore their difference, $2m$.
Now $2m > 0$, and $2m < 2s$ because $n \geq 1$. The only multiple of s that is greater than 0 and less than $2s$ is s itself, so $2m = m + n$, giving $n = m$. Finally, with $n = m$ the condition is $2m \mid m(m + 1)$, and dividing by m this says $2 \mid m + 1$, so m is odd.
Marker's eye: this paragraph is the hard step. Writing "$s \mid 2m$, so $s = 2m$" with no reason would be a gap, because a divisor of $2m$ can be smaller than $2m$. The bound $0 < 2m < 2s$ is what justifies the step, and it is written out.
Conclusion. The pairs are exactly (m, m) with m odd.
Marker's eye: the conclusion repeats the answer in the form the question asked for.

Common ways to lose marks (all drawn from recent BMO1 markers' reports):

- giving the answer with no proof that nothing else works, or never checking that the claimed answers work;
- dividing by a quantity that might be zero;
- missing cases, often small ones such as $p = 2$ or $n = 1$;
- assuming a special case, such as treating real variables as integers;
- completing a table by pattern and calling it a proof;
- claims like "exponentials grow faster" with no starting point;
- geometry with angle facts that are stated but not justified by a named theorem;
- misreading the question, and so proving something else.

How to spot this in a contest

On BMO the write-up is what is marked. Before you write, know the answer and the outline of the proof, and leave real time for writing: a solution scrawled out in the last five minutes is often marked "0 plus".

When you finish, reread as a marker and check the following.

1. The answer is stated.
2. Both directions are proved.
3. Every division is safe.
4. The cases are exhaustive, with small ones shown.
5. Every "clearly" is justified.

Kevinvariant
kevinvariant08.github.io

Fixing a gap now costs two minutes; leaving it can cost five marks.

Practice

Try each problem before turning to the hints and solutions at the back.

1. ⋆ Find all positive integers n for which n^4+4 is prime.

   A student writes: *"We have $n^4+4=(n^2+2)^2-(2n)^2=(n^2+2n+2)(n^2-2n+2)$. Both factors are greater than 1, so n^4+4 is never prime, and there are no such n."*

   Find the gap, correct the answer and write a complete solution.

2. ⋆⋆ Find all functions $f:\mathbb{Z}\to\mathbb{Z}$ such that

$$f(x+y)=f(x)+f(y)+2xy$$

   for all integers x and y. Write a complete solution to BMO standard.

3. ⋆⋆ Find all real solutions (x,y) of the system

$$x^2+y=3,\qquad y^2+x=3.$$

   A student writes: *"Subtracting, $x^2-y^2-(x-y)=0$, so $(x-y)(x+y-1)=0$. Dividing by $x-y$ gives $x+y=1$. Then $y=1-x$, so $x^2+1-x=3$, that is, $x^2-x-2=0$, giving $x=2$ or $x=-1$. The solutions are $(2,-1)$ and $(-1,2)$."*

   Find the gap and give a complete solution.

4. ⋆⋆⋆ Prove that for every positive integer n,

$$1+\frac{1}{2^2}+\frac{1}{3^2}+\cdots+\frac{1}{n^2}<2.$$

   A student writes: *"Let S_n be the sum. Then $S_1=1<2$. Suppose $S_k<2$. Then $S_{k+1}=S_k+\frac{1}{(k+1)^2}<2+\frac{1}{(k+1)^2}$, and since $\frac{1}{(k+1)^2}$ is very small, $S_{k+1}<2$. By induction, $S_n<2$ for all n."*

   Explain why this is not a proof, and write a correct one.

5. ⋆⋆⋆ Determine all positive integers n for which the set $\{1,2,\ldots,n\}$ can be split into three sets with equal sums. Write a complete solution to BMO standard.

6. ⋆⋆⋆⋆ Find all pairs (a,b) of positive integers such that a^2+5b and b^2+5a are both perfect squares. Write a complete solution to BMO standard.

Contest ladder

Real past problems that use this idea, easiest first. Open them on the official paper or the AoPS wiki, or online at kevinvariant08.github.io.

1. UKMT BMO1 2022 Problem 1 ⋆⋆
2. UKMT BMO1 2023 Problem 4 ⋆⋆
3. USAJMO 2011 Problem 1 ⋆⋆

Kevinvariant
kevinvariant08.github.io

4. UKMT BMO1 2025 Problem 2 ⋆⋆⋆
5. UKMT BMO1 2025 Problem 1 ⋆⋆⋆
6. USAJMO 2023 Problem 1 ⋆⋆⋆
7. UKMT BMO1 2024 Problem 2 ⋆⋆⋆
8. USAJMO 2013 Problem 1 ⋆⋆⋆
9. UKMT BMO1 2023 Problem 2 ⋆⋆⋆⋆
10. USAJMO 2015 Problem 4 ⋆⋆⋆⋆

Kevinvariant
kevinvariant08.github.io

Hints

1. The factorisation is right. Test the claim about the factors on the smallest possible n.
2. Compare with x^2. What equation does $g(x) = f(x) - x^2$ satisfy? Then use induction carefully, not forgetting negative integers.
3. What did the student assume when they divided? Treat that possibility as a separate case.
4. The statement $S_k < 2$ is too weak to carry the induction. Try to prove something stronger, of the form $S_n \leq 2 -$ (something positive).
5. There are two necessary conditions: one from divisibility of the total, and one from the set that contains n. For sufficiency, find a block of six consecutive integers that splits into three pairs with equal sums.
6. Assume $a \geq b$, which is allowed by symmetry. Then $a^2 + 5b$ is trapped between a^2 and $(a+3)^2$. Each of the resulting cases leads to a second equation that can be turned into a difference of two squares.

Solutions

Problem 1

Answer: $n = 1$

The claim that both factors exceed 1 is asserted without proof, and it is false for $n = 1$, where $n^2 - 2n + 2 = 1$. In that case $n^4 + 4 = 5$, which is prime, so the student's answer is wrong.

Here is a complete solution. The answer is $n = 1$. For $n = 1$ we get 5, which is prime. Now let $n \geq 2$. We have $n^4 + 4 = (n^2 + 2n + 2)(n^2 - 2n + 2)$, and $n^2 - 2n + 2 = (n-1)^2 + 1 \geq 2$ since $n - 1 \geq 1$. The other factor is larger, since it exceeds the first by $4n$. So $n^4 + 4$ is a product of two integers that are each at least 2, and it is not prime. Hence $n = 1$ is the only solution.

Problem 2

Answer. The functions $f(x) = x^2 + cx$, where c is any integer.

These functions work. If $f(x) = x^2 + cx$ then f maps integers to integers, and

$$f(x) + f(y) + 2xy = x^2 + cx + y^2 + cy + 2xy = (x+y)^2 + c(x+y) = f(x+y).$$

No other functions work. Suppose f satisfies the equation and let $g(x) = f(x) - x^2$, which is again a function from $\mathbb{Z}$ to $\mathbb{Z}$. Substituting $f(x) = g(x) + x^2$ and cancelling $(x+y)^2 = x^2 + y^2 + 2xy$ gives $g(x+y) = g(x) + g(y)$ for all integers x, y. Putting $x = y = 0$ gives $g(0) = 2g(0)$, so $g(0) = 0$. Let $c = g(1)$, an integer.

We prove $g(n) = cn$ for all $n \geq 0$ by induction. It holds for $n = 0$. If $g(n) = cn$ then $g(n+1) = g(n) + g(1) = cn + c = c(n+1)$. For negative integers, put $x = n$ and $y = -n$ with $n > 0$: then $0 = g(0) = g(n) + g(-n)$, so $g(-n) = -cn = c \cdot (-n)$. Hence $g(x) = cx$ for every integer x, and $f(x) = x^2 + cx$.

Conclusion. The solutions are exactly $f(x) = x^2 + cx$ with $c \in \mathbb{Z}$.

Problem 3

Answer: $(2, -1)$, $(-1, 2)$, $\left(\frac{-1+\sqrt{13}}{2}, \frac{-1+\sqrt{13}}{2}\right)$, $\left(\frac{-1-\sqrt{13}}{2}, \frac{-1-\sqrt{13}}{2}\right)$

Kevinvariant
kevinvariant08.github.io

Dividing by $x-y$ assumes $x \neq y$, and the case $x = y$ was never considered. That case contains two more solutions.

Here is a complete solution. Subtracting the equations shows that any solution satisfies $(x-y)(x+y-1) = 0$. Conversely, the system is equivalent to the pair of equations consisting of the first equation and $(x-y)(x+y-1) = 0$, since $y^2 + x = (x^2 + y) - (x-y)(x+y-1)$. So we find all solutions of $x^2 + y = 3$ with $x = y$ or with $x + y = 1$.

Case 1: $x = y$**.** Then $x^2 + x - 3 = 0$, so $x = \frac{-1 \pm \sqrt{13}}{2}$, giving the solutions $\left(\frac{-1+\sqrt{13}}{2}, \frac{-1+\sqrt{13}}{2}\right)$ and $\left(\frac{-1-\sqrt{13}}{2}, \frac{-1-\sqrt{13}}{2}\right)$.

Case 2: $x + y = 1$**.** Then $y = 1 - x$ and $x^2 - x - 2 = 0$, so $x = 2$ or $x = -1$, giving $(2, -1)$ and $(-1, 2)$.

The cases cover every possibility because a product is zero only if a factor is. As a check, $(2, -1)$ gives $4 - 1 = 3$ and $1 + 2 = 3$, and when $x = y$ both equations read $x^2 + x = 3$. So there are exactly four solutions.

Problem 4

From $S_k < 2$ the student correctly deduces $S_{k+1} < 2 + \frac{1}{(k+1)^2}$, but that does not give $S_{k+1} < 2$. However small the added term is, it is positive, and nothing in the hypothesis leaves room for it. The phrase "very small" hides the fact that the induction does not close. Instead we prove a stronger statement (strengthening the induction hypothesis).

Claim. For every positive integer n, $S_n \leq 2 - \frac{1}{n}$.

Proof. For $n = 1$ both sides equal 1. Suppose $S_k \leq 2 - \frac{1}{k}$ for some $k \geq 1$. Then

$$S_{k+1} \leq 2 - \frac{1}{k} + \frac{1}{(k+1)^2}.$$

We want this to be at most $2 - \frac{1}{k+1}$, which is equivalent to

$$\frac{1}{(k+1)^2} \leq \frac{1}{k} - \frac{1}{k+1} = \frac{1}{k(k+1)}.$$

This holds because $k(k+1) < (k+1)^2$. So $S_{k+1} \leq 2 - \frac{1}{k+1}$, completing the induction.

Hence $S_n \leq 2 - \frac{1}{n} < 2$ for all n. (Equivalently, one can bound $\frac{1}{j^2} < \frac{1}{j-1} - \frac{1}{j}$ for $j \geq 2$ and telescope; the strengthened hypothesis is exactly the telescoped sum.)

Problem 5

Answer. Exactly the integers $n \geq 5$ with $n \equiv 0$ or $2 \pmod 3$.

Necessity. The total is $\frac{n(n+1)}{2}$, so each set must have sum $\frac{n(n+1)}{6}$, and this must be an integer. Since $n(n+1)$ is always even, that means $3 \mid n(n+1)$. Since 3 is prime, $3 \mid n$ or $3 \mid n+1$, that is, $n \equiv 0$ or $2 \pmod 3$. Also, the set containing n has sum at least n, so $\frac{n(n+1)}{6} \geq n$, which gives $n \geq 5$.

Sufficiency. We give splittings for $n = 5, 6, 8, 9$:

$n = 5$: $\{5\}$, $\{1, 4\}$, $\{2, 3\}$, each with sum 5. $n = 6$: $\{1, 6\}$, $\{2, 5\}$, $\{3, 4\}$, each with sum 7. $n = 8$: $\{4, 8\}$, $\{5, 7\}$, $\{1, 2, 3, 6\}$, each with sum 12. $n = 9$: $\{6, 9\}$, $\{7, 8\}$, $\{1, 2, 3, 4, 5\}$, each with sum 15.

Now suppose $\{1, \ldots, n\}$ has a splitting. The six numbers $n+1, \ldots, n+6$ form the pairs $\{n+1, n+6\}$, $\{n+2, n+5\}$ and $\{n+3, n+4\}$, each with sum $2n+7$. Adding one pair to each set gives a splitting of $\{1, \ldots, n+6\}$.

Kevinvariant
kevinvariant08.github.io

The integers $n \geq 5$ with $n \equiv 0$ or $2 \pmod 3$ are those with $n \geq 5$ and $n \equiv 0, 2, 3$ or $5 \pmod 6$. In these four residue classes the smallest such n are 6, 8, 9 and 5 respectively, and every other such n is obtained from one of these by adding 6 repeatedly. So every such n has a splitting.

Conclusion. The set can be split into three sets with equal sums exactly when $n \geq 5$ and $n \equiv 0$ or $2 \pmod 3$.

Problem 6

Answer. $(4, 4)$, $(11, 27)$, $(27, 11)$, $(56, 69)$, $(69, 56)$, $(77, 192)$ and $(192, 77)$.

These pairs work. For $(4, 4)$: $16 + 20 = 36 = 6^2$. For $(27, 11)$: $27^2 + 55 = 784 = 28^2$ and $11^2 + 135 = 256 = 16^2$. For $(69, 56)$: $69^2 + 280 = 5041 = 71^2$ and $56^2 + 345 = 3481 = 59^2$. For $(192, 77)$: $192^2 + 385 = 37249 = 193^2$ and $77^2 + 960 = 6889 = 83^2$. The conditions are symmetric in a and b, so the swapped pairs work too.

No other pairs work. The conditions are symmetric, so we may assume $a \geq b$ and restore the swapped pairs at the end. Then

$$a^2 < a^2 + 5b \leq a^2 + 5a < a^2 + 6a + 9 = (a + 3)^2,$$

so $a^2 + 5b$ equals $(a + 1)^2$ or $(a + 2)^2$. Write $b^2 + 5a = k^2$ with k a positive integer.

Case 1: $a^2+5b = (a+1)^2$. Then $5b = 2a+1$, so b is odd and $a = \frac{5b-1}{2}$. Substituting, $b^2+\frac{5(5b-1)}{2} = k^2$. Multiplying by 16 and completing the square,

$$(4b + 25)^2 - (4k)^2 = 665 = 5 \cdot 7 \cdot 19.$$

Let $u = 4b + 25 - 4k$ and $v = 4b + 25 + 4k$. Then $uv = 665$ and $v > 0$, so $u > 0$, and $u < v$. Also $u + v = 8b + 50$. The factorisations of 665 give $(u, v) = (1, 665)$, $(5, 133)$, $(7, 95)$ or $(19, 35)$, with $u + v = 666$, 138, 102 or 54. Only the first two make $b = \frac{u+v-50}{8}$ a positive integer, giving $b = 77$ and $b = 11$, both odd. Then $a = 192$ and $a = 27$ respectively.

Case 2: $a^2 + 5b = (a + 2)^2$. Then $5b = 4a + 4$, so $5 \mid a + 1$. Write $a = 5t - 1$, so $b = 4t$, with t a positive integer. Substituting, $16t^2 + 25t - 5 = k^2$. Multiplying by 64 and completing the square,

$$(32t + 25)^2 - (8k)^2 = 945 = 3^3 \cdot 5 \cdot 7.$$

Let $u = 32t + 25 - 8k$ and $v = 32t + 25 + 8k$. As before $0 < u < v$, $uv = 945$ and $u + v = 64t + 50$. The factorisations are $(1, 945)$, $(3, 315)$, $(5, 189)$, $(7, 135)$, $(9, 105)$, $(15, 63)$, $(21, 45)$ and $(27, 35)$, with $u + v - 50$ equal to 896, 268, 144, 92, 64, 28, 16 and 12. This is a positive multiple of 64 only for 896 and 64, giving $t = 14$ and $t = 1$, so $(a, b) = (69, 56)$ and $(4, 4)$.

In both cases the pairs found satisfy $a \geq b$, as assumed.

Conclusion. With $a \geq b$ the solutions are $(4, 4)$, $(27, 11)$, $(69, 56)$ and $(192, 77)$. Restoring the order gives the seven pairs listed in the answer.

Kevinvariant
kevinvariant08.github.io