# Area, Pythagoras and similar shapes

Kevin Hu
November 2026

L2-GE2 Level 2 Intermediate · Geometry

How to find lengths and areas by subtraction, Pythagoras and its triples, and the ratios of similar triangles.
**By the end you can** find awkward areas by subtraction and by comparing triangles that share a height; recognise Pythagorean triples and set up Pythagoras equations, including in folding problems; spot similar triangles and use their length and area ratios.
**Prerequisites:** Angle facts and polygons (L1-GE1), Angle chasing and polygons (L2-GE1), Factorising, substitution and systems (L2-AL1). **Contests:** IMC, Grey Kangaroo, Pink Kangaroo, Cayley, Hamilton, Maclaurin, AMC 10. **Time:** about 160 minutes.

## 1 Lesson

How far is corner $B$ of a 6 by 8 rectangle $ABCD$ from the diagonal $AC$? You could set up coordinates and find the equation of a line, but computing an area twice is shorter. The diagonal has length 10, because 6, 8, 10 is a right-angled triangle. Triangle $ABC$ has area $\frac{1}{2} \times 6 \times 8 = 24$. Its area is also $\frac{1}{2} \times AC \times h = 5h$, where $h$ is the distance we want. So $h = \frac{24}{5}$.

This unit covers computing one area in two ways, Pythagoras and its triples, and similar shapes. Between them they cover much of the geometry on the IMC, the Kangaroos and the AMC 10.

### 1.1 Area


The area of a triangle is $\frac{1}{2} \times$ base $\times$ height, where the height is measured perpendicular to the base, and may fall outside the triangle. We use three consequences constantly.

**Area facts.**
(1) Triangles on the same base with the same height have equal areas, so sliding the apex parallel to the base does not change the area. (2) Triangles with the same height have areas in the ratio of their bases. (3) A triangle has half the area of a parallelogram on the same base and between the same parallels.

When a region is awkward, find it by subtracting from something simple, such as the rectangle around it, or by cutting it into pieces whose bases and heights you know. Fact (2) is the one people forget: if $D$ lies on $BC$ with $BD : DC = 2 : 3$, then triangles $ABD$ and $ADC$ have areas in the ratio $2 : 3$, whatever the shape of the triangle, because they share the height from $A$.



Kevinvariant
kevinvariant08.github.io




**Example 1.**
Rectangle $ABCD$ has $AB = 12$ and $BC = 8$. $M$ is the midpoint of $BC$, and $N$ lies on $CD$ with $CN = 3$. Find the area of triangle $AMN$.

*Solution.*
Using $\frac{1}{2} \times$ base $\times$ height directly needs the distance from $N$ to the line $AM$, which is unpleasant to find. But the triangle lies inside the rectangle, and the three pieces left over are right-angled triangles with their legs along the sides, so we subtract their areas.
Triangle $ABM$ has legs 12 and 4, so area 24. Triangle $MCN$ has legs 4 and 3, so area 6. Triangle $ADN$ has legs 8 and 9, so area 36. The rectangle has area 96, so triangle $AMN$ has area $96 - 24 - 6 - 36 = 30$.
This method (enclose and subtract) works for any triangle whose vertices lie on the sides of a rectangle, and for any triangle drawn on a square grid: draw the smallest rectangle round it and remove the right-angled triangles (and any rectangles) in the corners.


## 1.2 Pythagoras and the triples worth knowing

**Pythagoras' theorem.**
In a right-angled triangle with legs $a$, $b$ and hypotenuse $c$, we have $a^2 + b^2 = c^2$. Conversely, if $a^2 + b^2 = c^2$ then the angle opposite $c$ is a right angle.

I will give a proof by area. Place four copies of the triangle inside a square of side $a + b$, with their right angles in the corners. At each point where two triangles meet on a side of the big square, they contribute the two different acute angles of the triangle, which add to $90°$, so the angle of the hole there is $180° - 90° = 90°$. The four sides of the hole are hypotenuses, so the hole is a square of side $c$. Counting the area two ways, $(a + b)^2 = c^2 + 4 \times \frac{1}{2}ab$, which simplifies to $a^2 + b^2 = c^2$.




Kevinvariant
kevinvariant08.github.io




Whole-number solutions are called Pythagorean triples. These are worth remembering: $(3,4,5)$, $(5,12,13)$, $(8,15,17)$, $(7,24,25)$, $(20,21,29)$ and $(9,40,41)$, together with multiples such as $(6,8,10)$ and $(15,20,25)$. You can generate more from $(m^2-n^2,\ 2mn,\ m^2+n^2)$, since $(m^2-n^2)^2+(2mn)^2 = (m^2+n^2)^2$; in fact every triple with no common factor arises this way, though that takes more work to prove.

Two right-angled triangles with known angles are worth memorising. Half a square has sides $1, 1, \sqrt{2}$ and angles $45°, 45°, 90°$. Half an equilateral triangle has sides $1, \sqrt{3}, 2$ and angles $30°, 60°, 90°$. Finally, one algebraic move: $c^2 - a^2 = b^2$ factorises as $(c-a)(c+a) = b^2$, which is the way to find whole-number solutions with one side fixed.




**Example 2.**
Rectangle $ABCD$ has $AB = 6$ and $BC = 10$. It is folded along a line $AE$, with $E$ on $CD$, so that $D$ lands on a point $D'$ of side $BC$. Find $DE$.

*Solution.*
A fold is a reflection in the crease, so lengths and angles are preserved: $AD' = AD = 10$, $ED' = ED$, and $\angle AD'E = \angle ADE = 90°$. We start with what is fully determined. In the right-angled triangle $ABD'$ we know $AB = 6$ and the hypotenuse $AD' = 10$, so $BD' = 8$ (a $6, 8, 10$ triangle) and $D'C = 2$.
Now let $DE = x$. Then $ED' = x$ and $EC = 6 - x$, and Pythagoras in triangle $ECD'$ gives $(6-x)^2 + 2^2 = x^2$. Expanding, $40 - 12x = 0$, so $x = \frac{10}{3}$.
There is also a method with no algebra. Since $\angle AD'E = 90°$, the angles $\angle AD'B$ and $\angle ED'C$ add to $90°$, so $\angle ED'C = \angle BAD'$. Triangles $ABD'$ and $D'CE$ therefore have the same angles, and the ratio of corresponding sides is $D'C : AB = 2 : 6 = 1 : 3$. So $ED' = \frac{1}{3}AD' = \frac{10}{3}$.
Right angles that meet on a straight line almost always produce similar triangles like this.




Kevinvariant
kevinvariant08.github.io

## 1.3 Similar shapes

Two triangles are similar if their angles are equal in pairs; then all corresponding lengths are in the same ratio. For triangles, two pairs of equal angles are enough, since the third pair follows. Write the vertices in matching order: if $\triangle ABC$ is similar to $\triangle PQR$, with $\angle A = \angle P$ and so on, then $\dfrac{AB}{PQ} = \dfrac{BC}{QR} = \dfrac{CA}{RP}$.

**Scale factors.**
If lengths are multiplied by $k$, areas are multiplied by $k^2$ and volumes by $k^3$.

For a triangle this is clear, since the base and the height are both multiplied by $k$, and any polygon is made of triangles. Three configurations come up often. A line parallel to one side of a triangle cuts off a smaller triangle similar to the whole (the A shape). Two parallel segments with their ends joined crosswise give two similar triangles meeting at a point (the X shape), which is what the diagonals of a trapezium do. And the altitude to the hypotenuse of a right-angled triangle splits it into two triangles similar to the whole; if the altitude from $C$ meets $AB$ at $D$, this gives $CD^2 = AD \times DB$.





**Example 3.**
Triangle $ABC$ has a right angle at $C$, with $CA = 8$ and $CB = 6$. A square $PQRS$ has $P$ and $Q$ on $AB$, $R$ on $CB$ and $S$ on $CA$. Find the side of the square.

*Solution.*
First, $AB = 10$. Coordinates would work, but the square is tilted and the equations are long. Instead, consider the three small triangles round the square. Each is right-angled and shares an angle with $ABC$, so each is similar to $ABC$.
Let the side be $s$. Triangle $APS$ has a right angle at $P$ and the angle $A$ of the big triangle, so $\dfrac{AP}{PS} = \dfrac{AC}{CB} = \dfrac{8}{6}$, giving $AP = \frac{4}{3}s$. Triangle $BQR$ has a right angle at $Q$ and the angle $B$, so $\dfrac{BQ}{QR} = \dfrac{BC}{CA} = \dfrac{6}{8}$, giving $BQ = \frac{3}{4}s$. Along the hypotenuse, $\frac{4}{3}s + s + \frac{3}{4}s = 10$, so $\frac{37}{12}s = 10$ and $s = \frac{120}{37}$.
As a check, use the A shape. The altitude from $C$ to $AB$ is $\frac{6\times 8}{10} = \frac{24}{5}$ (area two ways again). Triangle $CSR$ is similar to $CAB$ because $SR$ is parallel to $AB$, and its height is $\frac{24}{5} - s$, so



Kevinvariant
kevinvariant08.github.io

$\frac{s}{10} = \frac{24/5 - s}{24/5}$. This also gives $s = \frac{120}{37}$. For comparison, the square standing in the corner at $C$ has side $\frac{24}{7}$, which is larger.




**Example 4.**
$ABCD$ is a trapezium with $AB$ parallel to $DC$. Its diagonals meet at $X$. Triangle $AXB$ has area 50 and triangle $CXD$ has area 8. Find the area of the trapezium.

*Solution.*
No lengths are given, and $AB$, $CD$ and the height cannot be found, since infinitely many trapezia fit the data. So the answer must come from ratios alone.
Since $AB$ is parallel to $DC$, alternate angles give $\angle XAB = \angle XCD$ and $\angle XBA = \angle XDC$, so triangles $AXB$ and $CXD$ are similar (the X shape). Their areas are in the ratio $50 : 8 = 25 : 4$, so their lengths are in the ratio $5 : 2$. In particular $AX : XC = 5 : 2$.
Now we use a shared height. Triangles $AXD$ and $CXD$ have their bases $AX$ and $XC$ on the same line and share the apex $D$, so their areas are in the ratio $5 : 2$, which makes the area of $AXD$ equal to 20. Similarly triangle $BXC$ has area 20. (Alternatively, triangles $ABD$ and $ABC$ have the same base and the same height, so removing $AXB$ from each shows that $AXD$ and $BXC$ have equal areas.) The trapezium has area $50 + 8 + 20 + 20 = 98$.
In general, if the two triangles on the parallel sides have areas $P$ and $Q$, each side triangle has area $\sqrt{PQ}$ and the trapezium has area $(\sqrt{P} + \sqrt{Q})^2$.

**Common traps.**

- Scaling areas by $k$ instead of $k^2$. Lengths in the ratio $2 : 3$ give areas in the ratio $4 : 9$.
- Matching the wrong sides in similar triangles. Pair up the equal angles first, then read off the sides opposite them.
- Using Pythagoras without a right angle, or with the wrong side as the hypotenuse: it is always the side opposite the right angle.
- Taking a slanted length as a height. The height must be perpendicular to the base.
- Seeing a triple that is not there. Check that $a^2 + b^2 = c^2$ before you trust it.

## 1.4 How to spot this in a contest

If right angles appear and lengths are asked for, look for Pythagoras and for triples first, since setters often choose numbers that form triples. If an area is asked for and the region is awkward,



Kevinvariant
kevinvariant08.github.io

subtract it from a rectangle, or compare triangles that share a height. If there are parallel lines, there are similar triangles: mark the equal angles and write down the ratio. If the question gives areas and asks for areas, think in ratios: find a length ratio by similarity, square it, and use shared heights for the rest. A fold is a reflection, and folding problems almost always lead to a Pythagoras equation in one unknown. Whenever you need the distance from a point to a line, compute an area two ways.

## 2 Practice problems

The problems are roughly in order of difficulty, from ⋆ to ⋆⋆⋆⋆. Hints are in Section 4 and full solutions in Section 5.

**Problem 1** ⋆ A right-angled triangle has hypotenuse 25 and one leg 7. Find its area.

**Problem 2** ⋆⋆ $P$ is a point inside rectangle $ABCD$. Triangles $PAB$, $PBC$ and $PCD$ have areas 13, 9 and 20. Find the area of triangle $PDA$.

**Problem 3** ⋆⋆ Square $ABCD$ has side 6. $E$ is the point on $BC$ with $BE = 2$, and $AE$ meets the diagonal $BD$ at $F$. Find the area of triangle $BEF$.

**Problem 4** ⋆⋆⋆ How many right-angled triangles with whole-number side lengths have a leg of length 20? (Triangles that are congruent count once.)

**Problem 5** ⋆⋆⋆ Rectangle $ABCD$ has $AB = 15$ and $BC = 8$. Points $P$ on $AB$ and $Q$ on $CD$ are chosen so that $APCQ$ is a rhombus. Find the length $PQ$.

**Problem 6** ⋆⋆⋆⋆ A right-angled triangle has perimeter 120, and the altitude from the right angle to the hypotenuse has length 24. Find the area of the triangle.

## 3 Contest ladder

Real past problems on the ideas of this unit, easiest first. Each source is a link to the official paper or the AoPS wiki; the note under it says where to start, not how to finish.

1. **AMC 10B 2006 Problem 8** ⋆
   *Join the centre of the semicircle to a top corner of the square.*
2. **UKMT IMC 2018 Question 8** ⋆
   *The corner triangles are right-angled: think of a familiar triple.*
3. **UKMT IMC 2017 Question 13** ⋆⋆
   *Use the ratio of the two similar triangles.*
4. **AMC 10A 2018 Problem 13** ⋆⋆
   *The crease is the perpendicular bisector of the hypotenuse.*
5. **AMC 10A 2018 Problem 9** ⋆⋆
   *Areas of similar triangles scale with the square of the ratio.*



Kevinvariant
kevinvariant08.github.io

6. **AMC 10A 2008 Problem 20** ⋆⋆⋆
   *X shape at the diagonals, then triangles sharing a height.*
7. **UKMT IMC 2018 Question 22** ⋆⋆⋆
   *The gaps between the L-shape and the rectangle are similar right-angled triangles.*
8. **AMC 10A 2016 Problem 19** ⋆⋆⋆
   *Similar triangles on either side of the diagonal.*
9. **UKMT Maclaurin 2026 Question 2** ⋆⋆⋆
   *Pythagoras for the cut, area for the product; check for extra roots.*
10. **AMC 10B 2021 Problem 21** ⋆⋆⋆
    *A fold is a reflection: Pythagoras in one unknown.*
11. **UKMT Maclaurin 2026 Question 5** ⋆⋆⋆⋆
    *Look for similar triangles using the parallel sides.*
12. **UKMT Hamilton 2026 Question 4** ⋆⋆⋆⋆
    *Look for two different pairs of similar triangles.*



Kevinvariant
kevinvariant08.github.io


## 4 Hints

1. Find the other leg with Pythagoras, or recognise the triple.
2. Add the areas of the two triangles on opposite sides. What do their heights add up to?
3. $AD$ is parallel to $BE$, so look for an X shape at $F$.
4. If the other leg is $b$ and the hypotenuse is $c$, factorise $c^2 - b^2$.
5. Find the side of the rhombus with Pythagoras, then compute the rhombus's area in two ways.
6. Call the legs $a$, $b$ and the hypotenuse $c$. The area gives one equation, and squaring $a + b$ brings in Pythagoras.

## 5 Solutions

### Problem 1

*A right-angled triangle has hypotenuse* 25 *and one leg* 7*. Find its area.*





The other leg is $\sqrt{25^2 - 7^2} = \sqrt{625 - 49} = \sqrt{576} = 24$, so this is the $(7, 24, 25)$ triple. The legs are perpendicular, so the area is $\frac{1}{2} \times 7 \times 24 = 84$.

**Answer:** 84

### Problem 2

*$P$ is a point inside rectangle $ABCD$. Triangles $PAB$, $PBC$ and $PCD$ have areas* 13, 9 *and* 20*. Find the area of triangle $PDA$.*






Kevinvariant
kevinvariant08.github.io


Triangles $PAB$ and $PCD$ have bases $AB$ and $CD$, both equal to the length of the rectangle, and their heights are the distances from $P$ to the opposite sides $AB$ and $CD$, which add up to $BC$. So their areas add to $\frac{1}{2} \times AB \times BC$, half the rectangle. The same is true of triangles $PBC$ and $PDA$. Hence, writing $x$ for the area of $PDA$, we get $13 + 20 = 9 + x$, so $x = 24$.

(Such a point exists: for example, take $AB = 11$ and $BC = 6$, with $P$ at distance 3 from $BC$ and $\frac{26}{11}$ from $AB$.)

**Answer:** 24

### Problem 3

*Square $ABCD$ has side 6. $E$ is the point on $BC$ with $BE = 2$, and $AE$ meets the diagonal $BD$ at $F$. Find the area of triangle $BEF$.*




Since $AD$ is parallel to $BE$, triangles $FAD$ and $FEB$ have equal alternate angles and equal vertically opposite angles at $F$, so they are similar with ratio $AD : EB = 6 : 2 = 3 : 1$. Hence $AF : FE = 3 : 1$, and $FE$ is a quarter of $AE$.

Triangles $BEF$ and $BEA$ share the apex $B$, and their bases $FE$ and $AE$ lie on the same line, so the ratio of their areas is $FE : AE = 1 : 4$. Triangle $ABE$ has legs 6 and 2 and area 6, so triangle $BEF$ has area $\frac{6}{4} = \frac{3}{2}$.


**Answer:** $\frac{3}{2}$

### Problem 4

*How many right-angled triangles with whole-number side lengths have a leg of length 20? (Triangles that are congruent count once.)*



Kevinvariant
kevinvariant08.github.io





Let the other leg be $b$ and the hypotenuse $c$. Then $c^2 - b^2 = 400$, that is $(c-b)(c+b) = 400$. The two factors have the same parity (they differ by $2b$), and their product is even, so both are even. Write $c - b = 2u$ and $c + b = 2v$ with $u < v$; then $uv = 100$, and each such factor pair gives $b = v - u$ and $c = u + v$.

The pairs with $u < v$ are $(1, 100)$, $(2, 50)$, $(4, 25)$ and $(5, 20)$; the pair $(10, 10)$ would give $b = 0$. They give the triangles $(20, 99, 101)$, $(20, 48, 52)$, $(20, 21, 29)$ and $(15, 20, 25)$, so there are 4.

**Answer:** 4

### Problem 5

*Rectangle $ABCD$ has $AB = 15$ and $BC = 8$. Points $P$ on $AB$ and $Q$ on $CD$ are chosen so that $APCQ$ is a rhombus. Find the length $PQ$.*




Let the side of the rhombus be $x$, so $AP = PC = x$ and $PB = 15 - x$. Pythagoras in triangle $PBC$ gives $(15 - x)^2 + 8^2 = x^2$, so $289 - 30x = 0$ and $x = \frac{289}{30}$.

The rhombus is a parallelogram with base $AP$ and height $BC$, so its area is $\frac{289}{30} \times 8 = \frac{1156}{15}$. Its diagonals are perpendicular, so its area is also $\frac{1}{2} \times AC \times PQ$, and $AC = 17$ (an $8, 15, 17$ triangle). Hence $PQ = \dfrac{2 \times 1156}{15 \times 17} = \frac{136}{15}$.


**Answer:** $\frac{136}{15}$



Kevinvariant
kevinvariant08.github.io

### Problem 6

*A right-angled triangle has perimeter* 120*, and the altitude from the right angle to the hypotenuse has length* 24*. Find the area of the triangle.*




Let the legs be $a$ and $b$ and the hypotenuse $c$. Computing the area in two ways gives $\frac{1}{2}ab = \frac{1}{2} \times 24c$, so $ab = 24c$. The perimeter gives $a + b = 120 - c$.

Squaring, $a^2 + 2ab + b^2 = (120 - c)^2$. By Pythagoras $a^2 + b^2 = c^2$, so $c^2 + 48c = 14400 - 240c + c^2$, which gives $288c = 14400$ and $c = 50$. The area is $\frac{1}{2} \times 24 \times 50 = 600$. (The legs are 30 and 40: they satisfy $a + b = 70$ and $ab = 1200$.)

**Answer:** 600