

Spherification

Monday, April 28, 2025

10:49 PM



Soda or Juice Caviar Recipe

Ingredients

- 200 g. Fruit juice or soda (ph above 3.7%)
- 0.6 - 1.0 g. Sodium alginate
- 500 g. Tap water
- 1 g. Calcium chloride
- 6 fl. oz. Tap water as needed

Tools

- Stick blender, mixing tool
- Pipettes
- Spoons, slotted
- Bowls
- Tasting cups/spoons

Preparation

- Using a hand-held blender or mixing tool, mix the soda or flavored liquid and sodium alginate together. This will take longer than expected, 5-10 minutes.

Let rest to remove any trapped air bubbles. For optimum solution, let rest overnight.

- Using the blender or mixing tool, combine 500 g. water and calcium chloride until fully mixed. For optimum solution, let rest in refrigerator overnight.
- Set up a spherification station: soda + sodium alginate mixture; calcium chloride bath; tap water rinse; and holding vessel.
-
- These are time sensitive. Use within 15 minutes for maximum effect.

Procedure

- Fill pipettes with the juice or soda and sodium alginate solution.
- Gently squeeze the pipettes (syringes or squeeze bottles could also work) to release the solution drop by drop into calcium chloride bath. Be careful not to drop the liquid from too great of a height or they will flatten and resemble small pancakes instead of caviar spheres.
- After you have dropped a small quantity into the bath, remove the spheres from the calcium chloride solution with a small slotted spoon.
- Blot the bottom of the spoon with a towel to remove the calcium chloride solution.
- Place spheres in the water rinsing bath.
- Immediately after rinsing, remove the spheres from the tap water and sample the results.

Reverse Spherification - Yogurt Ravioli Recipe

Ingredients

- 1500 g. Distilled water
- 7.5 g. Sodium alginate
- 200 g. Yogurt
- 4 g. Calcium lactate gluconate
- 250 ml. Tap water

Tools

- Stick blender, mixing tool
- Pipettes
- Spoons, slotted

spoons, slotted

- Bowls
- Tasting cups/spoons

Preparation

- Using a hand-held blender or mixing tool, mix the distilled water and sodium alginate together. This will take longer than expected, 5-10 minutes. Let rest to remove any trapped air bubbles. For optimum solution, let rest overnight. Use distilled water if the calcium content is high in your tap water.
- Mix the yogurt with calcium lactate gluconate to boost the calcium content.
- Set up the three-part station: Yogurt + calcium lactate gluconate mixture; sodium alginate bath; tap water rinse; and holding vessel.
-

Procedure

- Use a small spoon (a 1/2 teaspoon measure) and gently turn the spoon so the yogurt will enter the sodium alginate bath.
- Let the sphere sit in the solution for up to two minutes so the membrane will form. Make sure not to let the spheres touch each other or they will stick together.
- Remove the sphere with a slotted spoon; make sure the sodium alginate solution has drained from the spoon.
- Place the sphere in the water to rinse the salt byproduct.
- Sample or store in juice until ready to sample.