

## MENTAL IMPAIRMENT QUESTIONNAIRE (LISTINGS)

Completed by: \_\_\_\_\_

Patient: \_\_\_\_\_

Date Completed: \_\_\_\_\_

Patient DOB: \_\_\_\_\_

Please answer the following questions concerning your patient's impairments.

1. Frequency and length of contact: \_\_\_\_\_

\_\_\_\_\_

2. DSM-5 Mental Disorders/Principal Diagnoses:

Other Conditions / Focus of Clinical  
Attention:

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

3. Treatment and response: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

4. a. List prescribed medications:

b. Identify your patient's side effects of medications that may affect working, *e.g.*, sedation, drowsiness, fatigue, lethargy, malaise, irritability, nausea, dizziness, *etc.*

5. List the *clinical findings* including those from mental status examination that demonstrate the severity of your patient's mental impairment and symptoms:

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

6. Prognosis: \_\_\_\_\_

7. Identify your patient's signs and symptoms:

|                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                                                                                                                                                                                          |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Significant cognitive decline from a prior level of functioning in <i>one</i> or more of the cognitive areas: Complex attention; Executive function; Learning and memory; Language; Perceptual-motor; or Social cognition.                                                                                                                                                                                                                                | Hyperactive and impulsive behavior (e.g., difficulty remaining seated, talking excessively, difficulty waiting, appearing restless, or behaving as if being “driven by a motor”)         |
| Delusions or hallucinations                                                                                                                                                                                                                                                                                                                                                                                                                               | Irritability                                                                                                                                                                             |
| Qualitative deficits in verbal communication, nonverbal communication, and social interaction                                                                                                                                                                                                                                                                                                                                                             | Preoccupation with having or acquiring a serious illness without significant symptoms present                                                                                            |
| Increase in goal-directed activity or psychomotor agitation                                                                                                                                                                                                                                                                                                                                                                                               | Disregard for and violation of the rights of others                                                                                                                                      |
| Depressed mood                                                                                                                                                                                                                                                                                                                                                                                                                                            | Detachment from social relationships                                                                                                                                                     |
| Diminished interest in almost all activities                                                                                                                                                                                                                                                                                                                                                                                                              | Distrust and suspiciousness of others                                                                                                                                                    |
| Restlessness                                                                                                                                                                                                                                                                                                                                                                                                                                              | Instability of interpersonal relationships                                                                                                                                               |
| Sleep disturbance                                                                                                                                                                                                                                                                                                                                                                                                                                         | Pressured speech                                                                                                                                                                         |
| Observable psychomotor agitation or retardation                                                                                                                                                                                                                                                                                                                                                                                                           | Preoccupation with perfectionism and orderliness                                                                                                                                         |
| Decreased energy                                                                                                                                                                                                                                                                                                                                                                                                                                          | Easily fatigued                                                                                                                                                                          |
| Feelings of guilt or worthlessness                                                                                                                                                                                                                                                                                                                                                                                                                        | Muscle tension                                                                                                                                                                           |
| Difficulty concentrating or thinking                                                                                                                                                                                                                                                                                                                                                                                                                      | Disorganized thinking (speech)                                                                                                                                                           |
| Thoughts of death or suicide                                                                                                                                                                                                                                                                                                                                                                                                                              | Recurrent motor movement or vocalization                                                                                                                                                 |
| Excessive emotionality and attention seeking                                                                                                                                                                                                                                                                                                                                                                                                              | Repetitive behaviors aimed at reducing anxiety                                                                                                                                           |
| Flight of ideas                                                                                                                                                                                                                                                                                                                                                                                                                                           | Inflated self-esteem                                                                                                                                                                     |
| Decreased need for sleep                                                                                                                                                                                                                                                                                                                                                                                                                                  | Distractibility                                                                                                                                                                          |
| Involvement in activities that have a high probability of painful consequences that are not recognized                                                                                                                                                                                                                                                                                                                                                    | One or more somatic symptoms that are distressing, with excessive thoughts, feelings, or behaviors related to the symptoms                                                               |
| Grossly disorganized behavior or catatonia                                                                                                                                                                                                                                                                                                                                                                                                                | Appetite disturbance with change in weight                                                                                                                                               |
| Panic attacks followed by a persistent concern or worry about additional panic attacks or their consequences                                                                                                                                                                                                                                                                                                                                              | Symptoms of altered voluntary motor or sensory function that are not better explained by another medical or mental disorder                                                              |
| Disproportionate fear or anxiety about at least two different situations (for example, using public transportation, being in a crowd, being in a line, being outside of your home, being in open spaces)                                                                                                                                                                                                                                                  | Persistent alteration in eating or eating-related behavior that results in a change in consumption or absorption of food and that significantly impairs physical or psychological health |
| Recurrent, impulsive, aggressive behavioral outbursts                                                                                                                                                                                                                                                                                                                                                                                                     | Significantly restricted, repetitive patterns of behavior, interests, or activities                                                                                                      |
| Frequent distractibility, difficulty sustaining attention, and difficulty organizing tasks                                                                                                                                                                                                                                                                                                                                                                | Significant difficulties learning and using academic skills                                                                                                                              |
| Involuntary, time-consuming preoccupation with intrusive, unwanted thought                                                                                                                                                                                                                                                                                                                                                                                |                                                                                                                                                                                          |
| Medical documentation of <i>all</i> of the following: 1. Exposure to actual or threatened death, serious injury, or violence; 2. Subsequent involuntary re-experiencing of the traumatic event (for example, intrusive memories, dreams, or flashbacks); 3. Avoidance of external reminders of the event; 4. Disturbance in mood and behavior; and 5. Increases in arousal and reactivity (for example, exaggerated startle response, sleep disturbance). |                                                                                                                                                                                          |

8. Is your patient's intellectual functioning limited?

☐ Yes ☐ No

If yes, please explain:

9. Rate the degree of your patient's expected limitations in a work setting using the following scale:

**Moderate** means the ability to function independently, appropriately, effectively, and on a sustained basis is **fair**;

**Marked** means the ability to function independently, appropriately, effectively, and on a sustained basis is **seriously limited**;

**Extreme** means **not able** to function independently, appropriately, effectively, and on a sustained basis, but it does not mean a total loss of ability to function.

| RATE THE DEGREE OF LIMITATION<br>IN THE FOLLOWING AREAS: |                                   |           |          |        |         |
|----------------------------------------------------------|-----------------------------------|-----------|----------|--------|---------|
| A.1.                                                     | Understanding information:        | None-Mild | Moderate | Marked | Extreme |
| A.2.                                                     | Remembering information           | None-Mild | Moderate | Marked | Extreme |
| A.3.                                                     | Applying information              | None-Mild | Moderate | Marked | Extreme |
| B.                                                       | Interacting with others:          | None-Mild | Moderate | Marked | Extreme |
| C.1.                                                     | Concentrating                     | None-Mild | Moderate | Marked | Extreme |
| C.2.                                                     | Persisting                        | None-Mild | Moderate | Marked | Extreme |
| C.3.                                                     | Maintaining pace                  | None-Mild | Moderate | Marked | Extreme |
| D.1.                                                     | Adapting in the workplace         | None-Mild | Moderate | Marked | Extreme |
| D.2.                                                     | Managing oneself in the workplace | None-Mild | Moderate | Marked | Extreme |

10. If your patient has a neurocognitive disorder; schizophrenia or other psychotic disorder; depressive, bipolar or related disorder; anxiety or obsessive compulsive disorder; or trauma or stress-related disorder, please indicate if any of the following apply to your patient:
- ☐ Your patient's chronic mental disorder is "serious and persistent;" that is, your patient has a medically documented history of the existence of the disorder over a period of at least 2 years.
  - ☐ Your patient relies on ongoing medical treatment, mental health therapy, psychosocial support, or a highly-structured setting to diminish the symptoms and signs of his or her mental disorder.
  - ☐ Despite your patient's diminished symptoms and signs, your patient has only marginal adjustment, that is, your patient has minimal capacity to adapt to changes in his or her environment or to demands that are not already part of daily life.
11. Assuming your patient was trying to work full time, on the average, how often do you anticipate that your patient's impairments would cause your patient to be absent from work?
- ☐ Never                      ☐ About two days per month                      ☐ About four days per month  
☐ About one day per month                      ☐ About three days per month                      ☐ More than four days per month
12. Has your patient's impairment lasted or can it be expected to last at least twelve months?
- ☐ Yes                      ☐ No
13. Please describe any additional reasons not covered above why your patient would have difficulty working at a regular job on a sustained basis.
14. Is your patient currently abusing substances?
- ☐ Yes                      ☐ No
15. If there has been substance abuse in the past, what is the patient's date of sobriety?
- \_\_\_\_\_
16. How would the patient's sobriety (potential or current) impact their impairments?
17. Can your patient manage benefits in his or her own best interest?
- ☐ Yes                      ☐ No
18. Has the patient been given either of these tests? If so, please record the result.
- CDR                      ☐ No                      ☐ Yes                      Results \_\_\_\_\_
- MMSE                      ☐ No                      ☐ Yes                      Results \_\_\_\_\_

19. What is the earliest date that the description of *symptoms and limitations* in this questionnaire applies? \_\_\_\_\_

20. Comments:

---

---

---

---

---

---

\_\_\_\_\_  
Date

\_\_\_\_\_  
Signature

Printed/Typed Name: \_\_\_\_\_

Address: \_\_\_\_\_

\_\_\_\_\_

NOTE: This form was provided to the above named medical care provider by representing attorney Geri N. Kahn to be completed in support of our client's disability claim.

RETURN TO YOUR PATIENT OR FAX TO 707-312-8271