

Contents lists available at SciVerse ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb



Centrality dependence of charged particle production at large transverse momentum in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV

ALICE Collaboration

ARTICLE INFO

Article history:
Received 11 September 2012
Received in revised form 21 January 2013
Accepted 24 January 2013
Available online 31 January 2013
Editor: V. Metag

ABSTRACT

The inclusive transverse momentum (p_T) distributions of primary charged particles are measured in the pseudo-rapidity range $|\eta| < 0.8$ as a function of event centrality in Pb-Pb collisions at $\sqrt{s_{\rm NN}} = 2.76$ TeV with ALICE at the LHC. The data are presented in the p_T range $0.15 < p_T < 50$ GeV/c for nine centrality intervals from 70–80% to 0–5%. The results in Pb-Pb are presented in terms of the nuclear modification factor $R_{\rm AA}$ using a pp reference spectrum measured at the same collision energy. We observe that the suppression of high- p_T particles strongly depends on event centrality. The yield is most suppressed in central collisions (0–5%) with $R_{\rm AA} \approx 0.13$ at $p_T = 6-7$ GeV/c. Above $p_T = 7$ GeV/c, there is a significant rise in the nuclear modification factor, which reaches $R_{\rm AA} \approx 0.4$ for $p_T > 30$ GeV/c. In peripheral collisions (70–80%), only moderate suppression ($R_{\rm AA} = 0.6$ –0.7) and a weak p_T dependence is observed. The measured nuclear modification factors are compared to other measurements and model calculations.

1. Introduction

High-energy collisions of heavy-ions enable the study of hot and dense strongly interacting matter [1–5]. At sufficiently high temperature, it is expected that partons (quarks and gluons) are the dominant degrees of freedom. During the very early stage of the collision, some of the incoming partons experience scatterings with large momentum transfers. These partons lose energy when they traverse the hot and dense medium that is formed. One of the major goals of the heavy-ion physics programme at the LHC is to understand the underlying mechanisms for parton energy loss and use this as a tool to probe the properties of the medium.

Parton energy loss in heavy-ion collisions was first observed at RHIC as the suppression of high- $p_{\rm T}$ particle production in Au–Au collisions compared to expectations from an independent superposition of nucleon–nucleon collisions [6–9]. At RHIC, the particle production in central (0–5%) Au–Au collisions at $\sqrt{s_{\rm NN}}=200$ GeV is suppressed by a factor of 5 at $p_{\rm T}=5$ –6 GeV/c [8,9], and is consistent with being independent of $p_{\rm T}$ over the measured range $5 < p_{\rm T} < 20$ GeV/c [10]. At RHIC, RAA

The increase of the charged particle density $(dN_{ch}/d\eta)$ at midrapidity from RHIC energies to actual LHC energies by a factor of around 2.2 [11] implies a similar increase in energy density. However, the observed suppression of high- p_T particle production also depends on the ratio of quarks to gluons due to their different color factors, and on the steepness of the p_T spectra of the scattered partons. At the LHC the initial parton p_T spectra are less Km, NR NR

steep than at RHIC and the ratio of gluons to quarks at a given p_T is higher [12]. The measurement of high- p_T hadron production at the LHC helps to disentangle the effects which cause the suppression and provides a critical test of existing energy loss calculations [13]. In particular, the large p_T reach provides a means to study the dependence of the energy loss on the initial parton energy.

We present a measurement of the $p_{\rm T}$ distributions of charged particles in 0.15 < $p_{\rm T}$ < 50 GeV/c with pseudo-rapidity $|\eta|$ < 0.8, where $\eta=-\ln[\tan(\theta/2)]$, with θ the polar angle between the charged particle direction and the beam axis. Results are presented for different centrality intervals in Pb–Pb collisions at $\sqrt{s_{\rm NN}}=2.76$ TeV. They are compared with measurements in pp collisions, by calculating the nuclear modification factor

$$R_{\rm AA}(p_{\rm T}) = \frac{\mathrm{d}^2 N_{\rm ch}^{\rm AA}/\mathrm{d}\eta \,\mathrm{d}p_{\rm T}}{\langle T_{\rm AA}\rangle \,\mathrm{d}^2 \sigma_{\rm ch}^{\rm pp}/\mathrm{d}\eta \,\mathrm{d}p_{\rm T}} \tag{1}$$

where $N_{\rm ch}^{\rm AA}$ and $\sigma_{\rm ch}^{\rm pp}$ represent the charged particle yield in nucleus–nucleus (AA) collisions and the cross section in pp collisions, respectively. The nuclear overlap function $T_{\rm AA}$ is calculated from the Glauber model [14] and averaged over each centrality interval, $\langle T_{\rm AA} \rangle = \langle N_{\rm coll} \rangle / \sigma_{\rm inel}^{\rm NN}$, where $\langle N_{\rm coll} \rangle$ is the average number of binary nucleon–nucleon collisions and $\sigma_{\rm inel}^{\rm NN}$ is the inelastic nucleon–nucleon cross section.

Early results from ALICE [15] showed that the production of charged particles in central (0–5%) Pb–Pb collisions at $\sqrt{s_{\rm NN}}=2.76$ TeV is suppressed by more than a factor of 6 at $p_{\rm T}=6-7$ GeV/c compared to an independent superposition of nucleonnucleon collisions, and that the suppression is stronger than that observed at RHIC. The present data extend the study of high- $p_{\rm T}$

Table 1 Average values of the number of participating nucleons $\langle N_{\text{part}} \rangle$ and the nuclear overlap function $\langle T_{\text{AA}} \rangle$ [14] for the centrality intervals used in the analysis.

Centrality	$\langle N_{ m part} \rangle$	$\langle T_{\rm AA} \rangle \ ({\rm mb}^{-1})$
0-5%	383 ± 3	26.4 ± 1.1
5-10%	330 ± 5	20.6 ± 0.9
10-20%	261 ± 4	14.4 ± 0.6
20-30%	186 ± 4	8.7 ± 0.4
30-40%	129 ± 3	5.0 ± 0.2
40-50%	85 ± 3	2.68 ± 0.14
50-60%	53 ± 2	1.32 ± 0.09
60-70%	30.0 ± 1.3	0.59 ± 0.04
70-80%	15.8 ± 0.6	0.24 ± 0.03

particle suppression in Pb–Pb out to $p_T = 50 \text{ GeV}/c$ with a systematic study of the centrality dependence.

Moreover, the systematic uncertainties related to the pp reference were significantly reduced with respect to the previous measurement by using the p_T distribution measured in pp collisions at $\sqrt{s} = 2.76$ TeV [16].

2. Experiment and data analysis

The ALICE detector is described in [17]. The Inner Tracking System (ITS) and the Time Projection Chamber (TPC) are used for vertex finding and tracking. The minimum-bias interaction trigger was derived from signals from the forward scintillators (VZERO), and the two innermost layers of the ITS (Silicon Pixel Detector — SPD). The collision centrality is determined using the VZERO. In addition, the information from two neutron Zero Degree Calorimeters (ZDCs) positioned at ± 114 m from the interaction point was used to remove contributions from beam-gas and electromagnetic interactions. The trigger and centrality selection are described in more detail in [11]. 事件被分为9个中心度,以占总强子散射截面的百分比表示。

The following analysis is based on $1.6 \cdot 10^7$ minimum-bias Pb-Pb events recorded by ALICE in 2010. For this study, the events are divided into nine centrality intervals from the 70–80% to the 0–5% most central Pb-Pb collisions, expressed in percentage of the total hadronic cross section. The event centrality can be related to the number of participating nucleons $N_{\rm part}$ and the nuclear overlap function $T_{\rm AA}$ by using simulations based on the Glauber model [14]. The average values of $N_{\rm part}$ and $T_{\rm AA}$ for each centrality interval, $\langle N_{\rm part} \rangle$ and $\langle T_{\rm AA} \rangle$, along with their corresponding systematic uncertainties, are listed in Table 1. The errors include the experimental uncertainties on the inelastic nucleon-nucleon cross section $\sigma_{\rm inel}^{\rm NN} = 64 \pm 5$ mb at $\sqrt{s_{\rm NN}} = 2.76$ TeV [18] and on the parameters of the nuclear density profile used in the Glauber simulations (more details in [11]).

The primary vertex position was determined from the tracks reconstructed in the ITS and the TPC by using an analytic χ^2 minimization method, applied after approximating each of the tracks by a straight line in the vicinity of their common origin. The event is accepted if the coordinate of the reconstructed vertex measured along the beam direction (z-axis) is within ± 10 cm around the nominal interaction point. The event vertex reconstruction is fully efficient for the event centralities covered.

Primary charged particles are defined as all prompt particles produced in the collision, including decay products, except those from weak decays of strange hadrons. A set of standard cuts based on the number of space points and the quality of the momentum fit in the TPC and ITS is applied to the reconstructed tracks. Track candidates in the TPC are required to have hits in at least 120 (out of a maximum of 159) pad-rows and χ^2 per point of the momentum fit smaller than 4. Such tracks are projected to the ITS and used for further analysis if at least 2 matching hits (out

of a maximum of 6) in the ITS, including at least one in the SPD, are found. In addition, the χ^2 per point of the momentum fit in the ITS must be smaller than 36. In order to improve the purity of primary track reconstruction at high $p_{\rm T}$ we developed a procedure where we compare tracking information from the combined ITS and TPC track reconstruction algorithm to that derived only from the TPC and constrained by the interaction vertex point. We calculated the $\chi^2_{\rm TPC-ITS}$ between these tracks using the following formula

$$\chi_{\text{TPC-ITS}}^2 = (\mathbf{v}_{\text{TPC}} - \mathbf{v}_{\text{TPC-ITS}})^T \cdot (\mathbf{C}_{\text{TPC}} + \mathbf{C}_{\text{TPC-ITS}})^{-1} \cdot (\mathbf{v}_{\text{TPC}} - \mathbf{v}_{\text{TPC-ITS}})$$
(2)

where \mathbf{v}_{TPC} , $\mathbf{v}_{TPC-ITS}$ and \mathbf{C}_{TPC} , $\mathbf{C}_{TPC-ITS}$ represent the measured track parameter vectors $\mathbf{v} = (x, y, z, \theta, \phi, 1/p_T)$ and their covariance matrices, respectively. If the $\chi^2_{TPC-ITS}$ is larger than 36 the track candidate is rejected. At $p_T = 0.15-50$ GeV/c, this procedure removes about 2–7% (1–3%) of the reconstructed tracks in the most central (peripheral) collisions. This procedure in fact removes high- p_T fake tracks, which originate from spurious matches of low p_T particles in the TPC to hits in the ITS, and would result in an incorrect momentum assignment.

Finally, tracks are rejected from the sample if their distance of closest approach to the reconstructed vertex in the longitudinal direction d_z is larger than 2 cm or $d_{xy} > 0.018$ cm + 0.035 cm $\cdot p_T^{-1}$ in the transverse direction with p_T in GeV/c, which corresponds to 7 standard deviations of the resolution in d_{xy} (see [19] for details). The upper limit on the d_z ($d_z < 2$ cm) was set to minimize the contribution of tracks coming from pileup and beam-gas background events. These cuts reject less than 0.5% of the reconstructed tracks independently of p_T and collision centrality.

The efficiency and purity of the primary charged particle selection are estimated using a Monte Carlo simulation with HIJING [20] events and a GEANT3 [21] model of the detector response. We used a HIJING tune which reproduces approximately the measured charged particle density in central collisions [11]. In the most central events, the overall primary charged particle reconstruction efficiency (tracking efficiency and acceptance) in $|\eta|$ < 0.8 is 36% at $p_{\rm T} = 0.15~{\rm GeV}/c$ and increases to 65% for $p_{\rm T} > 0.6~{\rm GeV}/c$. In the most peripheral events the efficiency is larger than that for the central events by about 1-3%. The contribution from secondary particles was estimated using the d_{xy} distributions of data and HI-JING and is consistent with the measured strangeness to charged particle ratio from the reconstruction of K_s^0 , Λ and $\overline{\Lambda}$ invariant mass peaks in Pb-Pb [22]. The total contribution from secondary tracks at $p_T = 0.15 \text{ GeV}/c$ is 13 (7)% for central (peripheral) events and decreases to about 0.6% above $p_T = 4 \text{ GeV}/c$ for both central and peripheral events. From a systematic variation of the $\chi^2_{TPC-ITS}$ cut and comparison of track properties in MC to data we conclude that the number of properly reconstructed tracks rejected as high- p_T fake tracks is around 1-2% (0.5-1%) in the most central (peripheral) collisions. We also conclude that the contribution from the high- p_T fake tracks to the p_T spectra is negligible independently of the collision centrality and p_T .

The transverse momentum of charged particles is reconstructed from the track curvature measured in the magnetic field B=0.5 T using the ITS and TPC detectors. The $p_{\rm T}$ resolution is estimated from the track residuals to the momentum fit and verified by cosmic muon events, and the width of the invariant mass peaks of Λ , $\overline{\Lambda}$ and $K_{\rm S}^0$ reconstructed from their decays to two charged particles. For the selected tracks the relative $p_{\rm T}$ resolution ($\sigma(p_{\rm T})/p_{\rm T}$) amounts to 3.5% at $p_{\rm T}=0.15$ GeV/c, has a minimum of 1% at $p_{\rm T}=1$ GeV/c, and increases linearly to 10% at $p_{\rm T}=50$ GeV/c. It is independent of the centrality of the selected events. From the study of the invariant mass distributions of Λ and $K_{\rm S}^0$ as a

Table 2 Contribution to the systematic uncertainties on the $p_{\rm T}$ spectra (0.15–50 GeV/c) for the most central and peripheral Pb–Pb collisions. Also listed are the systematic uncertainties on the pp reference (0.15–50 GeV/c) [16].

Centrality class	0-5%	70-80%
Centrality selection	0.4%	6.7%
Event selection	3.2%	3.4%
Track selection	4.1-7.3%	3.6-6.0%
Tracking efficiency	5%	5%
$p_{\rm T}$ resolution correction	< 1.8%	< 3%
Material budget	0.9-1.2%	0.5-1.7%
Particle composition	0.6-10%	0.5-7.7%
MC generator	2.5%	1.5%
Secondary particle rejection	< 1%	< 1%
Total for p_T spectra	8.2-13.5%	10.3-13.4%
Total for pp reference	6.3-18.8%	
pp reference normalization	1.9%	

function of $p_{\rm T}$ we estimate that the relative uncertainty on the $p_{\rm T}$ resolution is around 20%. From the mass difference between Λ and $\overline{\Lambda}$ and the ratio of positively to negatively charged tracks, assuming charge symmetry at high $p_{\rm T}$, the upper limit of the systematic uncertainty of the momentum scale is estimated to be $|\Delta(p_{\rm T})/p_{\rm T}| < 0.005$ at $p_{\rm T} = 50$ GeV/c. This has an effect of around 1.5% on the yield of the measured spectra at the highest $p_{\rm T}$. To account for the finite $p_{\rm T}$ resolution, correction factors for the reconstructed $p_{\rm T}$ spectra at $p_{\rm T} > 10$ GeV/c are derived using a folding procedure. The corrections depend on collision centrality due to the change of the spectral shape and reach 4 (8)% at $p_{\rm T} = 50$ GeV/c in the most central (peripheral) collisions.

The systematic uncertainties on the p_T spectra are summarized in Table 2. The systematic uncertainties related to centrality selection were estimated by a comparison of the p_T spectra when the limits of the centrality classes are shifted by $\pm 1\%$ (e.g. for the 70-80% centrality class, 70.7-80.8% and 69.3-79.2%), which is a relative uncertainty on the fraction of the hadronic cross section used in the Glauber fit [11] to determine the centrality classes. We also varied the event and track quality selection criteria and the Monte Carlo assumptions to estimate systematic uncertainties on the p_T spectra. In particular, we studied a variation of the most abundant charged particle species (pions, kaons, protons) by $\pm 30\%$ to match the measured ratios and their uncertainties [22]. The material budget was varied by $\pm 7\%$ [23], and the secondary yield from strangeness decays in the Monte Carlo by $\pm 30\%$ to match the measured d_{xy} distributions. Moreover, we used a different event generator, DPM[ET [24], to calculate MC correction maps. The systematic uncertainties on the $p_{\rm T}$ spectra, related to the high- $p_{\rm T}$ fake track rejection procedure, were estimated by varying the track matching criteria in the range $25 < \chi^2_{TPC\text{-}ITS} < 49,$ and amount to 1–4% (1–2%) in the most central (peripheral) collisions. The total systematic uncertainties on the corrected p_T spectra depend on p_T and event centrality and amount to 8.2–13.5% (10.3–13.4%) in the most central (peripheral) collisions.

A dedicated run of the LHC to collect pp reference data at $\sqrt{s} = 2.76$ TeV took place in March 2011. Data taken in this run were used to measure the charged particle $p_{\rm T}$ spectrum that forms the basis of the pp reference spectrum for $R_{\rm AA}$. Using these data the systematic uncertainties in $R_{\rm AA}$ related to the pp reference could be significantly improved (Table 2) compared to the previous publication [15], allowing for an exploration of high- $p_{\rm T}$ particle suppression in Pb-Pb out to 50 GeV/c. More details about the pp reference determination can be found in [16].




Fig. 1. Charged particle $p_{\rm T}$ distribution measured in Pb–Pb collisions in different centrality intervals. The spectra are scaled for better visibility. The dashed lines show the pp reference [16] spectra scaled by the nuclear overlap function determined for each centrality interval (Table 1) and by the Pb–Pb spectra scaling factors. The systematic and statistical uncertainties for Pb–Pb are added quadratically. The uncertainties on the pp reference are not shown.

3. Results

The fully corrected $p_{\rm T}$ spectra of inclusive charged particles measured in Pb-Pb collisions at $\sqrt{s_{\rm NN}}=2.76$ TeV in nine different centrality intervals, and the scaled pp reference spectra are shown in Fig. 1. At low $p_{\rm T}$, the transverse momentum spectra differ from the pp reference. This is in agreement with the previously observed scaling behavior of the total charged particle production as a function of centrality [11]. A marked depletion of the spectra at high transverse momentum $(p_{\rm T}>5~{\rm GeV}/c)$ develops gradually as centrality increases, indicating strong suppression of high- $p_{\rm T}$ particle production in central collisions.

The nuclear modification factors for nine centrality intervals are shown in Fig. 2. In peripheral collisions (70–80%), only moderate suppression ($R_{AA}=0.6$ –0.7) and a weak p_T dependence is observed. Towards more central collisions, a pronounced minimum at about $p_T=6$ –7 GeV/c develops while for $p_T>7$ GeV/c there is a significant rise of the nuclear modification factor. This rise becomes gradually less steep with increasing p_T . In the most central collisions (0–5%), the yield is most suppressed, $R_{AA}\approx 0.13$ at $p_T=6$ –7 GeV/c, and R_{AA} reaches ≈ 0.4 with no significant p_T dependence for $p_T>30$ GeV/c.

The dependence of R_{AA} on the collision centrality, expressed in terms of N_{part} and the charged particle multiplicity density $(dN_{ch}/d\eta)$, are shown in Fig. 3 for different intervals of p_T . Also shown are results from PHENIX at RHIC in Au–Au collisions at $\sqrt{s_{NN}} = 200$ GeV [9]. The strongest centrality dependence is observed for particles with $5 < p_T < 7$ GeV/c. At higher p_T , the




Fig. 2. Nuclear modification factor R_{AA} of charged particles measured in Pb–Pb collisions in nine centrality intervals. The boxes around data points denote p_T -dependent systematic uncertainties. The systematic uncertainties on the normalization which are related to $\langle T_{AA} \rangle$ and the normalization of the pp data are added in quadrature and shown as boxes at $R_{AA} = 1$.

centrality dependence weakens gradually. In comparison to results from RHIC, the LHC data in the same p_T window show a suppression which is larger by a factor of about 1.2 at all $\langle N_{part} \rangle$ (Fig. 3, top panel). This implies that the shape of the N_{part} dependence at RHIC and the LHC is very similar when the same p_T is compared, indicating a strong relation between collision geometry and energy loss. The overall increase of suppression at the LHC as compared to RHIC may be expected from the larger density and longer lifetime of the fireball. The suppression reaches similar values when results from RHIC are compared to results from the LHC in terms of $dN_{ch}/d\eta$, as shown in Fig. 3 (bottom panel). Larger values of suppression than at RHIC are observed in central collisions at the LHC, where the charged particle multiplicity exceeds that of the most central collisions at RHIC. It should be noted that the suppression at a given centrality results from a subtle interplay between the parton p_T spectrum, the quark-to-gluon ratio, and the medium density, all of which exhibit a significant energy dependence. Further model studies are needed to evaluate their relative

The ALICE measurement of R_{AA} in the most central Pb–Pb collisions (0–5%) is compared to the CMS result [25] in Fig. 4. Both measurements agree within their respective statistical and systematic uncertainties.

In Fig. 4, the measured $R_{\rm AA}$ for 0–5% central collisions is also compared to model calculations. All selected models use RHIC data to calibrate the medium density and were available before the preliminary version of the data reported in this Letter. All model calculations except WHDG [26] use a hydrodynamical description of the medium, but different extrapolation assumptions from RHIC to LHC. A variety of energy loss formalisms is used. An increase of


 R_{AA} due to a decrease of the relative energy loss with increasing p_T is seen for all the models.

The curves labeled WHDG, ASW, and Higher Twist (HT) are based on analytical radiative energy loss formulations that include interference effects. Of those curves, the multiple soft gluon approximation (ASW [27]) and the opacity expansion (WHDG [26]) show a larger suppression than seen in the measurement, while one of the HT curves (Chen [28]) with lower density provides a good description. The other HT (Majumder [29]) curve shows a stronger rise with $p_{\rm T}$ than measured. The elastic energy loss model by Renk (elastic) [30] does not rise steeply enough with $p_{\rm T}$ and overshoots the data at low $p_{\rm T}$. The YaJEM-D model [31], which is based on medium-induced virtuality increases in a parton shower, shows too strong a $p_{\rm T}$ -dependence of $R_{\rm AA}$ due to a formation time cut-off.

A more systematic study of the energy loss formalisms, preferably with the same model(s) for the medium density is needed to rule out or confirm the various effects. Deviations of the nuclear parton distribution functions (PDFs) from a simple scaling of the nucleon PDF with mass number A (e.g. shadowing) are also expected to affect the nuclear modification factor. These effects are predicted to be small for $p_{\rm T} > 10~{\rm GeV}/c$ at the LHC [26] and will be quantified in future p–Pb measurements.

4. Summary

We have reported the measurements of charged particle $p_{\rm T}$ spectra and nuclear modification factors $R_{\rm AA}$ as a function of event centrality in Pb–Pb collisions at $\sqrt{s_{\rm NN}}=2.76$ TeV. The results indicate a strong suppression of charged particle production in Pb–Pb






Fig. 3. Nuclear modification factor R_{AA} of charged particles as a function of $\langle N_{\rm part} \rangle$ (top panel) and $dN_{\rm ch}/d\eta$ (bottom panel) measured by ALICE in Pb–Pb collisions in different $p_{\rm T}$ -intervals, compared to PHENIX results in $5 < p_{\rm T} < 7$ GeV/c [9]. The boxes around the data represent the $p_{\rm T}$ -dependent uncertainties on the Pb–Pb $p_{\rm T}$ spectra. The boxes at $R_{\rm AA}=1$ represent the systematic uncertainties on the pp reference in different $p_{\rm T}$ -intervals ($p_{\rm T}$ -interval increases from left to right, the leftmost is for PHENIX). The systematic uncertainties on the overall normalization for ALICE and PHENIX are not shown.

collisions and a characteristic centrality and p_T dependence of the nuclear modification factors. In central collisions (0–5%) the yield is most strongly suppressed ($R_{AA} \approx 0.13$) at $p_T = 6-7$ GeV/c. Above $p_T = 7$ GeV/c, there is a significant rise in the nuclear modification factor, which reaches $R_{AA} \approx 0.4$ for $p_T > 30$ GeV/c. This result is in agreement with the CMS measurement within statistical and systematic uncertainties. The suppression is weaker in peripheral collisions (70–80%) with $R_{AA} = 0.6-0.7$ and no strong p_T dependence. The observed suppression of high- p_T particles in central Pb-Pb collisions provides evidence for strong parton energy loss and a large medium density at the LHC. We observe that the suppression of charged particles with $5 < p_T < 7$ GeV/c reaches similar values when results from RHIC are compared to results from LHC in terms of the $dN_{ch}/d\eta$. The measured R_{AA} in 0–5%




Fig. 4. Nuclear modification factor R_{AA} of charged particles measured by ALICE in the most central Pb-Pb collisions (0-5%) in comparison to results from CMS [25] and model calculations [26–31]. The boxes around the data denote p_T -dependent systematic uncertainties. For CMS statistical and systematic uncertainties on R_{AA} are added in quadrature. The systematic uncertainties on the normalization which are related to (T_{AA}) and the normalization of the pp data are added in quadrature and shown as boxes at $R_{AA} = 1$ (the right-most is for CMS).

central collisions is compared to model calculations. An increase of $R_{\rm AA}$ due to a decrease of the relative energy loss with increasing $p_{\rm T}$ is seen for all the models. The measurement presented here, together with measurements of particle correlations [32] and measurements using jet reconstruction [33], will help in understanding the mechanism of jet quenching and the properties of the medium produced in heavy-ion collisions.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex.

The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: Calouste Gulbenkian Foundation from Lisbon and Swiss Fonds Kidagan, Armenia; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (FINEP), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP); National Natural Science Foundation of China (NSFC), the Chinese Ministry of Education (CMOE) and the Ministry of Science and Technology of China (MSTC): Ministry of Education and Youth of the Czech Republic; Danish Natural Science Research Council, the Carlsberg Foundation and the Danish National Research Foundation; The European Research Council under the European Community's Seventh Framework Programme; Helsinki Institute of Physics and the Academy of Finland; French CNRS-IN2P3, the 'Region Pays de Loire', 'Region Alsace', 'Region Auvergne' and CEA, France; German BMBF and the Helmholtz Association; General Secretariat for Research and Technology, Ministry of Development, Greece; Hungarian OTKA and National Office for Research and Technology (NKTH); Department of Atomic Energy and Department of Science and Technology of the Government of India; Istituto Nazionale di Fisica Nucleare (INFN) of Italy; MEXT Grantin-Aid for Specially Promoted Research, Japan; Joint Institute for Nuclear Research, Dubna; National Research Foundation of Korea (NRF); CONACYT, DGAPA, México, ALFA-EC and the HELEN Program (High-Energy physics Latin-American-European Network); Stichting voor Fundamenteel Onderzoek der Materie (FOM) and the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; Research Council of Norway (NFR); Polish Ministry of Science and Higher Education; National Authority for Scientific Research — NASR (Autoritatea Natională pentru Cercetare Stiintifică — ANCS): Federal Agency of Science of the Ministry of Education and Science of Russian Federation. International Science and Technology Center, Russian Academy of Sciences, Russian Federal Agency of Atomic Energy, Russian Federal Agency for Science and Innovations and CERN-INTAS; Ministry of Education of Slovakia; Department of Science and Technology, South Africa; CIEMAT, EELA, Ministerio de Educación y Ciencia of Spain, Xunta de Galicia (Consellería de Educación), CEADEN, Cubaenergía, Cuba, and IAEA (International Atomic Energy Agency); Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW); Ukraine Ministry of Education and Science; United Kingdom Science and Technology Facilities Council (STFC); The United States Department of Energy, the United States National Science Foundation, the State of Texas, and the State of Ohio.

Open access

This article is published Open Access at sciencedirect.com. It is distributed under the terms of the Creative Commons Attribution License 3.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are credited.

References

- [1] I. Arsene, et al., BRAHMS Collaboration, Nucl. Phys. A 757 (2005) 1.
- [2] B.B. Back, et al., PHOBOS Collaboration, Nucl. Phys. A 757 (2005) 28.

- [3] J. Adams, et al., STAR Collaboration, Nucl. Phys. A 757 (2005) 102.
- [4] K. Adcox, et al., PHENIX Collaboration, Nucl. Phys. A 757 (2005) 184.
- [5] D. d'Enterria, Phys. Lett. B 596 (2004) 32.
- [6] K. Adcox, et al., PHENIX Collaboration, Phys. Rev. Lett. 88 (2001) 022301.
- [7] C. Adler, et al., STAR Collaboration, Phys. Rev. Lett. 89 (2002) 202301.
- [8] J. Adams, et al., STAR Collaboration, Phys. Rev. Lett. 91 (2003) 172302.
- [9] S.S. Adler, et al., PHENIX Collaboration, Phys. Rev. C 69 (2004) 034910.
- [10] A. Adare, et al., PHENIX Collaboration, Phys. Rev. Lett. 101 (2008) 232301.
 [11] K. Aamodt, et al., ALICE Collaboration, Phys. Rev. Lett. 105 (2010) 252301;
 K. Aamodt, et al., ALICE Collaboration, Phys. Rev. Lett. 106 (2011) 032301;
- B. Abelev, et al., ALICE Collaboration, in press.
 [12] K.J. Eskola, H. Honkanen, C.A. Salgado, U.A. Wiedemann, Nucl. Phys. A 747 (2005) 511
- [13] N. Armesto, et al., arXiv:1106.1106v1 [hep-ph]. Glauber Model
- [14] M. Miller, K. Reygers, S. Sanders, P. Steinberg, Annu. Rev. Nucl. Part. Sci. 57 (2007) 205.
- [15] K. Aamodt, et al., ALICE Collaboration, Phys. Lett. B 696 (2011) 30.
- [16] K. Aamodt, et al., ALICE Collaboration, in press.
- [17] K. Aamodt, et al., ALICE Collaboration, JINST 3 (2008) S08002.
- [18] K. Nakamura, et al., Particle Data Group, J. Phys. G 37 (2010) 075021.
- [19] B. Abelev, et al., ALICE Collaboration, JHEP 1201 (2012) 128.
- [20] X.-N. Wang, M. Gyulassy, Phys. Rev. D 44 (1991) 3501;
 W.-T. Deng, X.-N. Wang, R. Xu, Phys. Rev. C 83 (2011) 014915.
- [21] R. Brun, et al., CERN Program Library Long Write-up, W5013, GEANT Detector Description and Simulation Tool, 1994.
- [22] M. Floris, et al., ALICE Collaboration, J. Phys. G: Nucl. Part. Phys. 38 (2011) 124025.
- [23] K. Koch, et al., ALICE Collaboration, Nucl. Phys. A 855 (2011) 281.
- [24] S. Roesler, R. Engel, J. Ranft, arXiv:hep-ph/0012252.
- [25] S. Chatrchvan, et al., CMS Collaboration, Eur. Phys. I. C 72 (2012) 1945.
- [26] W.A. Horowitz, M. Gyulassy, Nucl. Phys. A 872 (2011) 265.
- [27] C.A. Salgado, U.A. Wiedemann, Phys. Rev. D 68 (2003) 014008.
- [28] X.-F. Chen, T. Hirano, E. Wang, X.-N. Wang, H. Zang, Phys. Rev. C 84 (2011) 034902.
- [29] A. Majumder, B. Muller, Phys. Rev. Lett. 105 (2010) 252002.
- [30] T. Renk, H. Holopainen, R. Paatelainen, K.J. Eskola, Phys. Rev. C 84 (2011) 014906.
- [31] T. Renk, Phys. Rev. C 83 (2011) 024908.
- [32] K. Aamodt, et al., ALICE Collaboration, Phys. Rev. Lett. 108 (2012) 092301.
- [33] B. Abelev, et al., ALICE Collaboration, JHEP 1203 (2012) 053.

ALICE Collaboration

B. Abelev ⁶⁸, J. Adam ³⁴, D. Adamová ⁷³, A.M. Adare ¹²⁰, M.M. Aggarwal ⁷⁷, G. Aglieri Rinella ³⁰, A.G. Agocs ⁶⁰, A. Agostinelli ¹⁹, S. Aguilar Salazar ⁵⁶, Z. Ahammed ¹¹⁶, N. Ahmad ¹⁴, A. Ahmad Masoodi ¹⁴, S.A. Ahn ⁶², S.U. Ahn ³⁷, A. Akindinov ⁴⁶, D. Aleksandrov ⁸⁸, B. Alessandro ⁹⁴, R. Alfaro Molina ⁵⁶, A. Alici ⁹⁷, 10, A. Alkin ², E. Almaráz Aviña ⁵⁶, J. Alme ³², T. Alt ³⁶, V. Altini ²⁸, S. Altinpinar ¹⁵, I. Altsybeev ¹¹⁷, C. Andrei ⁷⁰, A. Andronic ⁸⁵, V. Anguelov ⁸², J. Anielski ⁵⁴, C. Anson ¹⁶, T. Antičić ⁸⁶, F. Antinori ⁹³, P. Antonioli ⁹⁷, L. Aphecetche ¹⁰², H. Appelshäuser ⁵², N. Arbor ⁶⁴, S. Arcelli ¹⁹, A. Arend ⁵², N. Armesto ¹³, R. Arnaldi ⁹⁴, T. Aronsson ¹²⁰, I.C. Arsene ⁸⁵, M. Arslandok ⁵², A. Asryan ¹¹⁷, A. Augustinus ³⁰, R. Averbeck ⁸⁵, T.C. Awes ⁷⁴, J. Äystö ³⁸, M.D. Azmi ¹⁴, ⁷⁹, M. Bach ³⁶, A. Badalà ⁹⁹, Y.W. Baek ^{63,37}, R. Bailhache ⁵², R. Bala ⁹⁴, R. Baldini Ferroli ¹⁰, A. Baldisseri ¹², A. Baldit ⁶³, F. Baltasar Dos Santos Pedrosa ³⁰, J. Bán ⁴⁷, R.C. Baral ⁴⁸, R. Barbera ²⁵, F. Barile ²⁸, G.G. Barnaföldi ⁶⁰, L.S. Barnby ⁹⁰, V. Barret ⁶³, J. Bartke ¹⁰⁴, M. Basile ¹⁹, N. Bastid ⁶³, S. Basu ¹¹⁶, B. Bathen ⁵⁴, G. Batigne ¹⁰², B. Batyunya ⁵⁹, C. Baumann ⁵², I.G. Bearden ⁷¹, H. Beck ⁵², N.K. Behera ⁴⁰, I. Belikov ⁵⁸, F. Bellini ¹⁹, R. Bellwied ¹¹⁰, E. Belmont-Moreno ⁵⁶, G. Bencedi ⁶⁰, S. Beole ²³, I. Berceanu ⁷⁰, A. Brecuci ⁷⁰, S. Beole ¹¹⁰, P. Berenyi ⁶⁰, A.A.E. Bergognon ¹⁰², D. Berzano ⁹⁴, L. Betev ³⁰, A. Bhasin ⁸⁰, A.K. Bhati ⁷⁷, J. Bhom ¹¹⁴, L. Bianchi ²³, N. Bianchi ⁶⁵, C. Bianchin ²⁰, J. Bielčík ³⁴, J. Bielčíková ⁷³, A. Bilandzic ⁷¹, S. Bjelogrlic ⁴⁵, F. Blanco ⁸, F. Blanco ¹¹⁰, D. Blau ⁸⁸, C. Blume ⁵², M. Boccioli ³⁰, N. Bock ¹⁶, S. Böttger ⁵¹, A. Bogdanov ⁶⁹, H. Bøggild ⁷¹, M. Bogolyubsky ⁴³, L. Boldizsár ⁶⁰, M. Bombara ³⁵, J. Book ⁵², H. Borel ¹², P. Braun-Munzin

C. Cavicchioli 30, C. Ceballos Sanchez ⁷, J. Cepila ³⁴, P. Cerello ⁹⁴, B. Chang ^{38,123}, S. Chapeland ³⁰, J.L. Charvet ¹², S. Chatopadhyay ¹⁶, S. Chatopadhyay ⁸⁶, I. Chawla ⁷⁷, M. Cherney ⁷⁶, C. Cheshkov ^{30,109}, B. Cheynis ¹⁰⁹, V. Chibante Barroso ³⁰, D.D. Chinellato ¹⁰⁸, P. Chochula ³⁰, M. Chojnack ³¹, M. Chojnack ³¹, S. Chaudhury ¹¹⁶, P. Christakoglou ⁷², C. H. Christensen ⁷¹, P. Christiansen ²³, T. Chujo ¹¹⁴, S.U. Chung ⁸⁴, C. Cicalo ³⁶, L. Cifarelli ^{30,301}, P. Circhou, P. Chorstantin ³², G. Contin ²¹, J.C. Contreas ³², T. Chung ³⁴, C. Conesa Balbastre ³⁴, Z. Conesa del Valle ³⁰, P. Cortest ³⁷, C. Cortest ³⁸, C. Contrest ³⁸, D. Colella ³⁸, D. Colella ³⁸, D. Colella ³⁸, E. Crace ³⁸, D. Cortes ³⁸, D. Cortes ³⁸, D. Colella ³⁸, D. Colella ³⁸, D. Colella ³⁸, E. Crace ³⁸, D. Dah ³⁸, S. Dah ³⁸, S. Dah ³⁸, D. Dah ³ C. Cavicchioli ³⁰, C. Ceballos Sanchez ⁷, J. Cepila ³⁴, P. Cerello ⁹⁴, B. Chang ^{38,123}, S. Chapeland ³⁰,

I. León Monzón ¹⁰⁶, H. León Vargas ⁵², P. Lévai ⁶⁰, J. Lien ¹⁵, R. Lietava ⁹⁰, S. Lindal ¹⁸, V. Lindenstruth ³⁶, C. Lippmann ^{85,30}, M.A. Lisa ¹⁶, L. Liu ¹⁵, V.R. Loggins ¹¹⁹, V. Loginov ⁶⁹, S. Lohn ³⁰, D. Lohner ⁸², C. Loizides ⁶⁷, K.K. Loo ³⁸, X. Lopez ⁶³, E. López Torres ⁷, G. Løvhøiden ¹⁸, X.-G. Lu ⁸², P. Luettig ⁵², M. Lunardon ²⁰, J. Luo ⁵, G. Luparello ⁴⁵, L. Luquin ¹⁰², C. Luzzi ³⁰, R. Ma ¹²⁰, K. Ma ⁵, D.M. Madagodahettige-Don ¹¹⁰, A. Maevskaya ⁴⁴, M. Mager ^{53,30}, D.P. Mahapatra ⁴⁸, A. Maire ⁸², M. Malaev ⁷⁵, I. Maldonado Cervantes ⁵⁵, L. Malinina ^{59,i}, D. Mal'Kevich ⁴⁶, P. Malzacher ⁸⁵, A. Mamonov ⁸⁷, L. Mangotra ⁸⁰, V. Manko ⁸⁸, F. Manso ⁶³, V. Manzari ⁹⁸, Y. Mao ⁵, M. Marchisone ^{63,23}, J. Mareš ⁴⁹, G.V. Margagliotti ^{21,92}, A. Margotti ⁹⁷, A. Marín ⁸⁵, C.A. Marin Tobon ³⁰, C. Markert ¹⁰⁵, I. Martashvili 112, P. Martinengo 30, M.I. Martínez 1, A. Martínez Davalos 56, G. Martínez García 102, I. Martashvili ¹¹², P. Martinengo ³⁰, M.I. Martínez ¹, A. Martínez Davalos ³⁶, G. Martínez García ¹⁰², Y. Martynov ², A. Mas ¹⁰², S. Masciocchi ⁸⁵, M. Masera ²³, A. Masoni ⁹⁶, L. Massacrier ¹⁰², A. Mastroserio ²⁸, Z.L. Matthews ⁹⁰, A. Matyja ^{104,102}, C. Mayer ¹⁰⁴, J. Mazer ¹¹², M.A. Mazzoni ⁹⁵, F. Meddi ²⁴, A. Menchaca-Rocha ⁵⁶, J. Mercado Pérez ⁸², M. Meres ³³, Y. Miake ¹¹⁴, L. Milano ²³, J. Milosevic ^{18,ii}, A. Mischke ⁴⁵, A.N. Mishra ⁸¹, D. Miśkowiec ^{85,30}, C. Mitu ⁵⁰, J. Mlynarz ¹¹⁹, B. Mohanty ¹¹⁶, L. Molnar ^{60,30}, L. Montaño Zetina ⁹, M. Monteno ⁹⁴, E. Montes ⁸, T. Moon ¹²³, M. Morando ²⁰, D.A. Moreira De Godoy ¹⁰⁷, S. Moretto ²⁰, A. Morsch ³⁰, V. Muccifora ⁶⁵, E. Mudnic ¹⁰³, S. Muhuri ¹¹⁶, M. Mukherjee ¹¹⁶, H. Müller ³⁰, M.G. Munhoz ¹⁰⁷, L. Musa ³⁰, A. Musso ⁹⁴, B.K. Nandi ⁴⁰, R. Nania ⁹⁷, E. Nappi ⁹⁸, C. Nattrass ¹¹², N.P. Naumov ⁸⁷, S. Navin ⁹⁰, T.K. Nayak ¹¹⁶, S. Nazarenko ⁸⁷, C. Nazarov ⁸⁷, A. Nodosekin ⁴⁶, M. Nicascio ²⁸, M. Niculoscu ^{50,30}, R.S. Nickson ⁷¹, T. Niida ¹¹⁴ G. Nazarov ⁸⁷, A. Nedosekin ⁴⁶, M. Nicassio ²⁸, M. Niculescu ^{50,30}, B.S. Nielsen ⁷¹, T. Niida ¹¹⁴, S. Nikolaev ⁸⁸, V. Nikolic ⁸⁶, S. Nikulin ⁸⁸, V. Nikulin ⁷⁵, B.S. Nilsen ⁷⁶, M.S. Nilsson ¹⁸, F. Noferini ^{97,10}, P. Nomokonov⁵⁹, G. Nooren⁴⁵, N. Novitzky³⁸, A. Nyanin⁸⁸, A. Nyatha⁴⁰, C. Nygaard⁷¹, J. Nystrand¹⁵, A. Ochirov¹¹⁷, H. Oeschler^{53,30}, S. Oh¹²⁰, S.K. Oh³⁷, J. Oleniacz¹¹⁸, C. Oppedisano⁹⁴, A. Ortiz Velasquez^{29,55}, G. Ortona²³, A. Oskarsson²⁹, P. Ostrowski¹¹⁸, J. Otwinowski^{85,*}, K. Oyama⁸², K. Ozawa ¹¹³, Y. Pachmayer ⁸², M. Pachr ³⁴, F. Padilla ²³, P. Pagano ²⁶, G. Paić ⁵⁵, F. Painke ³⁶, C. Pajares ¹³, S.K. Pal ¹¹⁶, A. Palaha ⁹⁰, A. Palmeri ⁹⁹, V. Papikyan ¹²¹, G.S. Pappalardo ⁹⁹, W.J. Park ⁸⁵, A. Passfeld ⁵⁴, S.K. Pal ¹¹⁶, A. Palaha ⁹⁰, A. Palmeri ⁹⁹, V. Papikyan ¹²¹, G.S. Pappalardo ⁹⁹, W.J. Park ⁸⁵, A. Passfeld ⁵⁴, B. Pastirčák ⁴⁷, D.I. Patalakha ⁴³, V. Paticchio ⁹⁸, A. Pavlinov ¹¹⁹, T. Pawlak ¹¹⁸, T. Peitzmann ⁴⁵, H. Pereira Da Costa ¹², E. Pereira De Oliveira Filho ¹⁰⁷, D. Peresunko ⁸⁸, C.E. Pérez Lara ⁷², E. Perez Lezama ⁵⁵, D. Perini ³⁰, D. Perrino ²⁸, W. Peryt ¹¹⁸, A. Pesci ⁹⁷, V. Peskov ^{30,55}, Y. Pestov ³, V. Petráček ³⁴, M. Petran ³⁴, M. Petris ⁷⁰, P. Petrov ⁹⁰, M. Petrovici ⁷⁰, C. Petta ²⁵, S. Piano ⁹², A. Piccotti ⁹⁴, M. Pikna ³³, P. Pillot ¹⁰², O. Pinazza ³⁰, L. Pinsky ¹¹⁰, N. Pitz ⁵², D.B. Piyarathna ¹¹⁰, M. Planinic ⁸⁶, M. Płoskoń ⁶⁷, J. Pluta ¹¹⁸, T. Pocheptsov ⁵⁹, S. Pochybova ⁶⁰, P.L.M. Podesta-Lerma ¹⁰⁶, M.G. Poghosyan ^{30,23}, K. Polák ⁴⁹, B. Polichtchouk ⁴³, A. Pop ⁷⁰, S. Porteboeuf-Houssais ⁶³, V. Pospíšil ³⁴, B. Potukuchi ⁸⁰, S.K. Prasad ¹¹⁹, R. Preghenella ^{97,10}, F. Prino ⁹⁴, C.A. Pruneau ¹¹⁹, I. Pshenichnov ⁴⁴, S. Puchagin ⁸⁷, G. Puddu ²², A. Pulvirenti ²⁵, V. Punin ⁸⁷, M. Putiš ³⁵, J. Putschke ^{119,120}, E. Quercigh ³⁰, H. Ovigstad ¹⁸, A. Rachevski ⁹², A. Rademakers ³⁰, T.S. Räihä ³⁸, L. Rak ³⁸, A. Rakotozafindrabe ¹² S. Puchagin ⁸⁷, G. Puddu ²², A. Pulvirenti ²⁵, V. Punin ⁸⁷, M. Putiš ³⁵, J. Putschke ^{119,120}, E. Quercigh ³⁰, H. Qvigstad ¹⁸, A. Rachevski ⁹², A. Rademakers ³⁰, T.S. Räihä ³⁸, J. Rak ³⁸, A. Rakotozafindrabe ¹², L. Ramello ²⁷, A. Ramírez Reyes ⁹, R. Raniwala ⁸¹, S. Raniwala ⁸¹, S.S. Räsänen ³⁸, B.T. Rascanu ⁵², D. Rathee ⁷⁷, K.F. Read ¹¹², J.S. Real ⁶⁴, K. Redlich ^{100,57}, P. Reichelt ⁵², M. Reicher ⁴⁵, R. Renfordt ⁵², A.R. Reolon ⁶⁵, A. Reshetin ⁴⁴, F. Rettig ³⁶, J.-P. Revol ³⁰, K. Reygers ⁸², L. Riccati ⁹⁴, R.A. Ricci ⁶⁶, T. Richert ²⁹, M. Richter ¹⁸, P. Riedler ³⁰, W. Riegler ³⁰, F. Riggi ^{25,99}, B. Rodrigues Fernandes Rabacal ³⁰, M. Rodríguez Cahuantzi ¹, A. Rodriguez Manso ⁷², K. Røed ¹⁵, D. Rohr ³⁶, D. Röhrich ¹⁵, R. Romita ⁸⁵, F. Ronchetti ⁶⁵, P. Rosnet ⁶³, S. Rossegger ³⁰, A. Rossi ^{30,20}, C. Roy ⁵⁸, P. Roy ⁸⁹, A.J. Rubio Montero ⁸, R. Rui ²¹, R. Russo ²³, E. Ryabinkin ⁸⁸, A. Rybicki ¹⁰⁴, S. Sadovsky ⁴³, K. Šafařík ³⁰, R. Sahoo ⁴¹, P.K. Sahu ⁴⁸, J. Saini ¹¹⁶, H. Sakaguchi ³⁹, S. Sakai ⁶⁷, D. Sakata ¹¹⁴, C.A. Salgado ¹³, J. Salzwedel ¹⁶, S. Sambyal ⁸⁰, V. Samsonov ⁷⁵, X. Sanchez Castro ⁵⁸, L. Šándor ⁴⁷, A. Sandoval ⁵⁶, S. Sano ¹¹³, M. Sano ¹¹⁴, R. Santo ⁵⁴, R. Santoro ^{98,30,10}, J. Sarkamo ³⁸, E. Scapparone ⁹⁷, F. Scarlassara ²⁰, R.P. Scharenberg ⁸³, C. Schiaua ⁷⁰, R. Schicker ⁸², C. Schwidt ⁸⁵, H.R. Schmidt ¹¹⁵, S. Schreiner ³⁰, S. Schuchmann ⁵², L. Schukraft ³⁰. R. Schicker 82, C. Schmidt 85, H.R. Schmidt 115, S. Schreiner 30, S. Schuchmann 52, J. Schukraft 30, Y. Schutz ^{30,102}, K. Schwarz ⁸⁵, K. Schweda ^{85,82}, G. Scioli ¹⁹, E. Scomparin ⁹⁴, R. Scott ¹¹², G. Segato ²⁰, I. Selyuzhenkov ⁸⁵, S. Senyukov ⁵⁸, J. Seo ⁸⁴, S. Serci ²², E. Serradilla ^{8,56}, A. Sevcenco ⁵⁰, A. Shabetai ¹⁰², G. Shabratova ⁵⁹, R. Shahoyan ³⁰, S. Sharma ⁸⁰, N. Sharma ⁷⁷, S. Rohni ⁸⁰, K. Shigaki ³⁹, M. Shimomura ¹¹⁴, K. Shtejer ⁷, Y. Sibiriak ⁸⁸, M. Siciliano ²³, E. Sicking ³⁰, S. Siddhanta ⁹⁶, T. Siemiarczuk ¹⁰⁰, D. Silvermyr ⁷⁴, S. Shapana ⁸⁰, N. Sharma ⁸⁰, N. Sharma ⁸⁰, R. Shimomura ¹¹⁴, Shimomura ¹¹⁵, Shimomura ¹¹⁶, Shimomura ¹¹⁷, Shimomura ¹¹⁸, Shimomura ¹ C. Silvestre ⁶⁴, G. Simatovic ^{55,86}, G. Simonetti ³⁰, R. Singaraju ¹¹⁶, R. Singh ⁸⁰, S. Singha ¹¹⁶, V. Singhal ¹¹⁶, B.C. Sinha ¹¹⁶, T. Sinha ⁸⁹, B. Sitar ³³, M. Sitta ²⁷, T.B. Skaali ¹⁸, K. Skjerdal ¹⁵, R. Smakal ³⁴, N. Smirnov ¹²⁰, R.J.M. Snellings ⁴⁵, C. Søgaard ⁷¹, R. Soltz ⁶⁸, H. Son ¹⁷, J. Song ⁸⁴, M. Song ¹²³, C. Soos ³⁰, F. Soramel ²⁰,

```
60
                                                                                                                                                                      ALICE Collaboration / Physics Letters B 720 (2013) 52-62
  I. Sputowska <sup>104</sup>, M. Spyropoulou-Stassinaki <sup>78</sup>, B.K. Srivastava <sup>83</sup>, J. Stachel <sup>82</sup>, I. Stan <sup>50</sup>, I. Stan <sup>50</sup>, G. Stefanek <sup>100</sup>, M. Steinpreis <sup>16</sup>, E. Stenlund <sup>29</sup>, G. Steyn <sup>79</sup>, J.H. Stiller <sup>82</sup>, D. Stocco <sup>102</sup>, M. Stolpovskiy <sup>43</sup>,
G. Stefanek <sup>100</sup>, M. Steinpreis <sup>16</sup>, E. Stenlund <sup>29</sup>, G. Steyn <sup>79</sup>, J.H. Stiller <sup>82</sup>, D. Stocco <sup>102</sup>, M. Stolpovskiy <sup>43</sup>, K. Strabykin <sup>87</sup>, P. Strmen <sup>33</sup>, A.A.P. Suaide <sup>107</sup>, M.A. Subieta Vásquez <sup>23</sup>, T. Sugitate <sup>39</sup>, C. Suire <sup>42</sup>, M. Sukhorukov <sup>87</sup>, R. Sultanov <sup>46</sup>, M. Šumbera <sup>73</sup>, T. Susa <sup>86</sup>, T.J.M. Symons <sup>67</sup>, A. Szanto de Toledo <sup>107</sup>, I. Szarka <sup>33</sup>, A. Szczepankiewicz <sup>104,30</sup>, A. Szostak <sup>15</sup>, M. Szymański <sup>118</sup>, J. Takahashi <sup>108</sup>, J.D. Tapia Takaki <sup>42</sup>, A. Tauro <sup>30</sup>, G. Tejeda Muñoz <sup>1</sup>, A. Telesca <sup>30</sup>, C. Terrevoli <sup>28</sup>, J. Thäder <sup>85</sup>, D. Thomas <sup>45</sup>, R. Tieulent <sup>109</sup>, A.R. Timmins <sup>110</sup>, D. Tlusty <sup>34</sup>, A. Toia <sup>36,20,93</sup>, H. Torii <sup>113</sup>, L. Toscano <sup>94</sup>, V. Trubnikov <sup>2</sup>, D. Truesdale <sup>16</sup>, W.H. Trzaska <sup>38</sup>, T. Tsuji <sup>113</sup>, A. Tumkin <sup>87</sup>, R. Turrisi <sup>93</sup>, T.S. Tveter <sup>18</sup>, J. Ulery <sup>52</sup>, K. Ullaland <sup>15</sup>, J. Ulrich <sup>61,51</sup>, A. Uras <sup>109</sup>, J. Urbán <sup>35</sup>, G.M. Urciuoli <sup>95</sup>, G.L. Usai <sup>22</sup>, M. Vajzer <sup>34,73</sup>, M. Vala <sup>59,47</sup>, L. Valencia Palomo <sup>42</sup>, S. Vallero <sup>82</sup>, P. Vande Vyvre <sup>30</sup>, M. van Leeuwen <sup>45</sup>, L. Vannucci <sup>66</sup>, A. Vargas <sup>1</sup>, R. Varma <sup>40</sup>, M. Vasileiou <sup>78</sup>, A. Vasiliev <sup>88</sup>, V. Vechernin <sup>117</sup>, M. Veldhoen <sup>45</sup>, M. Venaruzzo <sup>21</sup>, E. Vercellin <sup>23</sup>, S. Vergara <sup>1</sup>, R. Vernet <sup>6</sup>, M. Verweij <sup>45</sup>, L. Vickovic <sup>103</sup>, G. Viesti <sup>20</sup>, O. Vikhlyantsev <sup>87</sup>, Z. Vilakazi <sup>79</sup>, O. Villalobos Baillie <sup>90</sup>, Y. Vinogradov <sup>87</sup>, L. Vinogradov <sup>117</sup>, A. Vinogradov <sup>88</sup>, T. Virgili <sup>26</sup>, Y.P. Viyogi <sup>116</sup>, A. Vodopyanov <sup>59</sup>, K. Voloshin <sup>46</sup>, S. Voloshin <sup>119</sup>, G. Volpe <sup>28,30</sup>, B. von Haller <sup>30</sup>, D. Vranic <sup>85</sup>, G. Øvrebekk <sup>15</sup>, J. Vrláková <sup>35</sup>, B. Vulpescu <sup>63</sup>, A. Vyushin <sup>87</sup>, V. Wagner <sup>34</sup>, B. Wagner <sup>15</sup>, R. Wan <sup>5</sup>, D. Wang <sup>5</sup>, M. Wang <sup>5</sup>, Y. Wang <sup>5</sup>, Y. Wang <sup>5</sup>, Y. Wang <sup>82</sup>, K. Watanabe <sup>114</sup>, M. Weber <sup>110</sup>, J.P. Wessels <sup>30,54</sup>, U. Westerhoff <sup>54</sup>, J. Wiechula <sup>115</sup>, J. Wikne <sup>18</sup>, M. Wilde <sup>54</sup>, A. Wilk <sup>54</sup>, G. Wilk <sup>100</sup>, M.C.S. Williams <sup>97</sup>, B. Windelband <sup>82</sup>, L. Xaplanteris Karampatsos <sup>105</sup>, C.G. Yaldo <sup>119</sup>, Y. Yamaguchi <sup>113</sup>, S. Yang <sup>15</sup>, H.
  N. Zaviyalov <sup>87</sup>, H. Zbroszczyk <sup>118</sup>, P. Zelnicek <sup>51</sup>, I.S. Zgura <sup>50</sup>, M. Zhalov <sup>75</sup>, X. Zhang <sup>63,5</sup>, H. Zhang <sup>5</sup>, F. Zhou <sup>5</sup>, Y. Zhou <sup>45</sup>, D. Zhou <sup>5</sup>, J. Zhu <sup>5</sup>, X. Zhu <sup>5</sup>, J. Zhu <sup>5</sup>, A. Zichichi <sup>19,10</sup>, A. Zimmermann <sup>82</sup>, G. Zinovjev <sup>2</sup>, Y. Zoccarato <sup>109</sup>, M. Zynovyev <sup>2</sup>, M. Zyzak <sup>52</sup>
    <sup>1</sup> Benemérita Universidad Autónoma de Puebla, Puebla, Mexico
    <sup>2</sup> Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine
    <sup>3</sup> Budker Institute for Nuclear Physics, Novosibirsk, Russia
    <sup>4</sup> California Polytechnic State University, San Luis Obispo, CA, United States
    <sup>5</sup> Central China Normal University, Wuhan, China
    <sup>6</sup> Centre de Calcul de l'IN2P3, Villeurbanne, France
    <sup>7</sup> Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba
   <sup>8</sup> Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain
    <sup>9</sup> Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico
    10 Centro Fermi – Centro Studi e Ricerche e Museo Storico della Fisica "Enrico Fermi", Rome, Italy
    <sup>11</sup> Chicago State University, Chicago, United States
    12 Commissariat à l'Energie Atomique, IRFU, Saclay, France
    13 Departamento de Física de Partículas and IGFAE, Universidad de Santiago de Compostela, Santiago de Compostela, Spain
    <sup>14</sup> Department of Physics Aligarh Muslim University, Aligarh, India
    <sup>15</sup> Department of Physics and Technology, University of Bergen, Bergen, Norway
    <sup>16</sup> Department of Physics, Ohio State University, Columbus, OH, United States
    <sup>17</sup> Department of Physics, Sejong University, Seoul, South Korea
    <sup>18</sup> Department of Physics, University of Oslo, Oslo, Norway
    <sup>19</sup> Dipartimento di Fisica dell'Università and Sezione INFN, Bologna, Italy
    <sup>20</sup> Dipartimento di Fisica dell'Università and Sezione INFN, Padova, Italy
    <sup>21</sup> Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy
    <sup>22</sup> Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy
    <sup>23</sup> Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy
    <sup>24</sup> Dipartimento di Fisica dell'Università 'La Sapienza' and Sezione INFN, Rome, Italy
```

²⁵ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy
²⁶ Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy

³³ Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava, Slovakia

⁴² Institut de Physique Nucléaire d'Orsay (IPNO), Université Paris-Sud, CNRS-IN2P3, Orsay, France

38 Helsinki Institute of Physics (HIP) and University of Jyväskylä, Jyväskylä, Finland

³⁴ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic

³⁶ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany

²⁸ Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy
²⁹ Division of Experimental High Energy Physics, University of Lund, Lund, Sweden
³⁰ European Organization for Nuclear Research (CERN), Geneva, Switzerland

³² Faculty of Engineering, Bergen University College, Bergen, Norway

³⁷ Gangneung-Wonju National University, Gangneung, South Korea

³⁵ Faculty of Science, P.J. Šafárik University, Košice, Slovakia

⁴⁰ Indian Institute of Technology Bombay (IIT), Mumbai, India ⁴¹ Indian Institute of Technology Indore (IIT), Indore, India

⁴³ Institute for High Energy Physics, Protvino, Russia

³¹ Fachhochschule Köln, Köln, Germany

³⁹ Hiroshima University, Hiroshima, Japan

²⁷ Dipartimento di Scienze e Innovazione Tecnologica dell'Università del Piemonte Orientale and Gruppo Collegato INFN, Alessandria, Italy

- ⁴⁴ Institute for Nuclear Research, Academy of Sciences, Moscow, Russia
- ⁴⁵ Nikhef, National Institute for Subatomic Physics and Institute for Subatomic Physics of Utrecht University, Utrecht, Netherlands
- ⁴⁶ Institute for Theoretical and Experimental Physics, Moscow, Russia
- ⁴⁷ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia
- ⁴⁸ Institute of Physics, Bhubaneswar, India
- ⁴⁹ Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic
- ⁵⁰ Institute of Space Sciences (ISS), Bucharest, Romania
- ⁵¹ Institut für Informatik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- ⁵² Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- ⁵³ Institut für Kernphysik, Technische Universität Darmstadt, Darmstadt, Germany
- ⁵⁴ Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, Münster, Germany
- ⁵⁵ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁵⁶ Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁵⁷ Institut of Theoretical Physics, University of Wroclaw, Poland
- ⁵⁸ Institut Pluridisciplinaire Hubert Curien (IPHC), Université de Strasbourg, CNRS-IN2P3, Strasbourg, France
- ⁵⁹ Joint Institute for Nuclear Research (JINR), Dubna, Russia
- 60 KFKI Research Institute for Particle and Nuclear Physics, Hungarian Academy of Sciences, Budapest, Hungary
- ⁶¹ Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- ⁶² Korea Institute of Science and Technology Information, Daejeon, South Korea
- 63 Laboratoire de Physique Corpusculaire (LPC), Clermont Université, Université Blaise Pascal, CNRS-IN2P3, Clermont-Ferrand, France
- 64 Laboratoire de Physique Subatomique et de Cosmologie (LPSC), Université Joseph Fourier, CNRS-IN2P3, Institut Polytechnique de Grenoble, Grenoble, France
- ⁶⁵ Laboratori Nazionali di Frascati, INFN, Frascati, Italy
- 66 Laboratori Nazionali di Legnaro, INFN, Legnaro, Italy
- ⁶⁷ Lawrence Berkeley National Laboratory, Berkeley, CA, United States
- ⁶⁸ Lawrence Livermore National Laboratory, Livermore, CA, United States
- ⁶⁹ Moscow Engineering Physics Institute, Moscow, Russia
- ⁷⁰ National Institute for Physics and Nuclear Engineering, Bucharest, Romania
- ⁷¹ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- ⁷² Nikhef, National Institute for Subatomic Physics, Amsterdam, Netherlands
- ⁷³ Nuclear Physics Institute, Academy of Sciences of the Czech Republic, Řež u Prahy, Czech Republic
- ⁷⁴ Oak Ridge National Laboratory, Oak Ridge, TN, United States
- ⁷⁵ Petersburg Nuclear Physics Institute, Gatchina, Russia
- ⁷⁶ Physics Department, Creighton University, Omaha, NE, United States
- ⁷⁷ Physics Department, Panjab University, Chandigarh, India
- ⁷⁸ Physics Department, University of Athens, Athens, Greece
- ⁷⁹ Physics Department, University of Cape Town, iThemba LABS, Cape Town, South Africa
- ⁸⁰ Physics Department, University of Jammu, Jammu, India
- ⁸¹ Physics Department, University of Rajasthan, Jaipur, India
- ⁸² Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- 83 Purdue University, West Lafayette, IN, United States
- ⁸⁴ Pusan National University, Pusan, South Korea
- 85 Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany
- ⁸⁶ Rudjer Bošković Institute, Zagreb, Croatia
- ⁸⁷ Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
- ⁸⁸ Russian Research Centre Kurchatov Institute, Moscow, Russia
- 89 Saha Institute of Nuclear Physics, Kolkata, India
- 90 School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
- 91 Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
- ⁹² Sezione INFN, Trieste, Italy
- 93 Sezione INFN, Padova, Italy
- ⁹⁴ Sezione INFN, Turin, Italy
- ⁹⁵ Sezione INFN, Rome, Italy
- ⁹⁶ Sezione INFN, Cagliari, Italy
- 97 Sezione INFN, Bologna, Italy
- 98 Sezione INFN, Bari, Italy
- ⁹⁹ Sezione INFN, Catania, Italy
- 100 Soltan Institute for Nuclear Studies, Warsaw, Poland
- ¹⁰¹ Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
- ¹⁰² SUBATECH, Ecole des Mines de Nantes, Université de Nantes, CNRS-IN2P3, Nantes, France
- ¹⁰³ Technical University of Split FESB, Split, Croatia
- ¹⁰⁴ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
- 105 The University of Texas at Austin, Physics Department, Austin, TX, United States
- 106 Universidad Autónoma de Sinaloa, Culiacán, Mexico
- ¹⁰⁷ Universidade de São Paulo (USP), São Paulo, Brazil
- ¹⁰⁸ Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
- 109 Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, France
- 110 University of Houston, Houston, TX, United States
- ¹¹¹ University of Technology and Austrian Academy of Sciences, Vienna, Austria
- 112 University of Tennessee, Knoxville, TN, United States
- 113 University of Tokyo, Tokyo, Japan
- 114 University of Tsukuba, Tsukuba, Japan
- ¹¹⁵ Eberhard Karls Universität Tübingen, Tübingen, Germany
- ¹¹⁶ Variable Energy Cyclotron Centre, Kolkata, India
- 117 V. Fock Institute for Physics, St. Petersburg State University, St. Petersburg, Russia
- 118 Warsaw University of Technology, Warsaw, Poland
- 119 Wayne State University, Detroit, MI, United States
- 120 Yale University, New Haven, CT, United States
- ¹²¹ Yerevan Physics Institute, Yerevan, Armenia
- ¹²² Yildiz Technical University, Istanbul, Turkey

- 123 Yonsei University, Seoul, South Korea
 124 Zentrum für Technologietransfer und Telekommunikation (ZTT), Fachhochschule Worms, Worms, Germany
- * Corresponding author. *E-mail address:* j.otwinowski@gsi.de (J. Otwinowski).
- i Also at: M.V. Lomonosov Moscow State University, D.V. Skobeltsyn Institute of Nuclear Physics, Moscow, Russia.
- ii Also at: University of Belgrade, Faculty of Physics and "Vinča" Institute of Nuclear Sciences, Belgrade, Serbia.