

Fractional Brownian Motion and its Application in financial mathematics

Ke Zhu

Supervisor: Prof. Dr. rer. nat. M. Keller-Ressel

Institute of Mathematical Stochastics
TU Dresden

July ??, 2015

Introduction of fBm

Introduction of fBm

Definition and Properties of fBm

Introduction of fBm

Definition and Properties of fBm

Applications

Introduction of fBm

Definition and Properties of fBm

Applications

OU-Process

Introduction of fBm

Definition and Properties of fBm

Applications

OU-Process

Fractional Black-Scholes Model

Introduction of fBm

Definition and Properties of fBm

Applications

OU-Process

Fractional Black-Scholes Model

Fractional Volatility Model

Aims

Centered Gaussian process $(U_H(t))$

$\text{Cov}[U_H(t), U_H(s)] = \frac{1}{2}(|t|^{2H} + |s|^{2H} - |t - s|^{2H})$ with Hurst exponent $H \in (0, 1)$.

Mandelbrot and Van Ness

Definition

Let $(U_H(t))_{t \in \mathbb{R}}$ be a \mathbb{R} -valued stochastic process and H be a real number such that $0 < H < 1$. $(U_H(t))$ is said to be *fractional Brownian motion* if

$$\begin{aligned} & U_H(t) - U_H(s) \\ &= \frac{1}{\Gamma(H + \frac{1}{2})} \left(\int_{\mathbb{R}} \mathbb{1}_{\{t > u\}} \cdot (t - u)^{H - \frac{1}{2}} - \mathbb{1}_{\{s > u\}} (-u)^{H - \frac{1}{2}} dB_u \right) (1) \end{aligned}$$

for $t \geq s, t, s \in \mathbb{R}$, where (B_u) is defined as two-sides Brownian motion and the integral is defined in the sense of stable integral.

Theorem

$U_H(0) = 0$, then

$$U_H(t) = \frac{1}{\Gamma(H + \frac{1}{2})} \left(\int_{\mathbb{R}} \underbrace{\mathbb{1}_{\{t>u\}} \cdot (t-u)^{H-\frac{1}{2}} - \mathbb{1}_{\{u<0\}}(-u)^{H-\frac{1}{2}}}_{f_t(u)} dB_u \right) \quad (2)$$

$$J(f_t) : f_t \rightarrow \int_{\mathbb{R}} f_t dB_u$$

If the integrand f is quadratic integrable then (1) is well-defined in the sense of stable integral.

Theorem

$$U_H(t) \sim \mathcal{N}(0, \frac{1}{\Gamma(H+\frac{1}{2})^2} (\int_{\mathbb{R}} |f_t(u)|^2 du))$$

Proposition

The stable integral is linear:

$$J(f_t + f_s) \stackrel{a.s.}{=} J(f_t) + J(f_s)$$

Theorem

$$U_H(t) - U_H(s) \sim \mathcal{N}(0, \frac{1}{\Gamma(H+\frac{1}{2})^2} (\int_{\mathbb{R}} |f_t(u) - f_s(u)|^2 du))$$

Theorem

The variance of $U_H(t)$ is $\frac{1}{(\Gamma(H+\frac{1}{2}))^2} t^{2H} \mathbb{E} U_H^2(1)$ for any $t \in \mathbb{R}$.

Divide with $\frac{1}{(\Gamma(H+\frac{1}{2}))^2} \mathbb{E} U_H^2(1)$ then,

$$\text{Var}[U_H(t)] = t^{2H}.$$

What is a fBm ?

Theorem

Let $(U_H(t))_t$ be a fBm. The covariance of $U_H(t)$ and $U_H(s)$ is $\frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H})$ for $t, s \in \mathbb{R}$.

Proof.

$$\begin{aligned}\text{Cov}[U_H(t), U_H(s)] &= \mathbb{E}[U_H(t)U_H(s)] \\ &= \frac{1}{2}(\mathbb{E}[U_H(t)^2] + \mathbb{E}[U_H(s)^2] \\ &\quad - \mathbb{E}[(U_H(t) - U_H(s))^2]) \\ &= \frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H})\end{aligned}\tag{3}$$



Theorem

$(U_H(t))_t$ is Gaussian process.

Theorem

Let $(U_H(t))_t$ be a fBm, then $(U_H(t))_t$ has stationary and H -self similar increments .

Theorem

$$(U_H(t_1), \dots, U_H(t_n)) \sim (U_H(t_1 + \tau), \dots, U_H(t_n + \tau))$$

Theorem

$$(U_H(ct_1), \dots, U_H(ct_k)) \sim (c^H U_H(t_1), \dots, c^H U_H(t_k))$$

Definition

A stationary stochastic process $(X_t)_t$ is said to have *long memory* if its autocovariance $\varsigma_X(\tau)$ tends to 0 so slowly such that $\sum_{\tau=0}^{\infty} \varsigma_X(\tau)$ diverges.

$$S_H(k) = U_H(k+1) - U_H(k) \text{ for } k \in \mathbb{R}.$$

Theorem

The fractional Brownian noise $S_H(k)$ with $H \in (\frac{1}{2}, 1)$ has long memory.

fBm is not semimartingal for $H \neq \frac{1}{2}$.

$$dX_t = -aX_t dt + \gamma dU_H(t),$$

where $(X_t)_{t \geq 0}$ is a stochastic process, $a, \gamma \in \mathbb{R}_+$ and $(U_H(t))_{t \geq 0}$ fBm with Hurst exponent H . In fact, given an initial condition $X_0(\omega) = b(\omega)$, then in the theory of SDE, (4) is understood as

$$X_t(\omega) = b(\omega) - a \int_0^t X_u(\omega) du + \gamma U_H(t)(\omega)$$

for $t \geq 0$.

In order to have a stationary solution, we assume that the initial value is centered Gaussian that

$$\hat{X}_{H,t} := \hat{X}_t^{\gamma \int_{-\infty}^0 e^{au} dU_H(u), H} := e^{-at} \left(\gamma \int_{-\infty}^t e^{au} dU_H(u) \right).$$

Theorem

$(\hat{X}_{H,t})_{t \geq 0}$ has long memory for $H \in (\frac{1}{2}, 1)$.

Fractional Black-Scholes model

$$A_t = \exp(rt)$$

$$S_t = \exp(rt + \mu(t) + \sigma U_H(t)), t \in [0, T],$$

where $r \in \mathbb{R}, \sigma \in \mathbb{R}_+, \sup_{t \in [0, T]} \mu(t) < \infty$.

Theorem

Let $(S_t)_{t \in [0, T]}$ be a stochastic process such that

$$\tilde{S}_t = \exp(\mu(t) + \sigma U_H(t)), \quad (4)$$

where μ, σ are as in (4), $U_H(t)$ is a fBm. If there exist

$$\xi_t^1 = f_0 \mathbb{1}_{\{0\}}(t) + \sum_{k=1}^{n-1} f_k \mathbb{1}_{(\tau_k, \tau_{k+1}]}(t)$$

where $t \in [0, T]$, f_k is family of $\mathcal{F}_k^{U_H}$ -measurable function for $k \in \{1, \dots, n-1\}$. $0 = \tau_1 < \dots < \tau_n = T$ are stopping times with respect to $\mathcal{F}_{\tau_k}^{U_H}$ respectively, with $\tau_{k+1} - \tau_k \geq m$ for some $m > 0$. If there exists a $k \in \{0, \dots, n-1\}$ such that $\mathcal{P}[f_k \neq 0] > 0$, then

$$\mathcal{P}[(\xi^1 \cdot \tilde{S})_T < 0] > 0,$$

where $(\xi^1 \cdot \tilde{S})_T := \sum_{k=1}^n \xi_{\tau_k}^1 (\tilde{S}_k - \tilde{S}_{k-1})$.

The market (4) is arbitrage-free, if there exists a minimal amount of time between two successive transactions,.

$$dS_t = r_t S_t dt + \sigma_t S_t dB_t, \quad (5)$$

$$\sigma_t = \exp\{X_t\}$$

$$dX_t = -aX_t dt + \gamma dU_H(t), \quad (6)$$

where $a, \gamma \in \mathbb{R}_+$. In the proceeding section, we have a stationary solution

$$\hat{X}_{H,t} = e^{-at} \gamma \int_{-\infty}^t e^{au} dU_H(u) \quad (7)$$

Theorem

$(\hat{\sigma}_{H,t})$ has long memory for $H \in (\frac{1}{2}, 1)$.

Smoothness of σ_t .

$$s(\tau, \sigma) = \frac{1}{N} \sum_{k=1}^N |\log(\sigma_{k\tau}) - \log(\sigma_{(k-1)\tau})|^2,$$

Theorem

$$\text{Var}[\hat{X}_{H,t,a}] - \text{Cov}[\hat{X}_{H,t,a}, \hat{X}_{H,t+\tau,a}] \rightarrow \frac{1}{2} \gamma^2 \tau^{2H} \quad (8)$$

as a goes to zero, for $t > 0, \tau > 0$.

Definition

A *mixed fractional Brownian motion* is defined as follows

$$M_{\alpha,\beta,H_1,H_2}(t) = \alpha U_{H_1}(t) + \beta U_{H_2}(t) \quad (9)$$

for $t \in \mathbb{R}$, where α, β are real numbers and U_{H_1}, U_{H_2} are two independent fBm's with Hurst exponents $H_1 \in (0, \frac{1}{2}), H_2 \in (\frac{1}{2}, 1)$ respectively.

$$X_{\alpha,\beta,H_1,H_2}(t) = X_{\alpha,\beta,H_1,H_2}(0) - a \int_0^t X_u du + \gamma M_{\alpha,\beta,H_1,H_2}(t).$$

$$\hat{X}_{\alpha,\beta,H_1,H_2}(t) = \alpha\gamma e^{-at} \int_{-\infty}^t e^{au} dU_{H_1} + \beta\gamma e^{-at} \int_{-\infty}^t e^{au} dU_{H_2}.$$

Proposition

\hat{X}_t satisfies following properties:

- (i) $(\hat{X}_t)_{t \geq 0}$ is a centered Gaussian stationary process.
- (ii) \hat{X}_t has long memory.

Proposition

Let M_{α,β,H_1,H_2} be a weighted fractional brownian motion with respect to U_{H_1} and U_{H_2} . $T, \tau > 0$, a, γ are defined by (??). J_{H_1}, J_{H_2} are defined by (10). $\phi = H_1 - \frac{1}{2}, \psi = H_2 - \frac{1}{2}$. Then, for $t \in [0, T]$,

- (i) $E[\sup_{t \in [0, T]} |\hat{X}_{\alpha,\beta,H_1,H_2}(t) - U_{H_1}(t)|] \rightarrow 0$ as $a \rightarrow 0, \alpha \rightarrow 1$.
- (ii) $E[|\hat{X}_{\alpha,\beta,H_1,H_2}(t + \tau) - \hat{X}_{\alpha,\beta,H_1,H_2}(t)|^2] \rightarrow \gamma^2 \tau^{2H}$ as $a \rightarrow 0, \alpha \rightarrow 1$.