

Fractional Brownian Motion and its Application in financial mathematics

Ke Zhu

Supervisor: Prof. Dr. rer. nat. M. Keller-Ressel

July 22, 2015

Contents

Introduction of fBm

Contents

Introduction of fBm

Definition and Properties of fBm

Contents

Introduction of fBm

Definition and Properties of fBm

Applications

Contents

Introduction of fBm

Definition and Properties of fBm

Applications

fOU-Process

Contents

Introduction of fBm

Definition and Properties of fBm

Applications

fOU-Process

Fractional Black-Scholes Model

Contents

Introduction of fBm

Definition and Properties of fBm

Applications

fOU-Process

Fractional Black-Scholes Model

Fractional Stochastic Volatility Model

Introduction of fBm

What is a fBm ?

Classical Definition

A centered Gaussian process $(U_H(t))_{t \in \mathbb{R}}$ with real number $0 < H < 1$ such that $E[U_H(t)U_H(s)] = \frac{1}{2}(s^{2H} + t^{2H} - |t - s|^{2H})$

Motivation

Problem: Is there a Representation of fBM ?

Motivation

Problem: Is there a Representation of fBM ?

Yes, Integral Representation.

Integral Representation

Mandelbrot and Van Ness[1]

Definition

$(U_H(t))$ is said to be *fractional Brownian motion* if

$$U_H(t) - U_H(s) = \frac{1}{\Gamma(H + \frac{1}{2})} \left(\int_{\mathbb{R}} \mathbb{1}_{\{t > u\}} \cdot (t - u)^{H - \frac{1}{2}} - \mathbb{1}_{\{s > u\}} (-u)^{H - \frac{1}{2}} dB_u \right)$$

for $t \geq s, t, s \in \mathbb{R}$, where (B_u) is defined as two-sides Brownian motion and the integral is defined in the sense of stable integral.

Theorem

$U_H(0) = 0$, then

$$U_H(t) = \frac{1}{\Gamma(H + \frac{1}{2})} \left(\int_{\mathbb{R}} \underbrace{\mathbb{1}_{\{t > u\}} \cdot (t - u)^{H - \frac{1}{2}} - \mathbb{1}_{\{u < 0\}} (-u)^{H - \frac{1}{2}}}_{f_t(u)} dB_u \right).$$

$$J(f_t) : f_t \rightarrow \int_{\mathbb{R}} f_t dB_u$$

If the integrand f_t is quadratic integrable then integral is well-defined in the sense of stable integral.

Introduction of fBm

Proposition

The stable integral is linear[2] :

$$J(af_t + bf_s) \stackrel{a.s.}{=} aJ(f_t) + bJ(f_s)$$

Theorem

$$U_H(t) \sim \mathcal{N}(0, \frac{1}{\Gamma(H+\frac{1}{2})^2} (\int_{\mathbb{R}} |f_t(u)|^2 du))$$

Introduction of fBm

Proposition

The stable integral is linear[2] :

$$J(af_t + bf_s) \stackrel{a.s.}{=} aJ(f_t) + bJ(f_s)$$

Theorem

$$U_H(t) \sim \mathcal{N}(0, \frac{1}{\Gamma(H+\frac{1}{2})^2} (\int_{\mathbb{R}} |f_t(u)|^2 du))$$

Theorem

$$U_H(t) - U_H(s) \sim \mathcal{N}(0, \frac{1}{\Gamma(H+\frac{1}{2})^2} (\int_{\mathbb{R}} |f_t(u) - f_s(u)|^2 du))$$

Standardization

Corollary

The variance of $U_H(t)$ is $\frac{1}{(\Gamma(H+\frac{1}{2}))^2} \mathbb{E}U_H^2(1)t^{2H}$ for any $t \in \mathbb{R}$.

Divide with $\frac{1}{(\Gamma(H+\frac{1}{2}))^2} \mathbb{E}U_H^2(1)$ then,

$$\text{Var}[U_H(t)] = t^{2H}.$$

fBm properties

Theorem

Let $(U_H(t))_t$ be a fBm. The covariance of $U_H(t)$ and $U_H(s)$ is $\frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H})$ for $t, s \in \mathbb{R}$.

fBm properties

Theorem

Let $(U_H(t))_t$ be a fBm. The covariance of $U_H(t)$ and $U_H(s)$ is $\frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H})$ for $t, s \in \mathbb{R}$.

Proof.

$$\begin{aligned}\text{Cov}[U_H(t), U_H(s)] &= \mathbb{E}[U_H(t)U_H(s)] \\ &= \frac{1}{2}(\mathbb{E}[U_H(t)^2] + \mathbb{E}[U_H(s)^2] \\ &\quad - \mathbb{E}[(U_H(t) - U_H(s))^2]) \\ &= \frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H})\end{aligned}$$

fBm properties

Theorem

$(U_H(t))_t$ is Gaussian process.

Theorem

$(U_H(t))_t$ has stationary and H -self similar increments .

fBm properties

Theorem

$$\begin{aligned} & ((U_H(t_1 + \tau) - U_H(t_0 + \tau)), \dots, (U_H(t_n + \tau) - U_H(t_{n-1} + \tau))) \\ & \sim (U_H(t_1 - t_0), \dots, U_H(t_n - t_{n-1})) \end{aligned}$$

Theorem

$$\begin{aligned} & (U_H(c(t_1 - t_0)), \dots, U_H(c(t_n - t_{n-1}))) \\ & \sim (c^H U_H(t_1 - t_0), \dots, c^H U_H(t_n - t_{n-1})) \end{aligned}$$

for $c > 0$.

Hurst exponent

H plays role.

Hurst exponent

Case $H = \frac{1}{2}$

fBM is BM.

Hurst exponent

Definition

A stationary stochastic process $(X_t)_t$ is said to have *long memory* if its autocovariance $\varsigma_X(\tau)$ tends to 0 so slowly such that $\sum_{\tau=0}^{\infty} \varsigma_X(\tau)$ diverges.

Hurst exponent

Case $H > \frac{1}{2}$

$$S_H(k) = U_H(k+1) - U_H(k) \text{ for } k \in \mathbb{R}$$

Theorem

The fractional Brownian noise $S_H(k)$ with $H \in (\frac{1}{2}, 1)$ has long memory.

Hurst exponent

Case $H \neq \frac{1}{2}$

Theorem

fBm is not semimartingale for $H \neq \frac{1}{2}$.

$$dX_t = -aX_t dt + \gamma dU_H(t),$$

where $(X_t)_{t \geq 0}$ is a stochastic process, $a, \gamma \in \mathbb{R}_+$ and $(U_H(t))_{t \geq 0}$ fBm with Hurst exponent H .

$$X_t(\omega) = X_0(\omega) - a \int_0^t X_u(\omega) du + \gamma U_H(t)(\omega)$$

for $t \geq 0$.

Stationary solution: $\hat{X}_{H,t} := e^{-at} \left(\gamma \int_{-\infty}^t e^{au} dU_H(u) \right)$.

Theorem

$(\hat{X}_{H,t})_{t \geq 0}$ is centered Gaussian process.

Theorem

$(\hat{X}_{H,t})_{t \geq 0}$ has long memory for $H \in (\frac{1}{2}, 1)$.

Fractional Black-Scholes

$$A_t = \exp(rt)$$

$$S_t = \exp(rt + \mu(t) + \sigma U_H(t)), t \in [0, T],$$

where $r \in \mathbb{R}, \sigma \in \mathbb{R}_+, \sup_{t \in [0, T]} \mu(t) < \infty$.

Theorem

$$\tilde{S}_t = \exp(\mu(t) + \sigma U_H(t)),$$

if there exists

$$\xi_t^1 = f_0 \mathbb{1}_{\{0\}}(t) + \sum_{k=1}^{n-1} f_k \mathbb{1}_{(\tau_k, \tau_{k+1}]}(t),$$

where $t \in [0, T]$, f_k is family of $\mathcal{F}_k^{U_H}$ -measurable function for $k \in \{1, \dots, n-1\}$. $0 = \tau_1 < \dots < \tau_n = T$ are stopping times with respect to $\mathcal{F}_{\tau_k}^{U_H}$ respectively, with $\tau_{k+1} - \tau_k \geq m$ for some $m > 0$. If there exists a $k \in \{0, \dots, n-1\}$ such that $\mathcal{P}[f_k \neq 0] > 0$, then

$$\mathcal{P}[(\xi^1 \cdot \tilde{S})_T < 0] > 0.$$

Fractional Black Scholes

The fractional Black-Scholes market is arbitrage-free, if there exists a minimal amount of time between two successive transactions,.

FSV

$$H > \frac{1}{2}$$

$$\begin{aligned} dS_t &= r_t S_t dt + \sigma_t S_t dB_t, \\ \sigma_t &= \exp\{X_t\} \\ dX_t &= -aX_t dt + \gamma dU_H(t), \end{aligned}$$

where $a, \gamma \in \mathbb{R}_+$.

Stationary solution :

$$\hat{X}_{H,t} = e^{-at} \gamma \int_{-\infty}^t e^{au} dU_H(u)$$

FSV

$$\hat{\sigma}_{H,t} = \exp\{\hat{X}_{H,t}\}$$

Theorem

$(\hat{\sigma}_{H,t})$ has long memory for $H \in (\frac{1}{2}, 1)$.

RFSV

$$H < \frac{1}{2}$$

$$dS_t = r_t S_t dt + \sigma_t S_t dB_t,$$

$$\sigma_t = \exp\{X_t\}$$

$$dX_t = -aX_t dt + \gamma dU_H(t),$$

RFSV

Smoothness of σ_t .

$$s(\tau, \sigma) = \frac{1}{N} \sum_{k=1}^N |\log(\sigma_{k\tau}) - \log(\sigma_{(k-1)\tau})|^2,$$

Empirical result:

$$s(\tau, \sigma) = k\tau^z.$$

RFSV

Gatheral et la.[3]

Theorem

$$s(\tau, \hat{X}_H) \rightarrow \gamma^2 \tau^{2H}$$

as a goes to zero, for $t > 0, \tau > 0$.

Motivation

Find a model which combine advatages on both FSV and RFSV.

Weighted fractional Brownian motion

Definition

A *weighted fractional Brownian motion* is defined as follows

$$M_{\alpha,\beta,H_1,H_2}(t) = \alpha U_{H_1}(t) + \beta U_{H_2}(t)$$

for $t \in \mathbb{R}$, where $\alpha, \beta \in \mathbb{R}_+$ such that $\alpha^2 + \beta^2 = 1$ and U_{H_1}, U_{H_2} are two independent fBm's with Hurst exponents $H_1 \in (0, \frac{1}{2})$, $H_2 \in (\frac{1}{2}, 1)$ respectively.

Weighted fractional Brownian motion

$$H_1 < \frac{1}{2}, H_2 > \frac{1}{2}$$

$$\hat{X}_{\alpha,\beta,H_1,H_2}(t) = \alpha\gamma e^{-at} \int_{-\infty}^t e^{au} dU_{H_1} + \beta\gamma e^{-at} \int_{-\infty}^t e^{au} dU_{H_2}.$$

Weighted fractional Brownian motion

Proposition

\hat{X}_t satisfies following properties:

- (i) $(\hat{X}_t)_{t \geq 0}$ is a centered Gaussian stationary process.
- (ii) $(\hat{X}_t)_{t \geq 0}$ has long memory.

Weighted fractional Brownian motion

Proposition

\hat{X}_t satisfies following properties:

- (i) $E[\sup_{t \in [0, T]} |\hat{X}_{\alpha, \beta, H_1, H_2}(t) - U_{H_1}(t)|] \rightarrow 0$ as $a \rightarrow 0, \alpha \rightarrow 1$.
- (ii) $E[|\hat{X}_{\alpha, \beta, H_1, H_2}(t + \tau) - \hat{X}_{\alpha, \beta, H_1, H_2}(t)|^2] \rightarrow \gamma^2 \tau^{2H_1}$ as $a \rightarrow 0, \alpha \rightarrow 1$.




Summary

FSV ensures the volatility process has long memory.

RFSV demonstrates a reasonable smoothness of volatility.

Weighted-FSV inherits long memory of FSV and 'good' smoothness of RFSV.

**Thank you for your Attention !
Vielen Dank für Ihre Aufmerksamkeit !**

-  MANDELBROT, B.B., & VAN NESS, J.W. (1968). Fractional Brownian Motions, Fractional Noises and Applications. SIAM Review 10(4):422–437.
-  SAMORODNITSKY, G., & TAQQU, M. (1994). Stable Non-Gaussian Random Processes: Stochastic Models with Infinite Variance. New York: Chapman & Hall.
-  GATHERAL, J. & JAISSON, T. & ROSENBAUM, M (2014). Volatility is Rough.