

WIP Title

Kexing Ying

January 1, 2023

Contents

1	Introduction	3
1.1	Structure of this essay	3
2	Stochastic Localization Scheme	4
2.1	Linear-tilt localization schemes	6
2.1.1	Linear-tilt localization driven by a Wiener process	6
2.1.2	Discrete time linear-tilt localization	8
3	Application: Markov Mixing	10
3.1	Mixing bounds	10
3.1.1	Ising model and Glauber dynamics	11
3.2	Dynamics of stochastic localizations	12
3.2.1	ACV of discrete time linear-tilt localization	14
3.2.2	ACV of linear-tilt localization driven by a Wiener process	15
3.3	Glauber dynamics as an associated Markov process	15
4	Application: The KLS and Thin-Shell Conjecture	17
4.1	Concentration	17
4.2	Example: concentration of the Gaussian	19
4.3	The KLS and thin-shell conjecture	19
4.3.1	Equivalent formulation of the KLS conjecture	20
4.4	Reduction of KLS to thin-shell	21
4.4.1	Differential of the quadratic variation	22
4.4.2	Analysis of the covariance matrix	23
4.4.3	Stopping the process early	26

1 Introduction

1.1 Structure of this essay

2 Stochastic Localization Scheme

We will in this section introduce the notion of stochastic localization schemes. To gain an intuition for these objects, we will present several examples which will be studied further in subsequent sections.

We will work in general Borel spaces (\mathcal{X}, Σ) for this section while for restrict our focus to either the Euclidean space \mathbb{R}^n or the Boolean hypercube $\{-1, 1\}^n$ in subsequent sections. We take $(\Omega, \mathcal{F}, \mathbb{P})$ our underlying probability space and we introduce the notation $\mathcal{M}(\mathcal{X})$ for the space of probability measures on \mathcal{X} .

Definition 2.1 (Prelocalization process). Given $\mu \in \mathcal{M}(\mathcal{X})$, a measure-valued stochastic process $(\mu_t)_{t \geq 0}$ is said to be a prelocalization of μ if

$$(L0) \quad \mu_0 = \mu.$$

$$(L1) \quad \text{For all } t \geq 0, \mu_t \text{ is a probability measure almost everywhere, i.e. } \mathbb{P}(\mu_t(\mathcal{X}) = 1) = 1.$$

$$(L2) \quad \text{For all } A \in \Sigma, (\mu_t(A))_{t \geq 0} \text{ is a martingale with respect to the natural filtration of } (\mu_t).$$

Where $\mathcal{M}(\mathcal{X})$ is equipped with the σ -algebra generated by maps of the form

$$\pi_A : \mathcal{M}(\mathcal{X}) \mapsto \mathbb{R}_{\geq 0} \cup \{\infty\} : \mu \mapsto \mu(A)$$

for all $A \in \Sigma$. Equivalently, this is the Borel σ -algebra on $\mathcal{M}(\mathcal{X})$ using the topology induced by the total variation norm.

Definition 2.2 (Stochastic localization process, [CE22]). Given $\mu \in \mathcal{M}(\mathcal{X})$, a measure-valued stochastic process $(\mu_t)_{t \geq 0}$ is said to be a stochastic localization of μ if in addition to being a prelocalization of μ , $(\mu_t)_{t \geq 0}$ also satisfies

$$(L3) \quad \text{For all } A \in \Sigma, \mu_t(A) \text{ converges almost everywhere to 0 or 1 as } t \rightarrow \infty.$$

Definition 2.3 (Stochastic localization scheme, [CE22]). Denoting $\mathcal{L}(\mu)$ the set of all stochastic localization processes of the measure μ , a stochastic localization scheme is a map

$$\Phi : \mathcal{M}(\mathcal{X}) \rightarrow \coprod_{\mu \in \mathcal{M}(\mathcal{X})} \mathcal{L}(\mu)$$

such that $\Phi(\mu) \in \mathcal{L}(\mu)$ for all $\mu \in \mathcal{M}(\mathcal{X})$.

We say a stochastic localization is discrete if t takes value in \mathbb{N} and continuous if t takes value in $\mathbb{R}_{\geq 0}$. For shorthand, we denote $(\mu_k)_k$ for a discrete stochastic localization of μ .

Proposition 2.1. Straightaway, by the martingale property, if $(\mu_t)_{t \geq 0}$ is a stochastic localization of μ , then

- $\mathbb{E}[\mu_t] = \mu$ for all $t \geq 0$.
- taking $X \sim \mu$ such that $\mu_t \rightarrow \delta_X$ almost everywhere as $t \rightarrow \infty$ (here weak and total variational convergence are equivalent and so \rightarrow can mean either).

Proof. The first statement is immediate as for all $A \in \Sigma$,

$$\mathbb{E}[\mu_t](A) \triangleq \mathbb{E}[\mu_t(A)] = \mathbb{E}[\mu_0(A)] = \mu(A).$$

To prove the second statement, let us first parse what the claim is. Fixing a realization ω of μ_t , we have by (L3) that μ_t converges to some Dirac measure based at some $x_\omega \in \mathcal{X}$. Thus, defining the random variable $X : \omega \mapsto x_\omega$, it suffices to show $X \sim \mu$. Indeed, by taking $\phi : \mathcal{X} \rightarrow \mathbb{R}$ to be any bounded and continuous function, by the definition of X

$$\int \phi(x) \mu_t(dx) \xrightarrow{\text{a.e.}} \int \phi(x) \delta_X(dx) = \phi(X) \text{ as } t \rightarrow \infty.$$

Thus, taking expectation on both sides, we have

$$\mathbb{E}[\phi(X)] = \mathbb{E}\left[\int \phi d\mu_t\right] = \int \phi d\mathbb{E}[\mu_t] = \int \phi d\mu$$

implying $X \sim \mu$ as required. \square

An example of a stochastic localization scheme is the coordinate by coordinate localization scheme on $\mathcal{X} = \{-1, 1\}^n$. This scheme relates to the Glauber dynamics for which the stochastic localization scheme provides a mixing bound. We shall examine the property in section 3.3, though we will construct the scheme now.

Given a probability measure μ on $\{-1, 1\}^n$, we introduce the random variable $X \sim \mu$, and Y a uniform random variable over all permutations of $[n] = \{1, \dots, n\}$ independent of X . Then, the coordinate by coordinate stochastic localization of μ is the process $(\mu_k)_k$ such that for all $x \in \{-1, 1\}^n$,

$$\mu_k(x) = \mathbb{P}(X = x \mid X_{Y_1}, \dots, X_{Y_{n \wedge k}}).$$

Namely, μ_k is the law of X conditioned on X_{Y_1}, \dots, X_{Y_k} .

$(\mu_k)_k$ is indeed a stochastic localization of μ . It is clear that (L0) and (L1) are satisfied. By construction of $(\mu_k)_k$, denoting $\mathcal{F}_k := \sigma(X_{Y_1}, \dots, X_{Y_{n \wedge k}})$, we have by the tower property

$$\mathbb{E}[\mu_{k+1}(x) \mid \mathcal{F}_k] = \mathbb{E}[\mathbb{E}[\mathbb{P}(X = x \mid X) \mid \mathcal{F}_{k+1}] \mid \mathcal{F}_k] = \mathbb{E}[\mathbb{P}(X = x \mid X) \mid \mathcal{F}_k] = \mu_k(x)$$

implying $(\mu_k(x))$ a martingale as required for (L2). Finally, it is clear that

$$\lim_{k \rightarrow \infty} \mu_k(x) = \mu_n(x) = \mathbb{P}(X = x \mid X) = \mathbf{1}_{\{X=x\}} \in \{0, 1\}$$

implying (L3).

An analogous construction of the coordinate by coordinate stochastic localization scheme in \mathbb{R}^n is the random subspace localization. Similar to before, for a probability measure μ on \mathbb{R}^n , we introduce the random variable $X \sim \mu$ and Y a uniform random variable on $O(n)$ (so the column vectors $\{Y_1, \dots, Y_n\}$ form an orthonormal basis of \mathbb{R}^n) independent of X . Then, we define the random subspace stochastic localization of μ as $(\mu_k)_k$ where μ_k is the law of X conditioned on $\langle X, Y_1 \rangle, \dots, \langle X, Y_{n \wedge k} \rangle$.

2.1 Linear-tilt localization schemes

An important class of stochastic localization schemes are the linear-tilt schemes. Introduced by Eldan in [Eld13], linear-tilt schemes has been vital in the recent progress regarding the KLS conjecture. More recently, a discrete version of the linear-tilt scheme was introduced in [CE22] and is used to provide a mixing bound for Glauber dynamics. We will in this section introduce these family of localizations and consider two specific examples of such linear-tilt schemes which are useful for our analysis later.

Informally, given a probability measure μ on $\mathcal{X} \subseteq \mathbb{R}^n$, the linear-tilt scheme of μ is constructed recursively in which at each step, we pick a random direction and multiply the density at this time with a linear function along this direction (i.e. a tilt along a random direction).

Let μ be a probability measure on $\mathcal{X} \subseteq \mathbb{R}^n$, we introduce the following definition.

Definition 2.4 (Barycenter). The barycenter of μ with respect to the function $F : \mathbb{R}^n \rightarrow \mathbb{R}$ is

$$\bar{\mu}(F) := \int_{\mathcal{X}} xF(x)\mu(dx).$$

In the case that $F = \text{id}$, we simply write $\bar{\mu} = \bar{\mu}(F) = \mathbb{E}_{X \sim \mu}[X]$.

Definition 2.5 (Linear-tilt localization). A measure-valued stochastic process $(\mu_t)_{t \geq 0}$ is said to be a linear-tilt localization of the probability measure μ if

1. $\mu_t \ll \mu$ for each $t \geq 0$, and
2. denoting $F_t := d\mu_t/d\mu$, we have $F_0 = 1$ and

$$dF_t(x) = \langle x - \bar{\mu}(F_t), dZ_t \rangle F_t(x) \quad (1)$$

for some stochastic process $(Z_t)_{t \geq 0}$ such that $\mathbb{E}[dZ_t | \mu_t] = 0$ for all $t \geq 0$.

It is clear that $(\mu_t(A))_t$ is a martingale for all $A \in \Sigma$ by observing that equation (1) has no drift term. Hence, it follows $(\mu_t(\mathcal{X}))$ is a martingale. However, since $\mu_t(\mathcal{X}) = \int F_t d\mu$ is differentiable by construction, $(\mu_t(\mathcal{X}))$ has zero quadratic variation and thus is constant in t . With this in mind, as $\mu_0 = \mu$ is a probability measure, it follows:

Proposition 2.2. If $(\mu_t)_t$ is a linear-tilt localization of μ , then μ_t is a probability measure for each t .

Corollary 2.3. A linear-tilt localization $(\mu_t)_t$ of μ is a prelocalization of μ .

We remark that in general, a linear-tilt localization is not necessarily a stochastic localization as (L3) might not be satisfied. It is possible to impose sufficient conditions on (Z_t) for which (L3) holds, e.g. by requiring $\|\text{Cov}(Z_t)\|_{\text{op}}$ to decrease sufficiently fast. However, for generality, we will not restrict ourselves to one of these conditions. Instead, we will consider (L3) case by case in the following examples of linear-tilt schemes.

I think Eldan's proof in [Eld13] has a mistake

2.1.1 Linear-tilt localization driven by a Wiener process

A natural choice of $(Z_t)_{t \geq 0}$ is the standard Wiener process on \mathbb{R}^n . Denoting $(W_t)_{t \geq 0}$ a standard Wiener process on \mathbb{R}^n , we define the random functions $(F_t)_{t \geq 0}$ to be the solution of the following infinite system of SDEs (existence and uniqueness is established by theorem 5.2 in [Øks03]):

$$F_0 = 1, dF_t(x) = \langle x - \bar{\mu}(F_t), dW_t \rangle F_t(x), \quad (2)$$

for all $x \in \mathbb{R}^n$. We shall from this point forward denote the random variables $a_t := \bar{\mu}(F_t)$. By applying Itô's formula, we make the following useful observation: for all $x \in \mathbb{R}^n$,

$$d \log F_t(x) = \frac{dF_t(x)}{F_t(x)} - \frac{d[F(x)]_t}{2F_t(x)^2} = \langle x - a_t, dW_t \rangle - \frac{1}{2} \|x - a_t\|^2 dt \quad (3)$$

where the second equality follows by the construction of F . Hence, as $\log F_0(x) = 0$, we observe

$$\begin{aligned} \log F_t(x) &= \int_0^t \langle x - a_s, dW_s \rangle - \frac{1}{2} \int_0^t \|x - a_s\|^2 ds \\ &= \left(\langle x, W_t \rangle - \int_0^t \langle a_s, dW_s \rangle \right) - \left(\frac{t}{2} \|x\|^2 + \frac{1}{2} \int_0^t \|a_s\|^2 ds - \int_0^t \langle x, a_s \rangle ds \right) \\ &= - \left(\int_0^t \langle a_s, dW_s \rangle + \frac{1}{2} \|a_s\|^2 ds \right) + \langle x, a_t + W_t \rangle - \frac{t}{2} \|x\|^2. \end{aligned}$$

Thus, taking $dz_t := \langle a_t, dW_t \rangle + \frac{1}{2} \|a_t\|^2 dt$ and $v_t := a_t + W_t$, we observe $F_t(x)$ is of the form

$$F_t(x) = e^{z_t + \langle x, v_t \rangle - \frac{t}{2} \|x\|^2}, \quad (4)$$

for given Itô processes $(z_t), (v_t)$.

With this formulation of $F_t(x)$ in mind, it follows F_t is non-negative, and so, we may define $(\mu_t)_t$ to be the process such that $d\mu_t = F_t d\mu$. It is clear that $(\mu_t)_t$ is a linear-tilt localization of μ and so, is a prelocalization of μ . The remainder of this section is devoted to showing $(\mu_t)_t$ is furthermore a stochastic localization of μ if μ is log-concave (namely we will show (L3) for this special case), and prove some basic properties about this process useful for our analysis later.

Definition 2.6 (Log-concave measure). A measure μ on \mathbb{R}^n is said to log-concave if it is of the form $d\mu = \exp(-H) d\text{Leb}^n$ for some convex function $H : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$

To show (μ_t) satisfies (L3) if μ is log-concave, we study the limiting behavior of (μ_t) as $t \rightarrow \infty$ by considering their covariances:

$$A_t := \text{Cov}[\mu_t] = \int (x - a_t) \otimes (x - a_t) \mu_t(dx), \quad (5)$$

where \otimes denotes the Kronecker product. In particular, we will show $(A_t)_{ij} \rightarrow 0$ for all $i, j \in \{1, \dots, n\}$ as $t \rightarrow \infty$ allowing us to conclude (μ_t) converges weakly to some Dirac measure. Indeed, this is a direct consequence of the following lemma.

Lemma 2.4 (Brascamp-Lieb inequality, [BL76]). Given $V : \mathbb{R}^n \rightarrow \mathbb{R}$ convex and $K > 0$, if ν is an isotropic probability measure on \mathbb{R}^n of the form

$$d\nu = Z e^{-V(x) - \frac{1}{2K} \|x\|^2} d\text{Leb}^n$$

with Z being the normalization constant, then ν satisfy the Poincaré inequality, i.e. for all differentiable ϕ ,

$$K \text{Var}_\nu[\phi] \leq \int \|\nabla \phi\|^2 d\nu.$$

With this lemma in mind, by taking $\nu = \mu_t$ using equation (4) and defining $\pi_i(x) := x_i$, we have by the Cauchy-Schwarz inequality

$$(A_t)_{ij} \leq \sqrt{\text{Var}_{\mu_t}[\pi_i]} \sqrt{\text{Var}_{\mu_t}[\pi_j]} \leq \max_{k=1, \dots, n} \frac{1}{t} \int \|\nabla \pi_k\|^2 d\mu_t$$

Again, using equation (4), we note that any realizations of $(F_t(x))$ is eventually decreasing in t for all $x \neq 0$, implying

$$\sup_{t>0} \max_{k=1, \dots, n} \int \|\nabla \pi_k\|^2 d\mu_t = \sup_{t>0} \max_{k=1, \dots, n} \int x_k^2 d\mu_t < \infty.$$

Thus, by taking $t \rightarrow \infty$ we have $(A_t)_{ij} \rightarrow 0$ for all $i, j \in \{1, \dots, n\}$ as claimed and we have (μ_t) satisfying (L3).

Corollary 2.5. (μ_t) converges set-wise to some Dirac measure almost everywhere. We denote this limiting (random) Dirac measure by δ_{a_∞} where a_∞ is some \mathbb{R}^n -valued random variable.

As a result of 2.1, we have the following useful corollary.

Corollary 2.6. The massive point a_∞ of the limiting Dirac measure is the limit of a_t as $t \rightarrow \infty$ and has law μ .

Proof. Since convergence implies relatively compact, applying the Dunford-Pettis theorem it follows that any realizations of (F_t) is uniformly integrable. Thus, the result follows by the Vitali convergence theorem. \square

Corollary 2.7. Similarly, taking ϕ to be any continuous function (not necessarily bounded as we have uniform integrability), defining $M_t = \int \phi d\mu_t$, $(M_t)_t$ is a martingale and

$$M_t \xrightarrow{\text{a.e.}} M_\infty \sim \phi_* \mu \quad (6)$$

where $\phi_* \mu$ denotes the push-forward measure of μ along ϕ .

2.1.2 Discrete time linear-tilt localization

We may construct an analogous version of the linear-tilt localization for discrete time. By utilizing the little- o notation, equation (1) can be rewritten as

$$\frac{d\mu_{t+h}}{d\mu}(x) = \frac{d\mu_t}{d\mu}(x) + \langle x - \bar{\mu}_t, h dZ_t \rangle \frac{d\mu_t}{d\mu}(x) + o(h).$$

Hence, an discrete analog of the linear tilt localization is defined as the following.

Definition 2.7 (Discrete time linear-tilt localization). Given a measure $\mu \in \mathcal{M}(\mathcal{X})$, the discrete time linear-tilt localization is the sequence of random measures $(\mu_k)_k$ defined by $\mu_0 = \mu$ and

$$d\mu_{k+1} = (1 + \langle x - \bar{\mu}_k, Z_k \rangle) d\mu_k \quad (7)$$

for some sequence of random variables such that $\mathbb{E}[Z_k | \mu_k] = 0$ for all $k \in \mathbb{N}$.

Using the discrete time linear-tilt localization, let us now provide an alternative construction of the coordinate by coordinate localization.

Given μ a probability measure on $\{-1, 1\}^n$, we recall that the coordinate by coordinate localization is defined by “pinning” an additional random coordinate after each time step. To phrase this as a linear-tilt localization, we take the random variables Z_k to be

$$Z_k := e_{Y_k} \cdot \begin{cases} \frac{1}{1+(\bar{\mu}_k)_{Y_k}} & \text{with probability } \frac{1+(\bar{\mu}_k)_{Y_k}}{2} \\ \frac{-1}{1-(\bar{\mu}_k)_{Y_k}} & \text{with probability } \frac{1-(\bar{\mu}_k)_{Y_k}}{2} \end{cases} \quad (8)$$

where e_1, \dots, e_n are the standard basis of \mathbb{R}^n and again Y is a uniform random variable over all permutations of $[n]$. Thus, the linear-tilt localization (μ_k) given by this choice of (Z_k) is defined by

$$\mu_{k+1}(\sigma) = (1 + \|Z_k\|(\sigma_{Y_k} - (\bar{\mu}_k)_{Y_k}))\mu_k(\sigma)$$

for all $k < n$. Similar to before, we terminate the process at time n and so we extend the process to all times by taking $\mu_k = \mu_{n \wedge k}$.

Let us parse this definition to see why this is equivalent to the coordinate by coordinate localization. Taking $k < n$, we have at the $k + 1$ -th step, Y_{k+1} chooses a random axis which had not been chosen before. Then, Z_k is chosen such that for each configuration σ , the probability of σ_{Y_k} being ± 1 is proportional the mass of μ_k on $\pm e_{Y_k}$. This is precisely the steps needed to construct the coordinate by coordinate localization as conditioning on an additional axis in this case is simply a projection on to said axis.

3 Application: Markov Mixing

An application of stochastic localizations is used to prove mixing bounds for Markov processes. Expanding on the work of Anari, Liu and Oveis in [ALG20], Chen and Eldan in [CE22] established a framework in which mixing bounds of a special class of Markov processes arises, namely Markov chains associated with stochastic localizations. We will in this section present this framework and describe its application to the Glauber dynamics.

Fix!

3.1 Mixing bounds

The motivation for Markov mixing bounds fundamentally comes from sampling. Suppose we wish to sample from some probability distribution μ . A common method to achieve this to through the use of the Markov chain Monte Carlo (MCMC):

Theorem 1. Given (X_n) an irreducible positively recurrent homogenous Markov process on \mathcal{X} with stationary distribution μ , for any $\phi : \mathcal{X} \rightarrow \mathbb{R}$ integrable,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \phi(X_k) = \int \phi d\mu$$

almost everywhere.

With this theorem in mind, MCMC allows us to sample μ by sampling from (X_n) instead. It is in general not difficult to come up with such Markov processes, although it is often difficult to show its rate of convergence. This motivates the notion of mixing bounds which quantifies the time for which the Markov process takes before its law is approximately stationary.

Definition 3.1 (Total variation mixing time). Given a probability measure $\nu \in \mathcal{M}(\mathcal{X})$, a Markov kernel K with stationary distribution μ and some $\epsilon > 0$, the ϵ -total variation mixing time is defined as

$$t_{\text{mix}}(P, \epsilon, \mu) := \inf\{t \geq 0 \mid \|P^t \nu - \mu\|_{\text{TV}} < \epsilon\},$$

Furthermore, we denote

$$t_{\text{mix}}(P, \epsilon) = \sup_{x \in \mathcal{X}} t_{\text{mix}}(P, \epsilon, \delta_x)$$

the worst mixing time starting at a point.

A standard method of analyzing the mixing times of Markov chains is through the use of the spectral gap.

Definition 3.2 (Spectral gap, [Lev17]). The spectral gap of a Markov kernel K is defined to be

$$\text{gap}(K) := 1 - \sup\{\lambda \mid \lambda \text{ is an eigenvalue of } K, \lambda \neq 1\}.$$

It is not difficult to show that

$$\text{gap}(K) = \inf_{\substack{\phi : \mathcal{X} \rightarrow \mathbb{R} \\ \int \phi d\mu = 0}} 1 - \frac{\int \phi K \phi d\mu}{\int \phi^2 d\mu} = \inf_{\phi : \mathcal{X} \rightarrow \mathbb{R}} \frac{1}{2\text{Var}_\mu[\phi]} \int (\phi(x) - \phi(y))^2 K(x, dy) \mu(dx), \quad (9)$$

where μ is the stationary measure of K and $K\phi = \int \phi(y)K(\cdot, dy)$. We will take equation (9) to be the defining property of the spectral gap in the case K is defined on a general state space.

Theorem 2 ([Lev17]). Given a reversible and irreducible Markov chain with kernel K on the state space \mathcal{X} with stationary distribution μ , denoting $\mu_{\min} = \inf_{x \in \mathcal{X}} \mu(x)$, we have

$$t_{\text{mix}}(K, \epsilon) \leq \left\lceil \frac{1}{\text{gap}(K)} \left(\frac{1}{2} \log \left(\frac{1}{\mu_{\min}} \right) + \log \left(\frac{1}{2\epsilon} \right) \right) \right\rceil.$$

We remark that this inequality is only meaningful whenever $\mu_{\min} > 0$ and thus, this theorem is only meaningful for Markov chains on finite state space (and we can replace \inf with \min).

A similar bound can also be established using the modified log-Sobolev inequality (MLSI).

Definition 3.3 (Entropy, [CLV20]). Given $\phi : \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}$ and a measure μ , we define the entropy of ϕ with respect to μ to be

$$\text{Ent}_{\mu}[\phi] := \mathbb{E}_{\mu} \left[\phi \log \left(\frac{1}{\mathbb{E}_{\mu}[\phi]} \phi \right) \right] = \int \phi \log \phi d\mu - \int \phi d\mu \log \left(\int \phi d\mu \right).$$

Definition 3.4.

TODO

3.1.1 Ising model and Glauber dynamics

Definition 3.5 (Ising model). Given a graph $G = (V, E)$, $\beta > 0$ and $h \in \mathbb{R}$, the Ising model on G with inverse temperature β and external field h is the probability measure $\mu_{\beta, h}$ on $\{-1, 1\}^V$ defined such that for all $\sigma \in \{-1, 1\}^V$,

$$\mu_{\beta, h}(\sigma) := \frac{1}{Z} \exp \left\{ \beta \sum_{xy \in E} \sigma_x \sigma_y + h \sum_{x \in V} \sigma_x \right\}$$

where $Z > 0$ is the normalizing constant.

Heuristically, the Ising model measures the probability that a graph is in a specific configuration of up and down spins in which neighboring vertices are more likely to have the same spin. This “likeliness” is controlled by β in which a larger β means that neighboring vertices are more likely to align.

As illustrated by theorem 1, in order to sample from the Ising model, we can construct a Markov chain which has the Ising model as its stationary distribution. One such Markov chain is known as the Glauber dynamics.

Definition 3.6 (Glauber dynamics). Given a measure μ of $\{-1, 1\}^n$, the Glauber dynamics of μ is the Markov chain with kernel

$$K(\sigma^1, \sigma^2) := \mathbf{1}_{\{\|\sigma^1 - \sigma^2\|_1 = 1\}} \frac{1}{n} \frac{\mu(\sigma^2)}{\mu(\sigma^1) + \mu(\sigma^2)} + \mathbf{1}_{\{\sigma^1 = \sigma^2\}} \frac{1}{n} \sum_{\|\tilde{\sigma} - \sigma^1\|_1 = 1} \frac{\mu(\sigma^1)}{\mu(\sigma^1) + \mu(\tilde{\sigma})}, \quad (10)$$

where we define $\|\sigma^1 - \sigma^2\|_1 := \frac{1}{2} \sum_{i=1}^n |\sigma_i^1 - \sigma_i^2|$ for all $\sigma^1, \sigma^2 \in \{-1, 1\}^n$.

Parsing this definition, we see that the Glauber dynamics is the Markov chain such that, starting at a configuration $\sigma_1 \in \{-1, 1\}^n$, the configuration at the next time step either remains the same or change at one vertex. Furthermore, the probability of this occurring is weighted according to μ . As we hoped, the Glauber dynamics of the Ising model $\mu_{\beta, h}$ has $\mu_{\beta, h}$ as its stationary distribution.

3.2 Dynamics of stochastic localizations

As alluded to previously, one may associate a Markov process at each time step of a stochastic localization process for which the original process is stationary. We will in this section define these Markov processes and show that the Glauber dynamics can be constructed using this method.

Definition 3.7 (Markov process associated with a stochastic localization, [CE22]). Let $(\mu_t)_{t \geq 0}$ be a prelocalization of μ such that μ_t is absolutely continuous (almost everywhere) with respect to μ for all t . For all $\tau > 0$, we define the dynamics associated with $(\mu_t)_t$ at τ to be the Markov process with kernel

$$K(x, A) := \mathbb{E}_{\mathbb{P}} \left[\frac{d\mu_{\tau}}{d\mu}(x) \mu_{\tau}(A) \right]$$

for all $x \in \mathcal{X}, A \in \Sigma$.

As alluded by the notation, rather than a deterministic time τ , τ can be taken to be an appropriate stopping time. In this case, the theorems below will remain to hold by invoking the optional stopping theorem whenever necessary.

This is indeed a kernel since for each $x \in \mathcal{X}$, as $\mathbb{E}_{\mathbb{P}}[\mu_{\tau}] = \mu$ by (L2),

$$K(x, \Omega) = \mathbb{E}_{\mathbb{P}} \left[\frac{d\mu_{\tau}}{d\mu}(x) \right] = \frac{d}{d\mu} \mathbb{E}_{\mathbb{P}}[\mu_{\tau}](x) = \frac{d\mu}{d\mu}(x) = 1$$

where the third equality follows by the uniqueness of the Radon-Nikodym derivative as for all $A \in \Sigma$, we have

$$\int_A \mathbb{E}_{\mathbb{P}} \left[\frac{d\mu_{\tau}}{d\mu} \right] d\mu = \mathbb{E}_{\mathbb{P}} \left[\int_A \frac{d\mu_{\tau}}{d\mu} d\mu \right] = \mathbb{E}_{\mathbb{P}}[\mu_{\tau}](A).$$

Proposition 3.1. The Markov process associated with a prelocalization $(\mu_t)_{t \geq 0}$ of μ is reversible and has stationary distribution μ .

Proof. Taking $\phi : \mathcal{X}^2 \rightarrow \mathbb{R}$ integrable, we have by Fubini's theorem

$$\begin{aligned} \int_{\mathcal{X}^2} \phi(x, y) K(x, dy) \mu(dx) &= \int \phi(x, y) \mathbb{E}_{\mathbb{P}} \left[\frac{d\mu_{\tau}}{d\mu}(x) \mu_{\tau}(dy) \right] \mu(dx) \\ &= \mathbb{E}_{\mathbb{P}} \left[\int \phi(x, y) \mu_{\tau}(dy) \frac{d\mu_{\tau}}{d\mu}(x) \mu(dx) \right] \\ &= \mathbb{E}_{\mathbb{P}} \left[\int \phi(x, y) \mu_{\tau}(dy) \mu_{\tau}(dx) \right]. \end{aligned} \tag{11}$$

Similarly, by the same calculation,

$$\int_{\mathcal{X}^2} \phi(x, y) K(y, dx) \mu(dy) = \mathbb{E}_{\mathbb{P}} \left[\int \phi(x, y) \mu_{\tau}(dy) \mu_{\tau}(dx) \right].$$

Thus,

$$\int_{\mathcal{X}^2} \phi(x, y) K(x, dy) \mu(dx) = \int_{\mathcal{X}^2} \phi(x, y) K(y, dx) \mu(dy)$$

for any integrable $\phi : \mathcal{X}^2 \rightarrow \mathbb{R}$ implying K is reversible.

On the other hand, for all $A \in \Sigma$, we compute using the martingale property

$$\begin{aligned} K\mu(A) &= \int K(x, A)\mu(dx) = \int \mathbb{E}_{\mathbb{P}} \left[\frac{d\mu_{\tau}}{d\mu}(x) \mu_{\tau}(A) \right] = \mathbb{E}_{\mathbb{P}} \left[\mu_{\tau}(A) \int \frac{d\mu_{\tau}}{d\mu} d\mu \right] \\ &= \mathbb{E}_{\mathbb{P}}[\mu_{\tau}(A)\mu_{\tau}(\Omega)] = \mathbb{E}_{\mathbb{P}}[\mu_{\tau}(A)] = \mu(A) \end{aligned}$$

implying μ is the stationary measure of K . \square

Proposition 3.2. Taking K to be the kernel of the Markov process associated with a prelocalization $(\mu_t)_{t \geq 0}$ of μ at time τ , we have

$$\text{gap}(K) = \inf_{\phi: \mathcal{X} \rightarrow \mathbb{R}} \frac{\mathbb{E}[\text{Var}_{\mu_{\tau}}[\phi]]}{\text{Var}_{\mu}[\phi]}.$$

Proof. By equation (11) (where we take $\phi(x, y) = \phi(x)\phi(y)$), we have

$$\int_{\mathcal{X}^2} \phi(x)\phi(y)K(x, dy)\mu(dx) = \mathbb{E}_{\mathbb{P}} \left[\left(\int_{\mathcal{X}} \phi d\mu_{\tau} \right)^2 \right], \quad (12)$$

for any integrable $\phi: \mathcal{X} \rightarrow \mathbb{R}$. On the other hand, we observe

$$\int_{\mathcal{X}^2} \phi(y)^2 K(x, dy)\mu(dx) = \int (K\phi^2)(x)\mu(dx) = \int \phi(x)^2 (K\mu)(dx) = \int \phi(x)^2 \mu(dx)$$

as μ is the stationary measure of K . Thus, for any integrable $\phi: \mathcal{X} \rightarrow \mathbb{R}$, by substituting the above two equations, we have

$$\begin{aligned} & \frac{1}{2\text{Var}_{\mu}[\phi]} \int_{\mathcal{X}^2} (\phi(x) - \phi(y))^2 K(x, dy)\mu(dx) \\ &= \frac{1}{2\text{Var}_{\mu}[\phi]} \left(\int \phi(x)^2 \mu(dx) - 2 \int \phi(x)\phi(y)K(x, dy)\mu(dx) + \int \phi(y)^2 K(x, dy)\mu(dx) \right) \\ &= \frac{1}{\text{Var}_{\mu}[\phi]} \left(\int \phi^2 d\mu - \mathbb{E}_{\mathbb{P}} \left[\left(\int_{\mathcal{X}} \phi d\mu_{\tau} \right)^2 \right] \right) \\ &= \frac{1}{\text{Var}_{\mu}[\phi]} \mathbb{E}_{\mathbb{P}} \left[\int \phi^2 d\mu_{\tau} - \left(\int_{\mathcal{X}} \phi d\mu_{\tau} \right)^2 \right] = \frac{\mathbb{E}[\text{Var}_{\mu_{\tau}}[\phi]]}{\text{Var}_{\mu}[\phi]}. \end{aligned}$$

Hence, recalling the equivalent form of the spectral gap as described by (9), the result follows by taking infimum on both sides. \square

With regards to theorem 2, in the case μ has full support on the finite state space \mathcal{X} (e.g. in the setting of Glauber dynamics) the above proposition provides a method for computing an upper bound for the mixing time. In particular, should the prelocalization $(\mu_k)_k$ satisfy

$$\mathbb{E}[\text{Var}_{\mu_{k+1}}[\phi] \mid \mu_t] \geq (1 - \epsilon)\text{Var}_{\mu_k}[\phi]$$

for given $\epsilon > 0$ and any integrable function $\phi: \mathcal{X} \rightarrow \mathbb{R}$, we have the telescoping product

$$\frac{\mathbb{E}[\text{Var}_{\mu_k}[\phi]]}{\text{Var}_{\mu}[\phi]} = \mathbb{E} \left[\prod_{i=1}^k \frac{\mathbb{E}[\text{Var}_{\mu_i}[\phi] \mid \mu_{i-1}]}{\text{Var}_{\mu_{i-1}}[\phi]} \right] \geq (1 - \epsilon)^k.$$

Hence, we have the bound

$$\text{gap}(K)^{-1} \leq (1 - \epsilon)^{-k}$$

which immediately provides an upper bound for the mixing time of K in light of theorem 2. This motivates the following definition.

Definition 3.8 (Discrete time approximate conservation of variance (ACV), [CE22]). A prelocalization process $(\mu_k)_k$ is said satisfy conserve (κ_k) -variance up to time n if for any integrable function $\phi : \mathcal{X} \rightarrow \mathbb{R}$, $1 \leq k \leq n$,

$$\mathbb{E}[\text{Var}_{\mu_{k+1}}[\phi] \mid \mu_k] \geq (1 - \kappa_k) \text{Var}_{\mu_k}[\phi].$$

By the same computation above, if $(\mu_k)_k$ conserves (κ_k) -variance up to time n , then it associated dynamics has a spectral gap of at least $\prod_{i=1}^n (1 - \kappa_i)$.

3.2.1 ACV of discrete time linear-tilt localization

In the case of the discrete time linear-tilt localization $(\mu_k)_k$ of some μ , we may compute an explicit value (η_k) for which $(\mu_k)_k$ conserves (η_k) -variance. Indeed, given integrable $\phi : \mathcal{X} \rightarrow \mathbb{R}$, we have

$$\begin{aligned} \mathbb{E}[\text{Var}_{\mu_{k+1}}[\phi] \mid \mu_k] &= \mathbb{E} \left[\int \phi^2 d\mu_{k+1} - \left(\int \phi d\mu_{k+1} \right)^2 \mid \mu_k \right] \\ &= \int \phi^2 d\mu_k - \mathbb{E} \left[\left(\int \phi(x) (1 + \langle x - \bar{\mu}_k, Z_k \rangle) \mu_k(dx) \right)^2 \mid \mu_k \right] \\ &= \int \phi^2 d\mu_k - \left(\int \phi d\mu_k \right)^2 - \mathbb{E} \left[\left(\int \phi(x) \langle x - \bar{\mu}_k, Z_k \rangle \mu_k(dx) \right)^2 \mid \mu_k \right] \\ &= \text{Var}_{\mu_k}[\phi] - \text{Var} \left[\int \phi(x) \langle x - \bar{\mu}_k, Z_k \rangle \mu_k(dx) \mid \mu_k \right] \end{aligned} \quad (13)$$

where the third and last equality follows as $\mathbb{E}[Z_k \mid \mu_k] = 0$. Then, denoting

$$C_k := \text{Cov}(Z_k \mid \mu_k) \text{ and } U_k := \int \phi(x) (x - \bar{\mu}_k) \mu_k(dx),$$

equation (13) becomes

$$\mathbb{E}[\text{Var}_{\mu_{k+1}}[\phi] \mid \mu_k] = \text{Var}_{\mu_k}[\phi] - \|C_k^{1/2} U_k\|^2.$$

With this reduction, our goal is reduced to bounding $\|C_k^{1/2} U_k\|^2$ by an expression of the form $\eta_k \text{Var}_{\mu_k}[\phi]$ where η_k is independent of ϕ . Indeed, taking θ to be the normalized vector $C_k^{1/2} U_k / \|C_k^{1/2} U_k\|$, we have

$$\begin{aligned} \|C_k^{1/2} U_k\|^2 &= \langle C_k^{1/2} U_k, \theta \rangle^2 = \left(\int \phi(x) \langle C_k^{1/2} (x - \bar{\mu}_k), \theta \rangle \mu_k(dx) \right)^2 \\ &= \left(\int \left(\phi(x) - \int \phi d\mu_k \right) \langle C_k^{1/2} (x - \bar{\mu}_k), \theta \rangle \mu_k(dx) \right)^2 \\ &\leq \text{Var}_{\mu_k}[\phi] \int \langle C_k^{1/2} (x - \bar{\mu}_k), \theta \rangle^2 \mu_k(dx) \leq \text{Var}_{\mu_k}[\phi] \|C_k^{1/2} \text{Cov}(\mu_k) C_k^{1/2}\|_{\text{op}} \end{aligned}$$

where the inequality is due to Cauchy-Schwarz. Hence, for any discrete linear-tilt localization $(\mu_k)_k$, we have

$$\mathbb{E}[\text{Var}_{\mu_{k+1}}[\phi] \mid \mu_k] \geq (1 - \|C_k^{1/2} \text{Cov}(\mu_k) C_k^{1/2}\|_{\text{op}}) \text{Var}_{\mu_k}[\phi] \quad (14)$$

for any integrable $\phi : \mathcal{X} \rightarrow \mathbb{R}$ and thus, conserve $(1 - \eta_k)$ -variance where we define $\eta_k := \|C_k^{1/2} \text{Cov}(\mu_k) C_k^{1/2}\|_{\text{op}}$.

3.2.2 ACV of linear-tilt localization driven by a Wiener process

In the setting of the linear-tilt localization driven by a Wiener process, the computation above remains to work with slight modifications. Namely, rather than considering the discrete differences of the variance, we instead compute and attempt to bound the time derivative of the variance from below. This computation turns out to be very useful with regards to the KLS conjecture which will be discussed in section 4.

Definition 3.9 (Approximate conservation of variance). A prelocalization process $(\mu_t)_{t \geq 0}$ satisfies approximate (κ_t) -variance conservation if for all integrable $\phi : \mathcal{X} \rightarrow \mathbb{R}$ and $t \geq 0$,

$$\frac{d\text{Var}_{\mu_t}[\phi]}{\text{Var}_{\mu_t}[\phi]} \geq -\kappa_t dt + N_t,$$

for some martingale (N_t) .

3.3 Glauber dynamics as an associated Markov process

As alluded to in section 2, Glauber dynamics can be constructed as an associated Markov chain of the coordinate by coordinate localization. Namely, taking $\mu \in \mathcal{M}(\{-1, 1\}^n)$ and $\tau = n - 1$, we will show the associate Markov kernel

$$\mathbb{E} \left[\frac{\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2)}{\mu(\sigma^1)} \right] = K(\sigma^1, \sigma^2)$$

for all $\sigma^1, \sigma^2 \in \{-1, 1\}^n$ where $K(\sigma^1, \sigma^2)$ as defined by equation (10).

First, taking $\sigma^1, \sigma^2 \in \{-1, 1\}^n$ such that $\|\sigma^1 - \sigma^2\| = 1$, say σ^1 and σ^2 differs at the m -th coordinate, we have

$$\mathbb{E} \left[\frac{\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2)}{\mu(\sigma^1)} \right] = \frac{1}{n} \sum_{k=1}^n \mathbb{E} \left[\frac{\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2)}{\mu(\sigma^1)} \mid Y_n = k \right].$$

Now, if $Y_n = k \neq m$, the configuration is fixed at the m -th coordinate. However, as σ^1 and σ^2 differs at the m -th coordinate, either $\mu_{n-1}(\sigma^1) = 0$ or $\mu_{n-1}(\sigma^2) = 0$. Thus, all but the m -th term

in the above sum vanishes and we have the kernel equals

$$\begin{aligned}
&= \frac{1}{n} \mathbb{E} \left[\frac{\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2)}{\mu(\sigma^1)} \middle| Y_n = m \right] \\
&= \frac{1}{n} \mathbb{E} \left[\frac{\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2)}{\mu(\sigma^1)} \mathbf{1}_{\{\text{supp } \mu_{n-1} = \{\sigma^1, \sigma^2\}\}} \middle| Y_n = m \right] \\
&= \frac{\mathbb{P}(\text{supp } \mu_{n-1} = \{\sigma^1, \sigma^2\})}{n} \mathbb{E} \left[\frac{\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2)}{\mu(\sigma^1)} \middle| Y_n = m, \text{supp } \mu_{n-1} = \{\sigma^1, \sigma^2\} \right]
\end{aligned}$$

Then, substituting in the following equalities:

$$\mathbb{P}(\text{supp } \mu_{n-1} = \{\sigma^1, \sigma^2\}) = \mu(\sigma^1) + \mu(\sigma^2),$$

and

$$\begin{aligned}
&\mathbb{E}[\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2) \mid Y_n = m, \text{supp } \mu_{n-1} = \{\sigma^1, \sigma^2\}] \\
&= \mathbb{E}[\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2) \mid \text{supp } \mu_{n-1} = \{\sigma^1, \sigma^2\}] \\
&= \frac{\mu(\sigma^1)}{\mu(\sigma^1) + \mu(\sigma^2)} \frac{\mu(\sigma^2)}{\mu(\sigma^1) + \mu(\sigma^2)}
\end{aligned}$$

we have

$$\mathbb{E} \left[\frac{\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2)}{\mu(\sigma^1)} \right] = \frac{\mu(\sigma^1) + \mu(\sigma^2)}{n \mu(\sigma^1)} \frac{\mu(\sigma^1)}{\mu(\sigma^1) + \mu(\sigma^2)} \frac{\mu(\sigma^2)}{\mu(\sigma^1) + \mu(\sigma^2)} = \frac{1}{n} \frac{\mu(\sigma^2)}{\mu(\sigma^1) + \mu(\sigma^2)}$$

which is precisely the kernel of the Glauber dynamics of two neighboring configurations. On the other hand, if $\|\sigma^1 - \sigma^2\| > 1$, then

$$\mathbb{E} \left[\frac{\mu_{n-1}(\sigma^1) \mu_{n-1}(\sigma^2)}{\mu(\sigma^1)} \right] = 0$$

as μ_{n-1} have fixed all but one coordinate. Thus, the dynamics associated with the coordinate is precisely the Glauber dynamics (we don't need to check the case $\sigma^1 = \sigma^2$ as they are kernels and so $K(\sigma, \sigma) = 1 - \sum_{\tilde{\sigma} \neq \sigma} K(\sigma, \tilde{\sigma})$).

4 Application: The KLS and Thin-Shell Conjecture

As an application of stochastic localizations, we will in this section introduce the KLS and thin-shell conjectures and by leveraging the linear-tilt localization, reduce the KLS conjecture (up to a logarithmic factor) into the thin-shell conjecture, i.e. we will describe a proof of theorem 4. The method presented in this section is due to Lee and Vempala [LV16] and reformulated in the language of concentration by Eldan [Eld18].

4.1 Concentration

Let us quickly introduce some preliminary definitions required to state the aforementioned conjectures.

Definition 4.1 (Concentration, [Eld18]). Let μ be a measure on \mathbb{R}^n , then μ is said to be C -(inversely)-concentrated if for all 1-Lipschitz function $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$,

$$\text{Var}_\mu[\phi] = \text{Var}_{X \sim \mu}[\phi(X)] \leq C^2. \quad (15)$$

We denote the least possible such C by C_{con}^μ .

Heuristically, the concentration measures the relation between μ and the Euclidean metric by providing a numerical control for the variance of its norm. This is perhaps best illustrated by the following proposition.

Proposition 4.1. Let X be a \mathbb{R}^n -valued random variable. Then for all K -Lipschitz function $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$,

$$\text{Var}[\phi(X)] \leq K^2 \text{Var}[\|X\|^2].$$

Proof. WLOG. by subtracting its expectation from X , we may assume $\mathbb{E}[X] = 0$. Let X' be a i.i.d. copy of X on the same probability space. Then for all K -Lipschitz function ϕ , we have

$$\begin{aligned} 2\text{Var}[\phi(X)] &= \text{Var}[\phi(X) - \phi(X')] && \text{(i.i.d.)} \\ &= \mathbb{E}[(\phi(X) - \phi(X'))^2] - \mathbb{E}[\phi(X) - \phi(X')]^2 \\ &= \mathbb{E}[(\phi(X) - \phi(X'))^2] && \text{(identically distributed)} \\ &\leq K^2 \mathbb{E}[\|X - X'\|^2] && \text{(as } \phi \text{ is } K\text{-Lipschitz)} \\ &= K^2 \mathbb{E}[X^T X + X'^T X' - X^T X' - X'^T X] \\ &= 2K^2 \text{Var}[\|X\|^2] - 2K^2 \text{Cov}(X, X') = 2K^2 \text{Var}[\|X\|^2]. && \text{(independence)} \end{aligned}$$

implying $\text{Var}[\phi(X)] \leq K^2 \text{Var}[\|X\|^2]$ as claimed. \square

With this proposition in mind, it is clear that for \mathbb{R} -valued random variables X , its law μ has concentration $C_{\text{con}}^\mu = \text{Var}[X]$. Furthermore, by considering the projection maps, it follows that the standard Gaussian measure on \mathbb{R}^n is 1-concentrated.

We note that the definition we are presenting here is slightly non-standard. However, utilising a remarkable result due to Milman, we show that this definition is equivalent to the following definitions in a specific sense.

Definition 4.2 (Exponential concentration, [Mil18]). Given a measure μ on \mathbb{R}^n , we say μ has exponential concentration if there exists some $c, D > 0$ such that for all 1-Lipschitz function $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$, $t > 0$, we have

$$\mu(|\phi - \mathbb{E}_\mu[\phi]| \geq t) \leq ce^{-Dt}. \quad (16)$$

Fixing $c = 1$, we denote the largest possible D as D_{exp}^μ .

Definition 4.3 (First-moment concentration, [Mil18]). Again, given μ a measure on \mathbb{R}^n , we say μ has first-moment concentration if there exists some $D > 0$ such that for all 1-Lipschitz function $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$, we have

$$\mathbb{E}_\mu[|\phi - \mathbb{E}_\mu[\phi]|] \leq \frac{1}{D}. \quad (17)$$

We denote the largest possible D by D_{FM}^μ .

It is clear that exponential concentration implies first-moment concentration. Indeed, if μ has exponential concentration with constant D (taking $c = 1$), then by the tail probability formula,

$$\mathbb{E}_\mu[|\phi - \mathbb{E}_\mu[\phi]|] = \int_0^\infty \mu(|\phi - \mathbb{E}_\mu[\phi]| \geq t) dt \leq \int_0^\infty e^{-Dt} dt = \frac{1}{D}.$$

On the other hand, Milman showed that for log-concave measures on \mathbb{R}^n , exponential concentration and first-moment concentration are equivalent in the following sense.

Theorem 3 (Milman, [Mil08]). For all log-concave measure μ on \mathbb{R}^n , μ has exponential concentration if and only if μ has first-moment concentration. Furthermore, $D_{\text{exp}}^\mu \simeq D_{\text{FM}}^\mu$ where we write $A \simeq B$ if there exists universal constants $C_1, C_2 > 0$ such that $C_1 A \leq B \leq C_2 A$.

With this theorem in mind, we establish the following correspondence.

Proposition 4.2. For all measures μ on \mathbb{R}^n , we have

Exponentially concentrated \implies Concentrated \implies First-moment concentrated

and $D_{\text{exp}}^\mu \leq \sqrt{2}(C_{\text{con}}^\mu)^{-1}$ and $(2C_{\text{con}}^\mu)^{-1} \leq D_{\text{FM}}^\mu$. Hence, if μ is log-concave, $D_{\text{exp}}^\mu \simeq D_{\text{FM}}^\mu \simeq (C_{\text{con}}^\mu)^{-1}$.

Proof. Assume first that μ is C -concentrated. Then by the Chebyshev inequality, we have

$$\mu(|\phi - \mathbb{E}_\mu[\phi]| \geq t) \leq \frac{1}{t^2} \text{Var}_\mu[\phi] \leq \frac{C^2}{t^2},$$

for all 1-Lipschitz ϕ . Thus, by tail probability,

$$\begin{aligned} \mathbb{E}_\mu[|\phi - \mathbb{E}_\mu[\phi]|] &= \int_0^\infty \mu(|\phi - \mathbb{E}_\mu[\phi]| \geq t) dt \\ &\leq \inf_{a>0} \left\{ \int_0^a \mu(|\phi - \mathbb{E}_\mu[\phi]| \geq t) dt + C^2 \int_a^\infty \frac{1}{t^2} dt \right\} \\ &\leq \inf_{a>0} \left\{ a + \frac{C^2}{a} \right\} = 2C, \end{aligned}$$

implying μ is first-moment concentrated with respect to the constant $(2C)^{-1}$.

On the other hand, if μ is exponential concentration with some constant D , then again by the tail probability,

$$\text{Var}_\mu[\phi] = \int_0^\infty \mu((\phi - \mathbb{E}_\mu[\phi])^2 \geq t) dt \leq \int_0^\infty e^{-D\sqrt{t}} dt = \frac{2}{D^2}$$

implying μ is $\sqrt{2}D^{-1}$ -concentrated. \square

4.2 Example: concentration of the Gaussian

4.3 The KLS and thin-shell conjecture

Informally, the KLS conjecture suggests that any log-concave measure on \mathbb{R}^n admits the same concentration as that of the Gaussian measure. However, unlike the Gaussian, as the concentration of measures is not invariant under linear functions, it is clear that the KLS conjecture would not hold without a suitable normalization. This leads us to the following formulation of the KLS conjecture.

Conjecture 1 (Kannan-Lovász-Simonovitz, [Eld18]). Denoting $\mathcal{M}_{\text{con}}^n$ the set of all log-concave probability measures μ on \mathbb{R}^n satisfying $\text{Var}_\mu[T] \leq 1$ for all 1-Lipschitz linear maps $T : \mathbb{R}^n \rightarrow \mathbb{R}$, there exists a *universal* constant C such that for all $\mu \in \mathcal{M}_{\text{con}}^n$, μ is C -concentrated.

We remark that C is universal in the sense that it does not depend on any parameter and in particular is independent of the dimension n .

Conjecture 2 (Thin-shell, [Eld13]). Taking $\mathcal{M}_{\text{con}}^n$ as above, there exists a universal constant C such that for all $\mu \in \mathcal{M}_{\text{con}}^n$, we have

$$\sqrt{\text{Var}_\mu[\|\cdot\|]} \leq C.$$

As the norm function is 1-Lipschitz, it is *a priori* that the thin-shell conjecture is weaker than that of the KLS conjecture. On the other hand, as we shall describe in the next section, as a consequence of the stochastic localization scheme, Eldan [Eld13] provides a reduction of the KLS conjecture to the thin-shell conjecture up to logarithmic factors.

Theorem 4 (Eldan, [Eld13]). Denoting $\mathcal{M}_{\text{con}}^n$ as above, we define

$$C_{\text{con}}^n := \inf \{C \mid \forall \mu \in \mathcal{M}_{\text{con}}^n, \mu \text{ is } C\text{-concentrated}\},$$

and

$$C_{\text{TS}}^n := \inf \left\{ C \mid \forall \mu \in \mathcal{M}_{\text{con}}^n, \sqrt{\text{Var}_\mu[\|\cdot\|]} \leq C \right\} = \sup_{\mu \in \mathcal{M}_{\text{con}}^n} \sqrt{\text{Var}_\mu[\|\cdot\|]},$$

we have,

$$C_{\text{TS}}^n \leq C_{\text{con}}^n \lesssim C_{\text{TS}}^n \log n.$$

The stochastic localization scheme has been wildly successful in making progress towards the KLS conjecture. Modifying the original arguments by Eldan, Lee and Vempala [LV16] obtained the bound $C_{\text{con}}^n \lesssim n^{-1/4}$. Further modifying their arguments, a recent breakthrough by Chen [Che20] improves the bound providing the following theorem.

Theorem 5 (Chen, [Che20]). $\log C_{\text{con}}^n \lesssim \sqrt{\log n \log \log n}$ and so $C_{\text{con}}^n = n^{-o(1)}$.

4.3.1 Equivalent formulation of the KLS conjecture

While we have formulated the KLS conjecture using the language of concentration, the conjecture itself is originally formulated as an isoperimetric problem. The isoperimetric problem is the problem in finding the set of unit volume with minimum surface area. In the case of the \mathbb{R}^n equipped with the Lebesgue measure, we have known since the ancient Greeks [B105] that the solution is the unit ball. With this in mind, it is natural for us to generalize the problem for arbitrary measures.

Definition 4.4 (Minkowski's boundary measure). Given a measure μ on \mathbb{R}^n and a Borel set $A \subseteq \mathbb{R}^n$, the Minkowski's boundary measure of A ,

$$\mu^+(\partial A) := \liminf_{\epsilon \downarrow 0} \frac{\mu(A_\epsilon) - \mu(A)}{\epsilon}.$$

where $A_\epsilon := \{x \in \mathbb{R}^n \mid \text{dist}(x, A) \leq \epsilon\}$ is the ϵ -thickening of some Borel set A .

The isoperimetric problem for the measure μ then becomes the problem of finding the set A satisfying $\mu(A) = 1$ with minimum $\mu^+(\partial A)$.

Definition 4.5 (Cheeger's inequality, [Mil08]). Given a measure μ on \mathbb{R}^n , we say μ satisfy Cheeger's inequality if there exists some D such that for all A ,

$$\mu(A) \wedge \mu(A^c) \leq D \mu^+(\partial A).$$

We call the largest such D the inverse Cheeger's constant (or the inverse isoperimetric constant) and denote it by D_C^μ .

With this, the KLS conjecture can be reformulated as the following.

Conjecture 3 (KLS, [Eld13]). Denoting $\mathcal{M}_{\text{iso}}^n$ the set of all log-concave and isotropic (i.e. $\mathbb{E}_{X \sim \mu}[X] = 0$ and $\text{Cov}_{X \sim \mu}(X) = \text{id}$) probability measures μ on \mathbb{R}^n , there exists a *universal* constant D such that for all $\mu \in \mathcal{M}_{\text{iso}}^n$, μ satisfy the Cheeger's inequality with constant D .

The equivalence of the reformulation follows by completing theorem 4.2 with two additional equivalences.

Theorem 6 (Milman, [Mil08]). For all log-concave measure μ on \mathbb{R}^n , the following are equivalent

- μ has exponential concentration with constant D_{exp}^μ .
- μ has first-moment concentration with constant D_{FM}^μ .
- μ satisfy the Cheeger's inequality with constant D_C^μ .
- μ satisfy the Poincaré inequality: there exists some $D > 0$ such that for all differentiable $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfying $\int \phi d\mu = 0$, we have

$$D \cdot \text{Var}_\mu[\phi] \leq \int \|\nabla \phi\|^2 d\mu.$$

We denote the largest such D by D_P^μ .

Furthermore, $D_{\text{exp}}^\mu \simeq D_{\text{FM}}^\mu \simeq D_{\text{p}}^\mu \simeq D_{\text{C}}^\mu$.

With this theorem and proposition 4.2 in mind, it is clear that the KLS conjecture can be instead formulated with any of these inequalities instead. Thus, the KLS conjecture can also be phrased using the constant provided by the Poincaré inequality.

Conjecture 4 (KLS, [Eld13]). Denoting $\mathcal{M}_{\text{iso}}^n$ as above, there exist a *universal* constant D such that for all $\mu \in \mathcal{M}_{\text{iso}}^n$, μ satisfy the Poincaré inequality with constant D .

We remark that isotropic measures satisfy the normalization condition in 1. Indeed, if $T : \mathbb{R}^n \rightarrow \mathbb{R}$ is a 1-Lipschitz linear function, i.e. is of the form $v \mapsto w^T v + d$ for some $w \in S^{n-1}$ and $d \in \mathbb{R}$, then we have

$$\text{Var}_\mu[T] = \text{Var}_{X \sim \mu} \left[\sum_{i=1}^n w_i X_i + d \right] = \sum_{i,j=1}^n w_i w_j \text{Cov}_{X \sim \mu}(X_i, X_j) = \sum_{i=1}^n w_i^2 = 1,$$

as $\text{Cov}_{X \sim \mu}(X) = \text{id}$.

4.4 Reduction of KLS to thin-shell

We will now present a proof of theorem 4. As a high level overview, recall that the linear-tilt localization of a given measure is a measure-valued martingale for which the original measure is recovered in the limit. Then, as the concentration of the measure relates to the covariance of said measure, we will stop the martingale before the covariance grows too large. This allows us to analyze the martingale in a more tractable manner. However, as the sequence is a martingale, some properties are invariant in time and hence allowing us to conclude that these properties also hold for the original measure.

We recall the goal of theorem 4 is to control $\text{Var}_\mu[\phi]$ by a logarithmic factor of $\text{Var}_\mu[\|\cdot\|]$. As translating the barycenter of μ does not affect its variance, we may assume μ has its barycenter $\bar{\mu}$ at the origin. Furthermore, we may assume μ is supported on $B_n(0) \subseteq \mathbb{R}^n$ with $B_n(0)$ the ball at the origin of radius n . Thus, we also have

$$\text{supp } \mu_t = \text{supp } F_t \mu \subseteq \text{supp } \mu \subseteq B_n(0)$$

for all $t > 0$.

Fix $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$ some 1-Lipschitz function and let (M_t) be the martingale as described above and in particular we recall equation (2.6) and so, $\text{Var}_\mu[\phi] = \text{Var}[M_\infty]$ where $M_t \xrightarrow{\text{a.e.}} M_\infty$. Hence, for all $t > 0$, by the martingale property we have

$$\begin{aligned} \text{Var}[M_t] + \mathbb{E}[\text{Var}[M_\infty | \mathcal{F}_t]] &= (\mathbb{E}[M_t^2] - \mathbb{E}[M_t]^2) + \mathbb{E}[\mathbb{E}[M_\infty^2 | \mathcal{F}_t] - \mathbb{E}[M_\infty | \mathcal{F}_t]^2] \\ &= \mathbb{E}[M_t^2] + (\mathbb{E}[\mathbb{E}[M_\infty^2 | \mathcal{F}_t]] - \mathbb{E}[M_t^2]) \\ &= \mathbb{E}[M_\infty^2] = \text{Var}[M_\infty], \end{aligned}$$

where the second equality follows as $\mathbb{E}[M_t] = \mathbb{E}[M_\infty] = \mathbb{E}_\mu[\phi] = 0$.

On the other hand, as (M_t) is a martingale, $M_t^2 - [M]_t$ is also a martingale implying $\mathbb{E}[M_t^2] = \mathbb{E}[M]_t$ and so $\text{Var}[M_t] = \mathbb{E}[M_t^2] - \mathbb{E}[M_t]^2 = \mathbb{E}[M]_t - \bar{\mu}^2 = \mathbb{E}[M]_t$. Hence, combining this with the above, we obtain the bound

$$\text{Var}_\mu[\phi] = \text{Var}[M_\infty] = \text{Var}[M_t] + \mathbb{E}[\text{Var}[M_\infty | \mathcal{F}_t]] = \mathbb{E}[M]_t + \mathbb{E}[\text{Var}_{\mu_t}[\phi]]. \quad (18)$$

explain

Now, observing that ϕ is 1-Lipschitz implies $\|\nabla\phi\|^2 \leq 1$, we have by lemma 2.4 the bound $\text{Var}_{\mu_t}[\phi] \leq t^{-1}$ (in fact $\text{Var}_{\mu_t}[\phi] \leq t^{-1} \wedge n^2$ as we have assumed $\text{supp } \mu_t \subseteq B_n(0)$). Thus, the second term $\mathbb{E}[\text{Var}_{\mu_t}[\phi]]$ is bounded by t^{-1} . With this in mind, by choosing an appropriate random time τ to stop the process such that $\mathbb{E}[M]_\tau$ is nicely bounded, the result follow by bounding $\mathbb{E}[\tau^{-1}]$. We dedicate the remainder of this section to describe said procedure in detail.

4.4.1 Differential of the quadratic variation

To bound the term $\mathbb{E}[M]_\tau$ we will compute its differential and bound it sufficiently such that we reobtain a bound for $[M]_\tau$ after integration. We will show $d[M]_t$ is bounded by a quantity concerning A_t . This should not be at all surprising as both $d[M]_t$ and A_t describes the variation of M_t in a infinitesimal time neighborhood of t .

We compute

$$\begin{aligned} dM_t &= d \int \phi(x) F_t(x) \mu_t(dx) = \int \phi(x) \langle x - a_t, dW_t \rangle \mu_t(dx) \\ &= \left\langle \int \phi(x)(x - a_t) \mu_t(dx), dW_t \right\rangle \end{aligned}$$

and so, by considering the component-wise quadratic variation, we have

$$d[M]_t = \left\| \int \phi(x)(x - a_t) \mu_t(dx) \right\|^2 dt. \quad (19)$$

Then, denoting θ the vector $\int \phi(x)(x - a_t) \mu_t(dx)$ normalized to have norm 1, so

$$\left\langle \theta, \int \phi(x)(x - a_t) \mu_t(dx) \right\rangle = \left\| \int \phi(x)(x - a_t) \mu_t(dx) \right\|$$

we observe,

$$\begin{aligned} d[M]_t &= \left\langle \theta, \int \phi(x)(x - a_t) \mu_t(dx) \right\rangle^2 dt = \left\langle \theta, \int \left(\phi(x) - \int \phi d\mu_t \right) (x - a_t) \mu_t(dx) \right\rangle^2 dt \\ &= \left(\int \left(\phi(x) - \int \phi d\mu_t \right) \langle \theta, x - a_t \rangle \mu_t(dx) \right)^2 dt \\ &\leq \left(\int \left(\phi(x) - \int \phi d\mu_t \right)^2 \mu_t(dx) \right) \left(\int \langle \theta, x - a_t \rangle^2 \mu_t(dx) \right) dt \\ &= \text{Var}_{\mu_t}[\phi] \left(\int \theta^T (x - a_t)^{\otimes 2} \theta \mu_t(dx) \right) dt = \text{Var}_{\mu_t}[\phi] (\theta^T A_t \theta) dt \\ &\leq \text{Var}_{\mu_t}[\phi] \|A_t\|_{\text{op}} dt. \end{aligned} \quad (20)$$

where the inequality follows by the Cauchy-Schwarz inequality and $\|\cdot\|_{\text{op}}$ denotes the operator norm. Thus, as we know $\text{Var}_{\mu_t}[\phi] \leq t^{-1}$, the problem is now reduced to that of bounding $\|A_t\|_{\text{op}}$.

4.4.2 Analysis of the covariance matrix

As demonstrated in section 2.1.1, we know the limiting behavior of the covariance matrices, namely $A_t \rightarrow 0$ point-wise as $t \rightarrow \infty$. This was important for us to establish the existence of the limit of (a_t) and (M_t) . However, as shown above, we now require some quantitative bounds for the operator norm of A_t . For this purpose, we first compute some useful properties of A_t .

Observing

$$\int dF_t(x)\mu(dx) = \int \langle x - a_t, dW_t \rangle \mu_t(dx) = \left\langle \int x\mu_t(dx) - a_t, dW_t \right\rangle = 0,$$

we have

$$\begin{aligned} da_t &= d \int x F_t(x) \mu(dx) = \int x dF_t(x) \mu(dx) = \int (x - a_t) dF_t(x) \mu(dx) \\ &= \int (x - a_t) \langle x - a_t, dW_t \rangle F_t(x) \mu(dx) = \int (x - a_t)^{\otimes 2} dW_t \mu_t(dx) = A_t dW_t \end{aligned} \quad (21)$$

where the second to last equality used the fact that $v \langle v, w \rangle = v^{\otimes 2} w$ for any appropriate v, w .

Similarly, computing using Itô's formula, we have

$$\begin{aligned} dA_t &= d \int (x - a_t)^{\otimes 2} F_t(x) \mu(dx) \\ &= \int (x - a_t)^{\otimes 2} dF_t(x) + F_t(x) d(x - a_t)^{\otimes 2} \\ &\quad - 2(x - a_t) \otimes d[a_t, F_t(x)]_t + F_t(x) d[a_t]_t \mu(dx). \end{aligned} \quad (22)$$

The second term vanishes as

$$\int F_t(x) d(x - a_t)^{\otimes 2} \mu(dx) = -2da_t \otimes \overbrace{\int (x - a_t) \mu_t(dx)}^{=0} = 0.$$

Also, by equation (21), $da_t = A_t dW_t$ implying $d[a_t]_t = A_t^2 dt$. Finally, as both (a_t) and $(F_t(x))$ are martingales, $d[a_t, F_t(x)]_t = F_t(x) A_t x dt$ and the third term becomes

$$\begin{aligned} -2 \int (x - a_t) \otimes d[a_t, F_t(x)] \mu_t(dx) &= -2A_t \left(\int (x - a_t) \otimes x \mu_t(dx) \right) dt \\ &= -2A_t \left(\overbrace{\int (x - a_t)^{\otimes 2} \mu_t(dx)}^{A_t} + \overbrace{\int (x - a_t) \mu_t(dx) \otimes a_t}^{=0} \right) dt \\ &= -2A_t^2 dt. \end{aligned}$$

Hence, combining these and equation (2) together in (22), we have

$$dA_t = \int (x - a_t)^{\otimes 2} \langle x - a_t, dW_t \rangle \mu_t(dx) - A_t^2 dt$$

However, since we wish to bound A_t from above, as the drift term $-A_t^2 dt$ only contributes negatively, an upper bound for the process of the form $\int (x - a_t)^{\otimes 2} \langle x - a_t, dW_t \rangle \mu_t(dx)$ is also sufficient for A_t . Hence, we proceed by ignoring the drift term and redefine the process A_t such that

$$dA_t = \int (x - a_t)^{\otimes 2} \langle x - a_t, dW_t \rangle \mu_t(dx). \quad (23)$$

With this justification, we now proceed to bound the operator norm of this new A_t . In particular, as A_t is symmetric, we recall that $\|A_t\|_{\text{op}} = \max_{i=1, \dots, n} \lambda_i(t) = \|(\lambda_i(t))_{i=1}^n\|_{\infty}$ where $\lambda_i(t)$ denotes the distinct eigenvalues of A_t . Hence, it suffices to find a bound for the potential

$$\Phi^\alpha(t) = \sum_{i=1}^n |\lambda_i(t)|^\alpha = \|(\lambda_i(t))_{i=1}^n\|_\alpha^\alpha \quad (24)$$

for some $\alpha > 0$. Furthermore, as A_t is positive semi-definite, $\lambda_i(t) \geq 0$ for all $i = 1, \dots, n$ and thus we have $\Phi^\alpha(t) = \sum_{i=1}^n \lambda_i(t)^\alpha$. Again, to proceed, we will attempt to compute $d\Phi^\alpha(t)$ at some $t = t_0 > 0$ utilizing the following lemma.

Lemma 4.3. If $A = [a_{ij}]$ is a diagonal matrix with distinct eigenvalues $\lambda_1, \dots, \lambda_n$, then for all $i, j, k, l, m \in 1, \dots, n$, we have

- $\frac{\partial \lambda_i}{\partial a_{jk}} = \delta_{ij} \delta_{ik}$;
- whenever $i \neq j$, $\frac{\partial^2 \lambda_i}{\partial a_{ij}^2} = 2(\lambda_i - \lambda_j)^{-1}$;
- and for $j \neq l, k \neq m$ or $i \neq j$ and $i \neq k$, $\frac{\partial^2 \lambda_i}{\partial a_{jk} \partial a_{lm}} = 0$,

where δ_{ij} denotes the Kronecker delta function.

As this lemma requires the matrix to be diagonal, denoting e_1, \dots, e_n as the normalized eigenbasis of A_{t_0} (they are in fact orthonormal as A_{t_0} is positive semi-definite), we will consider A_t with respect to this basis by considering the entries

$$a_{ij}(t) := \langle e_i, A_t e_j \rangle.$$

Using equation (23), we compute

$$\begin{aligned} da_{ij}(t) &= \left\langle e_i, \left(\int (x - a_t)^{\otimes 2} \langle x - a_t, dW_t \rangle \mu_t(dx) \right) e_j \right\rangle \\ &= \left\langle \int \langle e_i, (x - a_t)^{\otimes 2} e_j \rangle (x - a_t) \mu_t(dx), dW_t \right\rangle = \langle \xi_{ij}, dW_t \rangle \end{aligned}$$

where we introduce the notation $\xi_{ij} = \int \langle e_i, (x - a_t)^{\otimes 2} e_j \rangle (x - a_t) \mu_t(dx)$. Thus, combining this with lemma 4.3, denoting $\lambda_i = \lambda_i(t_0)$, we have by Itô's formula

$$\begin{aligned} d\lambda_i(t) &= \sum_{j,k=1}^n \frac{\partial \lambda_i}{\partial a_{jk}} da_{jk}(t) + \frac{1}{2} \sum_{j,k=1}^n \sum_{l,m=1}^n \frac{\partial^2 \lambda_i}{\partial a_{jk} \partial a_{lm}} d[a_{jk}, a_{lm}]_t \\ &= \langle \xi_{ii}, dW_t \rangle + \sum_{j \neq i} \frac{d[a_{ij}]_t}{\lambda_i - \lambda_j} = \langle \xi_{ii}, dW_t \rangle + \sum_{j \neq i} \frac{\|\xi_{ij}\|^2}{\lambda_i - \lambda_j} dt. \end{aligned} \quad (25)$$

at $t = t_0$. As a result, it is also clear that $d[\lambda_i(t)]_{t_0} = \|\xi_{ii}\|^2 dt$.

Again applying Itô's formula, we may finally compute

$$\begin{aligned}
d\Phi^\alpha(t) &= \sum_{i=1}^n \frac{\partial \Phi^\alpha}{\partial \lambda_i} \Big|_{t=t_0} d\lambda_i(t) + \frac{1}{2} \sum_{i,j=1}^n \frac{\partial^2 \Phi^\alpha}{\partial \lambda_i \partial \lambda_j} \Big|_{t=t_0} d[\lambda_i, \lambda_j]_t \\
&= \alpha \sum_{i=1}^n \lambda_i^{\alpha-1} d\lambda_i(t) + \frac{1}{2} \alpha(\alpha-1) \sum_{i=1}^n \lambda_i^{\alpha-2} d[\lambda_i(t)]_t \\
&= \alpha \sum_{i=1}^n \lambda_i^{\alpha-1} \left(\langle \xi_{ii}, dW_t \rangle + \sum_{j \neq i} \frac{\|\xi_{ij}\|^2}{\lambda_i - \lambda_j} dt \right) + \frac{1}{2} \alpha(\alpha-1) \sum_{i=1}^n \lambda_i^{\alpha-2} d[\lambda_i(t)]_t \\
&= \alpha \sum_{i \neq j} \lambda_i^{\alpha-1} \frac{\|\xi_{ij}\|^2}{\lambda_i - \lambda_j} dt + \frac{1}{2} \alpha(\alpha-1) \sum_{i=1}^n \lambda_i^{\alpha-2} \|\xi_{ii}\|^2 dt + \left\langle \underbrace{\alpha \sum_{i=1}^n \lambda_i^{\alpha-1} \xi_{ii}}_{=: v_t}, dW_t \right\rangle \\
&= \frac{1}{2} \alpha \sum_{i \neq j} \|\xi_{ij}\|^2 \frac{\lambda_i^{\alpha-1} - \lambda_j^{\alpha-1}}{\lambda_i - \lambda_j} dt + \frac{1}{2} \alpha(\alpha-1) \sum_{i=1}^n \lambda_i(t)^{\alpha-2} \|\xi_{ii}\|^2 dt + \langle v_t, dW_t \rangle \\
&\leq \frac{1}{2} \alpha(\alpha-1) \sum_{i \neq j} \|\xi_{ij}\|^2 (\lambda_i \vee \lambda_j)^{\alpha-2} dt + \frac{1}{2} \alpha(\alpha-1) \sum_{i=1}^n \lambda_i(t)^{\alpha-2} \|\xi_{ii}\|^2 dt + \langle v_t, dW_t \rangle \\
&= \frac{1}{2} \alpha(\alpha-1) \sum_{i,j=1}^n \|\xi_{ij}\|^2 (\lambda_i \vee \lambda_j)^{\alpha-2} dt + \langle v_t, dW_t \rangle \leq \alpha^2 \sum_{i,j=1}^n \|\xi_{ij}\|^2 \lambda_i^{\alpha-2} dt + \langle v_t, dW_t \rangle,
\end{aligned}$$

where the first inequality holds as

$$\frac{\lambda_i^{\alpha-1} - \lambda_j^{\alpha-1}}{\lambda_i - \lambda_j} = \lambda_i^{\alpha-2} + \lambda_i^{\alpha-3} \lambda_j + \cdots + \lambda_i^{\alpha-2} \leq (\alpha-1)(\lambda_i \vee \lambda_j)^{\alpha-2}.$$

Thus, we have shown

$$d\Phi^\alpha(t) \leq \alpha^2 \sum_{i=1}^n \lambda_i(t)^{\alpha-2} \sum_{j=1}^n \|\xi_{ij}\|^2 dt + \langle v_t, dW_t \rangle \quad (26)$$

where $v_t := \alpha \sum_{i=1}^n \lambda_i^{\alpha-1} \xi_{ii}$.

By recalling that our goal is to bound $\|A_t\|_{\text{op}}$ from above (c.f. equation (18) and (20)), we may assume without loss of generality that $\|A_t\|_{\text{op}} \geq 1$. Thus, applying the reverse Cauchy-Schwarz inequality to equation (26), we have

$$\begin{aligned}
d\Phi^\alpha(t) &\leq 2\alpha^2 \sum_{i=1}^n \lambda_i(t)^{\alpha-2} \sum_{j=1}^n \|\xi_{ij}\|^2 dt + \langle v_t, dW_t \rangle \\
&\leq 2\alpha^2 \|A_t\|_{\text{op}}^2 \sum_{i=1}^n \lambda_i(t)^{\alpha-2} \sum_{j=1}^n \|\xi_{ij}\|^2 dt + \langle v_t, dW_t \rangle \\
&\lesssim 2\alpha^2 \sum_{i=1}^n \lambda_i(t)^\alpha \sum_{j=1}^n \|\xi_{ij}\|^2 dt + \langle v_t, dW_t \rangle.
\end{aligned}$$

Thus, defining $K_t := \sup_i \sum_{j=1}^n \|\xi_{ij}\|^2$, we have the bound

$$d\Phi^\alpha(t) \lesssim 2\alpha^2 K_t \Phi^\alpha(t) dt + \langle v_t, dW_t \rangle. \quad (27)$$

4.4.3 Stopping the process early

As outlined in the beginning of this section, we will stop the process early in order to provide a bound for the right hand side of equation (18). By observing equation (20), we hypothesize that we should stop the process once $\|A_t\|_{\text{op}}$ grows too large. As a result we define the stopping time

$$\tau := \inf\{t > 0 \mid \|A_t\|_{\text{op}} > 2\} \wedge 1.$$

By the optional stopping theorem we have

$$\begin{aligned} [M]_\tau &= \int_0^\tau d[M]_t \leq \int_0^\tau \overbrace{\text{Var}_\mu[\phi]}^{\leq t^{-1} \wedge n^2} \overbrace{\|A_t\|_{\text{op}}}^{\leq 2} dt \\ &\leq 2 \int_0^\tau t^{-1} \wedge n^2 dt \leq 2 \int_0^1 t^{-1} \wedge n^2 dt = 2 + 4 \log n. \end{aligned}$$

Combining this with equation (18), we obtain

$$\text{Var}_\mu[\phi] \leq 2 + 4 \log n + \mathbb{E}[\tau^{-1}], \quad (28)$$

and it remains to find an upper bound for $\mathbb{E}[\tau^{-1}]$. Observing that $t < \tau$ whenever $\Phi^\alpha(t) < 2^\alpha$, we define the σ the first time for which the potential $\Phi^\alpha(t)$ reaches 2^α , namely

$$\sigma := \inf\{t > 0 \mid \Phi^\alpha(t) = 2^\alpha\},$$

we have $\sigma^{-1} \geq \tau^{-1}$ and so it suffices to bound σ from below.

For simplicity, let us ignore the stochastic term in equation (27) and regard it as an ODE. Then, by Gronwall's inequality, if we can find some constant K such that $K_t \leq K$ for all $t \leq \tau$, we have the bound

$$S_t \leq ne^{2\alpha^2 K t}.$$

Thus, substituting σ into the above, we have

$$2^\alpha = S_\sigma \leq ne^{2\alpha^2 K \sigma}$$

implying

$$\frac{\alpha \log 2 - \log n}{2\alpha^2 K} \leq \sigma \leq \tau.$$

Then, taking $\alpha = 10K \log n$, it is easy to check that

$$\frac{1}{10K \log n} \leq \frac{\alpha \log 2 - \log n}{2\alpha^2 K}$$

implying $\mathbb{E}[\tau^{-1}] \leq 10K \log n$. Of course, this deduction only holds while ignoring the stochastic term $\langle v_t, dW_t \rangle$. Nonetheless, this is justified as one can show that $\|v_t\|_2$ is bounded $\alpha \Phi^\alpha(t)$ and so the same analysis holds by applying the stochastic Gronwall's inequality (c.f. second part of lemma 34 in [LV18]).

Finally, to find a bound for (K_t) , we employ the following lemma.

Maybe not ignore the martingale term if we want a complete proof

Lemma 4.4 (Lemma 1.6 in [Eld13]). Denoting C_{TS}^n as in theorem 4, there exists a constant C such that for any log-concave, isotropic probability measure μ , we have

$$\sup_{\theta \in S^{n-1}} \sum_{i,j=1}^n \mathbb{E}_{X \sim \mu} [\langle X, e_i \rangle \langle X, e_j \rangle \langle X, \theta \rangle]^2 \leq C \sum_{k=1}^n \frac{(C_{\text{TS}}^n)^2}{k},$$

where $\{e_1, \dots, e_n\}$ is any orthonormal basis on \mathbb{R}^n .

Recalling that

$$\xi_{ij} = \mathbb{E}_{X+a_t \sim \mu_t} [\langle e_i, X^{\otimes 2} e_j \rangle X] = \mathbb{E}_{X+a_t \sim \mu_t} [\langle X, e_i \rangle \langle X, e_j \rangle X],$$

we have by Parseval's identity

$$\begin{aligned} K_t &= \sup_i \sum_{j=1}^n \|\xi_{ij}\|^2 = \sup_i \sum_{j=1}^n \|\mathbb{E}_{X+a_t \sim \mu_t} [\langle X, e_i \rangle \langle X, e_j \rangle X]\|^2 \\ &= \sup_i \sum_{j=1}^n \sum_{k=1}^n \langle \mathbb{E}_{X+a_t \sim \mu_t} [\langle X, e_i \rangle \langle X, e_j \rangle X], e_k \rangle^2 \\ &= \sup_i \sum_{j,k=1}^n \mathbb{E}_{X+a_t \sim \mu_t} [\langle X, e_i \rangle \langle X, e_j \rangle \langle X, e_k \rangle]^2 \\ &\leq \sup_{\theta \in S^{n-1}} \sum_{i,j=1}^n \mathbb{E}_{X+a_t \sim \mu} [\langle X, e_i \rangle \langle X, e_j \rangle \langle X, \theta \rangle]^2. \end{aligned}$$

We note that we cannot direct apply lemma 4.4 at this point since the measure μ_t might not be isotropic. Hence, to be able to use the lemma, we need to normalize the covariance of μ_t . Namely, taking $X + a_t \sim \mu_t$, we define $Y = A^{-1/2}X$ which by construction is isotropic. Thus, by observing that

$$\mathbb{E}_{X+a_t \sim \mu} [\langle X, e_i \rangle \langle X, e_j \rangle \langle X, \theta \rangle]^2 \leq \|A_t\|_{\text{op}}^3 \mathbb{E}_{X+a_t \sim \mu} [\langle Y, e_i \rangle \langle Y, e_j \rangle \langle Y, \theta \rangle]^2,$$

we have

$$\begin{aligned} K_t &\leq \sup_{\theta \in S^{n-1}} \sum_{i,j=1}^n \mathbb{E}_{X+a_t \sim \mu} [\langle X, e_i \rangle \langle X, e_j \rangle \langle X, \theta \rangle]^2 \\ &\leq \|A_t\|_{\text{op}}^3 \sup_{\theta \in S^{n-1}} \sum_{i,j=1}^n \mathbb{E}_{X+a_t \sim \mu} [\langle Y, e_i \rangle \langle Y, e_j \rangle \langle Y, \theta \rangle]^2 \leq 8C \sum_{k=1}^n \frac{(C_{\text{TS}}^n)^2}{k} \end{aligned} \tag{29}$$

where the last inequality follows as $\|A_t\|_{\text{op}} \leq 2$ for all $t < \tau$.

At last, combining equation (29) and (28), we have

$$\text{Var}_{\mu}[\phi] \leq 2 + \log n \left(4 + 80C \sum_{k=1}^n \frac{1}{k} (C_{\text{TS}}^n)^2 \right) = \Theta_n((C_{\text{TS}}^n \log n)^2)$$

implying there exists a constant $R > 0$ such that for all 1-Lipschitz ϕ , $\sqrt{\text{Var}_{\mu}[\phi]} \leq RC_{\text{TS}}^n \log n$, i.e. μ is $RC_{\text{TS}}^n \log n$ -concentrated and so, $C_{\text{con}}^n \leq RC_{\text{TS}}^n \log n$ as required.

References

- [ALG20] Nima Anari, Kuikui Liu, and Shayan Oveis Gharan. Spectral independence in high-dimensional expanders and applications to the hardcore model, 2020.
- [BL76] Herm Jan Brascamp and Elliott H Lieb. On extensions of the brunn-minkowski and prékopa-leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation. *Journal of Functional Analysis*, 22(4):366–389, 1976.
- [Bl05] Viktor Blåsjö. The isoperimetric problem. *The American Mathematical Monthly*, 112(6):526–566, 2005.
- [CE22] Yuansi Chen and Ronen Eldan. Localization schemes: A framework for proving mixing bounds for markov chains, 2022.
- [Che20] Yuansi Chen. An almost constant lower bound of the isoperimetric coefficient in the kls conjecture, 2020.
- [CLV20] Zongchen Chen, Kuikui Liu, and Eric Vigoda. Optimal mixing of glauber dynamics: Entropy factorization via high-dimensional expansion, 2020.
- [Eld13] Ronen Eldan. Thin shell implies spectral gap up to polylog via a stochastic localization scheme. *Geometric and Functional Analysis*, 23(2):532–569, mar 2013.
- [Eld18] Ronen Eldan. Lecture notes - from stochastic calculus to geometric inequalities, 2018.
- [Lev17] David Asher Levin. *Markov chains and mixing times / David A. Levin, Yuval Peres ; with contributions by Elizabeth L. Wilmer*. American Mathematical Society, second edition, 2017.
- [LV16] Yin Tat Lee and Santosh S. Vempala. Eldan’s stochastic localization and the kls conjecture: Isoperimetry, concentration and mixing, 2016.
- [LV18] Yin Tat Lee and Santosh S. Vempala. The kannan-lovász-simonovits conjecture, 2018.
- [Mil08] Emanuel Milman. Uniform tail-decay of lipschitz functions implies cheeger’s isoperimetric inequality under convexity assumptions. *Comptes Rendus Mathématique*, 346(17):989–994, 2008.
- [Mil18] Emanuel Milman. On the role of convexity in isoperimetry, spectral-gap and concentration. 2018.
- [Øks03] Bernt Øksendal. *Stochastic Differential Equations*, pages 65–84. Springer Berlin Heidelberg, Berlin, Heidelberg, 2003.