


Energy, Mass, Time, Space



To life.

Contents

License	vii
Preface	ix
I Foundations	1
1 The Complex Plane	2
2 The EMTS Framework: Core Definitions	4
2.1 The complex plane mapping	4
2.2 Polar representation	4
2.2.1 Physical interpretation of r and θ	5
2.2.2 The projection principle	5
2.3 Fundamental relationships	5
2.3.1 Time as phase	5
2.3.2 Energy-frequency connection	5
2.3.3 Measurement and collapse	6
2.4 Force quadrants	6
2.5 Notation conventions	6
2.6 Relationship to standard physics	6
2.7 Summary: The EMTS postulates	7
3 Polar Representation and Its Implications	8
3.1 Extended physical interpretation	8
4 Hermitian Operators on a Hilbert Space	9
5 The Quadrants and Their Implications	10

II	Framework and Mechanics	11
6	Standard Model	12
7	Mapping Reality	15
7.1	Reality as projection	15
7.2	Precision, dimensionality, and quantization	15
7.3	Quantization and the multiverse	16
7.3.1	Many-worlds from many-phases	16
7.3.2	Cross-branch interference	17
7.4	Implications for observers	17
7.5	Reconciling continuity and discreteness	17
7.6	The projection principle	18
8	Quantum Properties on the Complex Plane	19
8.1	Wave-particle duality as rotating projection	19
8.2	Superposition as geometric addition	19
8.3	Uncertainty as phase-projection complementarity	20
8.4	Quantization from closed-orbit conditions	20
8.5	Entanglement as correlated complex geometry	20
8.6	Summary	20
III	Implementation	21
9	Entanglement on the Complex Plane	22
9.1	Framing EMTS variables for entanglement	22
9.2	A minimal mathematical insertion	22
9.2.1	State, reduction, and θ -symmetry	22
9.2.2	Interaction kernels on the complex plane	22
9.3	Resonance as phase-locking in θ -time	23
9.4	Time independence and projection on the real axis	23
9.5	Testable consequences and a concrete ansatz	23
9.5.1	A simple EMTS entangled pair	23
9.5.2	Dynamics with resonance	23
9.6	Bridge to EMTS geometry	24
9.6.1	Core translation	24
9.6.2	Link to the geometry	24
10	Feynman Diagrams on the Complex Plane	25

10.1	Particles as trajectories in the plane	25
10.2	Vertices as junctions of complex vectors	25
10.3	Propagators as arcs or spirals	26
10.4	Curves and amplitudes	26
10.5	Loops and quantum corrections	26
10.6	Example: Electron–neutrino scattering	26
10.7	Why this is interesting	27
11	Periodic Table Analogy	28
11.1	How unknowns could emerge	28
11.1.1	Topology demands missing states	28
11.1.2	Symmetry completion	28
11.1.3	Forbidden gaps as clues	28
11.1.4	Density–phase resonance	28
11.2	What this could predict	28
11.3	How to search	29
12	Chemical Activation Analogue	30
12.1	Analogy	30
12.2	Activation mass–energy	30
12.3	Manipulating mass/energy	30
12.4	Diagrammatic representation	30
12.5	Formalization	31
13	Electron Angels and Demons	32
13.1	Motivation: the dual nature of the electron	32
13.2	Constructive interference: Angels	32
13.2.1	Definition	32
13.2.2	Physical content	33
13.2.3	EMTS geometry	33
13.3	Destructive interference: Demons	33
13.3.1	Definition	33
13.3.2	Electrons as plasmons	33
13.3.3	The Demon analogy	34
13.3.4	EMTS geometry	34
13.4	An EMTS taxonomy of collective electron modes	34
13.4.1	Angels: the natural order	35
13.4.2	Archangels: extreme coherence	35

13.4.3 Demons: the plasmon family	35
13.4.4 Devils: extreme destructive states	35
13.5 Dynamics: transitions between classes	36
13.6 A Pindaric Flight: Protons, Color, and the Strong Force	36
13.6.1 The proton as a collective state	36
13.6.2 The color-Devil condition	37
13.6.3 Why this is deeper than the electron Demon	37
13.6.4 Gluons as color Demons	37
13.6.5 Pions as two-body color Devils and the nuclear force	38
13.6.6 The quark-gluon plasma: melting the color Devil	38
13.6.7 A tentative EMTS conjecture	38
13.7 Summary: the celestial hierarchy in EMTS	39
IV Future Directions	40
14 Philosophy	41
15 Toward a Theory of Everything	42
15.1 State and geometry	42
15.2 Dynamics on spacetime and phase	42
15.3 Gauge sectors and interactions	42
15.4 Gravity via density-tied phase speed	43
15.5 Limiting cases and checks	43
15.6 Master equation with variable radius	43
16 Dark Matter in the Complex Plane	44
16.1 Off-axis or virtual trajectories	44
16.2 Phase-locked or resonant modes	44
16.3 High-radius, low-coupling orbits	45
16.4 Quadrant isolation	45
16.5 Density-driven phase speed modification	45
16.6 Missing geometric states	46
16.7 Composite interpretation	46
16.8 Testable predictions	46
16.8.1 Phase-independent gravitational lensing	46
16.8.2 Missing resonances in direct detection	46
16.8.3 Anomalous gravitational signatures	47
16.8.4 Indirect searches via phase transitions	47

<i>CONTENTS</i>	vi
16.9 Open questions	47
16.10 Connection to cosmology	47
Conclusion	48
Arguments and Notes	49

License

This work is licensed under the Creative Commons Attribution-ShareAlike 4.0 International (CC BY-SA 4.0).

SPDX-License-Identifier: CC-BY-SA-4.0

The full license text from the repository is reproduced below.

Creative Commons Attribution-ShareAlike 4.0 International (CC BY-SA 4.0)

SPDX-License-Identifier: CC-BY-SA-4.0

Copyright (c) 2025 Giancarlo Trevisan

This work is licensed under the Creative Commons Attribution-ShareAlike 4.0 International License.

You are free to:

- Share - copy and redistribute the material in any medium or format
- Adapt - remix, transform, and build upon the material for any purpose, even commercially

Under the following terms:

- Attribution - You must give appropriate credit, provide a link to the license, and indicate if changes were made.
- ShareAlike - If you remix, transform, or build upon the material, you must distribute your contributions under the same license as the original.

No additional restrictions - You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.

Full legal code: <https://creativecommons.org/licenses/by-sa/4.0/legalcode>

Human-readable summary: <https://creativecommons.org/licenses/by-sa/4.0/>

a. Adapted Material means material subject to Copyright and Similar Rights that is derived from or based upon the Licensed Material and in which the Licensed Material is translated, altered, arranged, transformed, or otherwise modified in a manner requiring permission under the Copyright and Similar Rights held by the Licensor. For purposes of this Public License, where the Licensed Material is a musical work, performance, or sound recording, Adapted Material is always produced where the Licensed Material is synched in timed relation with a moving image.

b. Adapter's License means the license You apply to Your Copyright and Similar Rights in Your contributions to Adapted Material in accordance with the terms and conditions of this Public License.

c. BY-SA Compatible License means a license listed at creativecommons.org/compatiblelicenses, approved by Creative Commons as essentially the equivalent of this Public License.

d. Copyright and Similar Rights means copyright and/or similar rights closely related to

copyright including, without limitation, performance, broadcast, sound recording, and Sui Generis Database Rights, without regard to how the rights are labeled or categorized. For purposes of this Public License, the rights specified in Section 2(b)(1)-(2) are not Copyright and Similar Rights.

e. Effective Technological Measures means those measures that, in the absence of proper authority, may not be circumvented under laws fulfilling obligations

Preface

In scientific discovery, pure mathematics often serves as the bedrock of our understanding of the universe. This essay is my attempt to develop, in real time with AI, an idea that has echoed in my mind for years. It is a perspective, not a declaration, and simply asks, *what if?*

Don't think it doesn't make sense; ask yourself what sense it could have.

First, a mathematical framework is defined that ties Energy, Mass, Time and Space (EMTS), then speculates how the framework could fit with our knowledge.

Giancarlo Trevisan

Visualize then Realize!

Part I

Foundations

Chapter 1

The Complex Plane

The connection between the complex plane and reality becomes especially significant when we consider the origins and applications of imaginary numbers. Historically, the introduction of imaginary numbers emerged from attempts to solve cubic equations, as first systematically explored by Gerolamo Cardano in the 16th century. Cardano discovered that even when all solutions to a cubic equation were real, the algebraic process sometimes required passing through intermediate steps involving the square roots of negative numbers—quantities that had no clear interpretation at the time.

This mathematical curiosity turned out to be much more than a formal trick. The development of complex numbers enabled mathematicians and physicists to describe phenomena that could not be captured by real numbers alone. A profound example is the discovery of electromagnetic waves. In the 19th century, James Clerk Maxwell formulated his famous equations, which describe how electric and magnetic fields propagate and interact. The solutions to Maxwell's equations are most naturally expressed using complex exponentials, where the imaginary unit i encodes oscillatory behavior—essential for describing wave phenomena such as light and radio waves (see Maxwell's equations[?]).

Thus, imaginary numbers are not merely abstract constructs; they are indispensable tools for modeling and understanding the physical world. Their use reveals that reality possesses layers and symmetries that extend beyond direct sensory perception, hinting at a deeper mathematical structure underlying the universe. The complex plane, therefore, is not just a mathematical convenience but a window into the hidden fabric of nature.

The complex plane, a mathematical construct that allows us to visualize complex numbers as points in a two-dimensional space. In this plane, the horizontal axis represents the real numbers, while the vertical axis represents the imaginary numbers. For our purposes, we will map mass onto the real axis and energy onto the imaginary axis. Why this choice? Well, mass seems more real than energy from my point of view (feel free to swap). With this picture in mind, let's pick a point z_0 on the complex plane and dive into its possible implications.




Figure 1.1: Complex plane illustration. m_z is what we perceive at time θ_0 and space r_0 .

Chapter 2

The EMTS Framework: Core Definitions

Having introduced the complex plane and its historical significance, we now establish the foundational elements of the Energy-Mass-Time-Space (EMTS) framework. This chapter consolidates the core mathematical definitions and physical interpretations that underpin the entire theory.

2.1 The complex plane mapping

The EMTS framework posits that physical states can be represented as points in the complex plane:

$$z = m + iE,$$

where:

- m is the mass component (mapped to the real axis)
- E is the energy component (mapped to the imaginary axis)
- $i = \sqrt{-1}$ is the imaginary unit

This choice reflects the intuition that mass is "more tangible" than energy in everyday experience, though the mapping could be reversed without loss of generality. What matters is the *relationship* between the components, not which axis they occupy.

2.2 Polar representation

Every complex point z can be expressed in polar form:

$$z = re^{i\theta} = r(\cos \theta + i \sin \theta),$$

where:

- $r = |z| = \sqrt{m^2 + E^2}$ is the **modulus** (unified magnitude)
- $\theta = \arg(z) = \arctan(E/m)$ is the **argument** (unified phase)

2.2.1 Physical interpretation of r and θ

The modulus r represents the total mass-energy content:

$$r = \sqrt{(mc^2)^2 + E^2} \quad \text{or in natural units} \quad r = \sqrt{m^2 + E^2}.$$

The argument θ encodes the **partition** between mass and energy:

$$m = r \cos \theta, \tag{2.1}$$

$$E = r \sin \theta. \tag{2.2}$$

Crucially, we identify θ with time evolution:

$$\theta(t) = \omega t,$$

where ω is the angular frequency related to energy by $E = \hbar\omega$ (Planck's quantum hypothesis).

2.2.2 The projection principle

Central postulate: Physical reality corresponds to the *projection* of the complex state onto the real axis:

$$\text{Observable reality} = \Re(z) = m = r \cos \theta.$$

This projection is **many-to-one**: infinitely many complex states

$$\{z_n = m + iE_n\}_n$$

all project to the same observable value m . The imaginary component E represents hidden degrees of freedom that influence dynamics but are not directly observed.

2.3 Fundamental relationships

2.3.1 Time as phase

The identification $\theta = \omega t$ unifies time evolution with rotation in the complex plane. A state evolves as:

$$z(t) = r e^{i\omega t}.$$

The observable projection oscillates:

$$m(t) = r \cos(\omega t),$$

explaining wave-like behavior from the geometry of rotation.

2.3.2 Energy-frequency connection

From quantum mechanics:

$$E = \hbar\omega \quad \Rightarrow \quad \omega = \frac{E}{\hbar}.$$

Thus the rotation rate in the complex plane is proportional to energy, linking dynamics to the imaginary component.

2.3.3 Measurement and collapse

In standard quantum mechanics, measurement "collapses" the wavefunction. In the EMTS framework:

1. Before measurement: Full state is $z = m + iE$ with complex dynamics
2. Measurement: Projects to $\Re(z) = m$
3. After measurement: The imaginary component E remains but is decohered from the observable m

Multiple branches with different E values persist along the imaginary axis (see Chapter 7 for the multiverse interpretation).

2.4 Force quadrants

The complex plane is divided into four quadrants based on the sign of m and E . We hypothesize that fundamental forces correspond to phase windows:

Quadrant	Force	Phase Range	Symbol
I	Electromagnetic	$0 < \theta < \frac{\pi}{2}$	Q_I
II	Weak	$\frac{\pi}{2} < \theta < \pi$	Q_{II}
III	Strong	$\pi < \theta < \frac{3\pi}{2}$	Q_{III}
IV	Gravitational	$\frac{3\pi}{2} < \theta < 2\pi$	Q_{IV}

Each force is characterized by a phase window $W_f(\theta)$ and coupling function $g_f(\theta, r)$. The observed interaction strength depends on where the state $z(t)$ resides in the phase cycle (see Chapter 5 for details).

2.5 Notation conventions

Throughout this work, we use the following notation:

- $z = m + iE = re^{i\theta}$ — Complex state (canonical form: $z = m + iE = re^{i\theta}$)
- $\theta = \omega t$ — Phase-time relation (shorthand: $\theta = \omega t$)
- $r = |z|$ — Modulus (total mass-energy)
- $\Re(z) = r \cos \theta = m$ — Real projection (observable)
- $\Psi(z, t)$ or $\Psi(r, \theta)$ — Wavefunction on complex state space
- $Q_I, Q_{II}, Q_{III}, Q_{IV}$ — Force quadrants (EM, Weak, Strong, Gravitational)

2.6 Relationship to standard physics

The EMTS framework is not a replacement for quantum mechanics or relativity, but a *geometric reinterpretation* that:

- Recovers the Schrödinger equation in appropriate limits
- Naturally incorporates special relativity through $r = \sqrt{m^2 + E^2}$
- Provides a unified language for discussing particles, forces, and spacetime
- Suggests extensions (e.g., multiverse structure, dark matter) that emerge from the geometry

Standard formulations are recovered by:

1. Restricting to the real axis (classical limit)
2. Quantizing the phase θ (quantum mechanics)
3. Coupling the modulus r to spacetime curvature (general relativity)

2.7 Summary: The EMTS postulates

We summarize the core framework:

1. **Complex representation:** Physical states are complex numbers $z = m + iE$
2. **Polar dynamics:** States evolve as $z(t) = re^{i\omega t}$ with $\omega = E/\hbar$
3. **Projection principle:** Observable reality is $\Re(z)$; imaginary components are hidden
4. **Phase-force mapping:** Forces correspond to phase windows in $[0, 2\pi)$
5. **Quantization:** Periodicity in θ leads to discrete spectra
6. **Dimensionality from precision:** Effective dimensionality increases with measurement precision

The remaining chapters develop these postulates in detail, exploring their implications for quantum mechanics (Part I), the Standard Model (Part II), specific phenomena (Part III), and future directions (Part IV).

Chapter 3

Polar Representation and Its Implications

Building on the core framework (Chapter 2), we explore deeper implications of the polar representation $z = m + iE = re^{i\theta}$. This chapter examines how the modulus r and argument θ relate to physical measurements and quantum phenomena.

3.1 Extended physical interpretation

The modulus and argument were defined in the core framework as:

$$r = \sqrt{m^2 + E^2}, \quad \theta = \arctan\left(\frac{E}{m}\right).$$

In units where $c = 1$, or more generally $r = \sqrt{(mc^2)^2 + E^2}$ and $\theta = \arctan(E/mc^2)$.

Space and Time. The key relation $\theta = \omega t$ with $E = \hbar\omega$ means probabilities or densities linked to projections oscillate with ω . The projection $m_z = \Re(z) = r \cos \theta$ can repeat with period 2π even as the underlying phase history differs.

Projection as Measurement. With the unit vector $|\psi\rangle = \cos \theta |m\rangle + i \sin \theta |E\rangle$, a mass measurement uses P_m and yields $\mathbb{P}(m) = \cos^2 \theta$, mirroring $m_z = r \cos \theta$. Standard treatments of such two-level mappings can be found in Griffiths[?].

Chapter 4

Hermitian Operators on a Hilbert Space

Let $H = \mathbb{C}^2$ with basis $\{|m\rangle, |E\rangle\}$. A normalized state is

$$|\psi\rangle = \cos\theta |m\rangle + i \sin\theta |E\rangle, \quad \langle\psi|\psi\rangle = 1.$$

Observables A are Hermitian with spectral decomposition $A = \sum_k a_k P_k$. Measurement of a_k occurs with probability $\langle\psi|P_k|\psi\rangle$ and the post-measurement state is $P_k|\psi\rangle/\sqrt{\langle\psi|P_k|\psi\rangle}$.

For a “mass” measurement, $P_m = |m\rangle\langle m|$, so $\mathbb{P}(m) = \cos^2\theta$, echoing the geometric projection $\Re(z) = r \cos\theta$.

Unitary evolution with $H = \frac{\hbar\omega}{2}\sigma_y$ yields $\mathbb{P}(m, t) = \cos^2(\omega t)$, bridging to $\theta = \omega t$. [?]

Chapter 5

The Quadrants and Their Implications

Let the pre-measurement state be $z(t_0) = re^{i\theta_0}$. A projection onto the real axis yields $m_z = r \cos \theta$. Hypothesis: the interaction channel observed in spacetime is tagged by the pre-collapse quadrant of $z(t_0)$.

- Quadrant I ($Q_I, 0 < \theta_0 < \frac{\pi}{2}$): Electromagnetic
- Quadrant II ($Q_{II}, \frac{\pi}{2} < \theta_0 < \pi$): Weak
- Quadrant III ($Q_{III}, \pi < \theta_0 < \frac{3\pi}{2}$): Strong
- Quadrant IV ($Q_{IV}, \frac{3\pi}{2} < \theta_0 < 2\pi$): Gravitational

With phase windows $W_f(\theta)$ and coupling $g(\theta, r)$, a cycle-averaged strength at radius r is

$$\bar{g}_f(r) = \frac{\omega(r)}{2\pi} \int_0^{2\pi} W_f(\theta) g(\theta, r) d\theta.$$

Part II

Framework and Mechanics

Chapter 6

Standard Model

Building on the EMTS framework (Chapter 2), we map Standard Model particles and forces onto the complex plane. In quantum theory, the state encodes probabilities.[?] In this framework, oscillations with frequency ω link to detection likelihoods. Particle vs. wave is reconciled: coherent evolution (*wave*) sets channel rates, while projection events (*particles*) are localized outcomes.

Particle vs. Wave

- **State vector:** carries phase and interferes.
- **Measurement:** projects to outcomes with Born probabilities.
- **Field view:** particles are quantized excitations; waves are coherent field amplitudes.

With $z(t) = re^{i\omega t}$ and windows $W_f(\theta)$ from Chapter 5, a minimal rate model is

$$R_f(r) = \frac{\omega}{2\pi} \int_0^{2\pi} W_f(\theta) g(\theta, r) d\theta.$$

Quarks in the complex plane

In the EMTS picture we write a generic excitation as $z = m + iE = re^{i\theta}$, with r encoding a space-like scale and $\theta = \omega t$ encoding time-like phase. Quarks can be viewed as excitations whose complex-plane behaviour is constrained by threefold structure:

- **Colour charge** corresponds to how the excitation winds in an internal copy of the complex plane, with three preferred phase orientations (“red, green, blue”) in an $SU(3)$ -like subspace.
- **Confinement** arises because a single quark’s complex trajectory cannot close in the observable m – E plane; only colour-neutral combinations lead to closed orbits and stable projections.
- **Flavour** (up, down, strange, etc.) is tied to distinct radii r and characteristic angular frequencies ω , leading to different effective masses and couplings.

In this view, quarks occupy specific bands in r and families of allowed phase patterns in θ , with hadrons emerging as composite closed paths whose joint projection appears as a single particle.

Leptons as simpler orbits

Leptons lack colour charge and so correspond to simpler trajectories on the same complex plane. Their key properties emerge from how their paths relate to the real and imaginary axes:

- **Charged leptons** (electron, muon, tau) follow orbits with a nonzero average real projection $\langle m_z \rangle$, giving rest mass, while their interaction with the electromagnetic field shifts θ and modulates E .
- **Neutrinos** are nearly lightlike: their trajectories lie close to the imaginary axis, with very small m but nontrivial phase evolution, which can support flavour oscillations as slow precessions between nearby orbits.
- **Generations** correspond to nested shells in r with similar angular patterns but different characteristic frequencies, reflecting the mass hierarchy without changing the basic complex geometry.

Leptons thus occupy cleaner, less composite regions of the complex plane, making them ideal probes of how m and E trade off along a single worldline.

Forces as phase symmetries

Forces in the Standard Model act by reshaping or constraining motion on the complex plane rather than by pushing in ordinary space alone. We can sketch them as follows:

- **Electromagnetism** tracks changes in the global phase of charged trajectories. A $U(1)$ gauge transformation is a shift $\theta \mapsto \theta + \alpha$, leaving r fixed but altering interference patterns and detection rates.
- **Weak interactions** mix components of z associated with different leptonic and quark flavours. Geometrically this can be pictured as rotations between nearby orbits in a multi-dimensional complex space, with massive gauge bosons mediating large, localized phase jumps.
- **Strong interactions** constrain colour phases so that only colour-neutral combinations yield closed, low-action paths. Gluon exchange continually rewires how individual quark trajectories share a joint complex-phase structure inside hadrons.

All three gauge forces can be summarised as rules about which deformations of the complex trajectory $z(t)$ leave physical rates invariant and which cost action, echoing the role of symmetries and gauge fields in the usual Standard Model.

A unified placement

In summary, the complex plane provides a common stage:

- **Quarks** occupy structured, colour-charged orbits whose composites form closed, observable paths.
- **Leptons** trace cleaner, single-particle trajectories distinguished mainly by radius and frequency.

- **Forces** appear as symmetries and constraints on allowed deformations of these complex paths.

This speculative mapping does not replace the field-theoretic Standard Model,[?] but it offers a geometric intuition: all fundamental matter and forces inhabit different patterns of motion and symmetry on an underlying complex m - E plane.

Chapter 7

Mapping Reality

7.1 Reality as projection

Recall from the core framework (Chapter 2) that physical states are represented as $z = m + iE = re^{i\theta}$, where r is the total mass-energy magnitude and θ is the phase encoding the partition between mass and energy components.

Reality emerges as the projection of the complex plane onto the real axis. For any complex state $z = a + ib$, what we observe—what we call "reality"—is the real component a . This projection is many-to-one: an infinite number of complex points

$$z_n = a + ib_n, \quad n \in \mathbb{Z} \text{ or } \mathbb{R},$$

all project to the same real value a . Each distinct imaginary component b_n represents a different hidden state, a different branch of possibility, yet all manifest identically in observable reality.

This degeneracy is profound: what appears as a single measurement outcome in our three-dimensional reality corresponds to an entire vertical line in the complex plane. The vast multiplicity of states "behind" each observation hints at the richness of the underlying structure.

7.2 Precision, dimensionality, and quantization

Conventional physics treats space and time as continuous manifolds with fixed dimensionality (3+1). The EMTS framework suggests a more subtle picture: **dimensionality is conveyed by precision.**

Consider a spatial coordinate x . At low precision (coarse-grained measurement), x appears one-dimensional. As we increase precision, we resolve finer structure:

- At scale Δx_1 , position is a single real number
- At scale $\Delta x_2 \ll \Delta x_1$, we resolve additional degrees of freedom: $x = x_{\text{coarse}} + \delta x$
- At Planck scale $\Delta x_P \sim 10^{-35}$ m, quantum geometry emerges, revealing discrete structure

The apparent dimensionality D_{eff} grows with precision ϵ :

$$D_{\text{eff}}(\epsilon) \sim D_0 + \alpha \log \left(\frac{L_{\text{max}}}{\epsilon} \right),$$

where D_0 is the macroscopic dimension, L_{\max} is the largest accessible scale, and α encodes how complexity unfolds at finer scales.

Similarly for time: what appears as a single temporal coordinate t at macroscopic precision becomes a rich structure at finer scales, potentially revealing branching, quantization, or phase-dependent flow rates (as encoded in $\mathcal{N}(x, \rho)$ from the unified framework).

7.3 Quantization and the multiverse

The projection degeneracy, combined with precision-dependent dimensionality, naturally leads to a **quantized multiverse structure**.

At each "point" in observable reality a , there exists a discrete (quantized) or continuous tower of states parameterized by the imaginary axis:

$$\{z_n = a + ib_n\}_n.$$

The quantization arises from the periodic structure in θ . Recall that physical states satisfy

$$\Psi(r, \theta + 2\pi) = \Psi(r, \theta),$$

leading to quantized phase modes:

$$\Psi_n(r, \theta) = R_n(r) e^{in\theta}, \quad n \in \mathbb{Z}.$$

Each integer n labels a different "branch" or "universe" characterized by:

- Winding number n around the origin in the complex plane
- Distinct energy spectrum from the Hamiltonian $\hat{H}_\theta = -\frac{\hbar^2}{2I_\theta} \partial_\theta^2 + U_{\text{quad}}(\theta)$
- Different coupling strengths to gauge forces via sector projectors $P_s(\theta)$

7.3.1 Many-worlds from many-phases

When a measurement occurs, the projection $z \rightarrow \text{Re}(z)$ collapses an infinite-dimensional phase space to a single real outcome. But the framework suggests that **all branches persist in the imaginary direction**. What appears as "wavefunction collapse" in conventional quantum mechanics is simply:

1. Projection: $z = a + ib \rightarrow a$ (observable outcome)
2. Persistence: The full state $z = a + ib$ continues to evolve
3. Branching: Different values of b (or equivalently θ) represent parallel realities, all projecting to the same a at the moment of measurement

The multiverse is not a set of disconnected universes, but rather a **continuum of phase-shifted realities**, all coexisting along the imaginary axis, distinguishable only by their internal phase structure θ , which determines:

- Which force quadrant dominates
- The rate of time flow via $\mathcal{N}(x, \rho, \theta)$
- Coupling to different gauge sectors
- Mass-energy partitioning via $\cos \theta$ and $\sin \theta$

7.3.2 Cross-branch interference

Although branches with different θ values project to the same real outcome, they can interfere quantum mechanically. The overlap integral

$$\mathcal{I}_{nm} = \int_0^{2\pi} d\theta \Psi_n^*(\theta) \hat{O} \Psi_m(\theta)$$

describes interference between branches n and m under observable \hat{O} . When $\mathcal{I}_{nm} \neq 0$, the branches are not fully independent—they constitute an entangled multiverse.

This framework recovers:

- **Decoherence:** Branches with $\Delta\theta \gg 1$ rapidly dephase, making $\mathcal{I}_{nm} \rightarrow 0$
- **Quantum superposition:** States with nearby θ values maintain coherence
- **Many-worlds interpretation:** Each θ -sector evolves independently when decoherence is complete

7.4 Implications for observers

An observer embedded in reality sees only the projection $\text{Re}(z)$. The observer's consciousness, measurement apparatus, and memory all exist in this projected space. Yet the full state $z = a + ib$ evolves according to the complex dynamics.

This creates an epistemic horizon: we can infer the existence of the imaginary component through:

1. **Quantum interference:** The imaginary part influences the evolution of the real part
2. **Entanglement:** Correlations between spatially separated real projections reveal hidden phase structure
3. **Gravitational effects:** The magnitude $r = \sqrt{a^2 + b^2}$ affects spacetime curvature, even though we only observe a
4. **Statistical ensembles:** Repeated measurements probe the distribution over b , revealing the underlying complex structure

7.5 Reconciling continuity and discreteness

The framework unifies apparently contradictory aspects of quantum mechanics and general relativity:

Aspect	Continuous	Discrete
Space	Manifold \mathcal{M}	Precision-dependent $D_{\text{eff}}(\epsilon)$
Time	Parameter t	Clock rate $\mathcal{N}(x, \rho)$, phase θ
Phase	Circle S^1	Quantized modes $e^{in\theta}$
Energy	Spectrum of \hat{H}	Eigenvalues E_n
Multiverse	Continuum of θ	Branches labeled by $n \in \mathbb{Z}$

At macroscopic scales and low precision, continuity dominates. At microscopic scales and high precision, discreteness emerges. The transition is smooth, governed by the characteristic scales \hbar , c , and the phase inertia I_θ .

7.6 The projection principle

We formalize the central insight:

Physical reality is the real projection of a complex state space. Observable phenomena correspond to $\text{Re}(z)$, while the imaginary component $\text{Im}(z)$ encodes hidden degrees of freedom—phase structure, interaction potentials, and alternate branches. Measurement selects a real outcome from an infinite multiplicity of complex pre-images, each representing a distinct universe in the quantized multiverse.

This projection principle explains:

- Why quantum mechanics is probabilistic (many z map to one a)
- Why entanglement is nonlocal (correlations exist in the imaginary direction)
- Why spacetime is dynamical (geometry responds to $|z|$, not just $\text{Re}(z)$)
- Why observers cannot access "other universes" directly (they live in the projection)
- Why precision matters (finer measurements probe finer structure in the complex plane)

Particles are points tracing oscillatory motion; forces emerge as structured relations across quadrants; spacetime geometry echoes these mathematical ties. But beneath it all lies a richer reality: a complex-valued ocean, of which we perceive only the surface.

Chapter 8

Quantum Properties on the Complex Plane

Quantum behaviour can be reframed in the EMTS picture established in Chapter 2. Recall that physical states are represented as $z = m + iE = re^{i\theta}$ with $\theta = \omega t$. Projection to the real axis ($r \cos \theta$) encodes what is classically observed, while the full complex motion remains hidden but dynamically essential.

8.1 Wave–particle duality as rotating projection

Take a single point $z(t) = re^{i\omega t}$. The real projection

$$m_z(t) = \Re z(t) = r \cos \theta(t)$$

oscillates, while discrete detection events correspond to sampling m_z at particular θ and locations in r . The “wave” is the smooth, complex rotation; the “particle” is the localized real-axis readout.

When several paths $z_k(t)$ are allowed, they add in the complex plane before projection:

$$z_{\text{tot}}(t) = \sum_k z_k(t).$$

Interference patterns arise because $\Re z_{\text{tot}}$ depends on relative angles θ_k , not just moduli r_k , mirroring standard complex-amplitude interference from part2-framework-mechanics/standard-model.tex.

8.2 Superposition as geometric addition

A superposed state of two alternatives A and B becomes a vector sum

$$z = \alpha z_A + \beta z_B,$$

with $\alpha, \beta \in \mathbb{C}$ encoding weights and phases. Only after projection does one obtain an outcome associated with z_A or z_B , analogous to

$$|\psi\rangle = \cos \theta |m\rangle + i \sin \theta |E\rangle$$

in part1-foundations/hermitian-operators.tex. The angle between z_A and z_B in the plane governs constructive or destructive interference in $\Re z$, giving a geometric visualization of the Born rule.

8.3 Uncertainty as phase–projection complementarity

If reality is read off as the real projection $m_z = r \cos \theta$, then sharp knowledge of m_z constrains θ to narrow windows where m_z takes that value. Conversely, if θ is highly delocalized over $[0, 2\pi)$, many different m_z values are sampled.

Formally, treat θ as an angle on a circle and its conjugate as an integer winding number n (counting how many 2π cycles the phase accumulates). Angle–number pairs satisfy an uncertainty relation of the form

$$\Delta n \Delta \theta \gtrsim \frac{1}{2},$$

which mirrors $\Delta E \Delta t$ once E is tied to angular frequency via $E = \hbar \omega$. In EMTS, the spread in θ (time-like phase) and the spread in m_z (measured mass-like projection) are thus inherently linked, explaining why precise localization in one degrades certainty in the other.

8.4 Quantization from closed-orbit conditions

Quantization appears naturally if allowed states correspond to closed or resonant trajectories on the complex circle. Requiring that after an evolution period T the point returns to itself,

$$\theta(T) - \theta(0) = 2\pi n, \quad n \in \mathbb{Z},$$

imposes discrete conditions on ω and hence on $E = \hbar \omega$. More generally, demanding single-valuedness of $\Psi(r, \theta)$ on S_θ^1 —as in the phase space discussed in part4-future/theory-of-everything.tex—forces integer winding numbers and a tower of allowed modes. The complex circle then acts as a geometric origin for energy levels and other quantized spectra.

8.5 Entanglement as correlated complex geometry

For two subsystems A, B with points $z_A = r_A e^{i\theta_A}$ and $z_B = r_B e^{i\theta_B}$, an entangled state can be represented by a joint amplitude $\Psi(z_A, z_B)$ on the product of two complex planes, as developed in part3-implementation/entanglement.tex. A simple form,

$$\Psi(z_A, z_B) \propto e^{im(\theta_A - \theta_B)},$$

locks their phase difference while leaving the common angle free. Measurements project each point separately to the real axis, but correlations in outcomes reflect the underlying constraint on (θ_A, θ_B) and hence on (m_{z_A}, m_{z_B}) .

In this view, entanglement is not mysterious action at a distance but a single geometric object on (z_A, z_B) space whose projections on separate real axes remain correlated even when r_A and r_B are widely separated.

8.6 Summary

Wave–particle duality, superposition, uncertainty, quantization, and entanglement all become features of how complex points move, add, and close on the EMTS plane, and how partial projections slice this richer geometry into the classical realities we observe.

Part III

Implementation

Chapter 9

Entanglement on the Complex Plane

9.1 Framing EMTS variables for entanglement

Recall from Chapter 2 that EMTS represents physical events as points on the complex plane $z = m + iE = re^{i\theta}$, with r encoding space and θ encoding time. Two-system entanglement can be modeled on pairs (z_A, z_B) by building joint states and correlators that respect EMTS' polar structure. In this view, time has two aspects: a monotone history variable (global θ -translation) and a periodic phase (modulo 2π), which naturally invites resonance phenomena and Floquet-like behavior in θ space.

9.2 A minimal mathematical insertion

9.2.1 State, reduction, and θ -symmetry

- **Global state:** Let $|\Psi\rangle$ live on a Hilbert space over EMTS points; for two subsystems A, B at $z_A = r_A e^{i\theta_A}$, $z_B = r_B e^{i\theta_B}$, write the joint amplitude $\Psi(z_A, z_B)$.^[?]
- **Entanglement test:** Compute $\rho_A = \text{Tr}_B |\Psi\rangle\langle\Psi|$ and a measure $S_A = -\text{Tr}(\rho_A \log \rho_A)$ (or a negativity). Entanglement is present iff ρ_A is mixed.
- **θ -translation invariance:** If dynamics and initial conditions are invariant under simultaneous shifts $\theta_A \rightarrow \theta_A + \alpha$, $\theta_B \rightarrow \theta_B + \alpha$, then any entanglement measure depends only on the phase difference $\Delta\theta = \theta_A - \theta_B$ and the radii (r_A, r_B) , not on absolute θ (stationarity):

$$\partial_\Theta S_A = 0, \quad \Theta = \frac{1}{2}(\theta_A + \theta_B).$$

9.2.2 Interaction kernels on the complex plane

- **Phase-sensitive coupling:**

$$H_{\text{int}} = \lambda f(r_A, r_B) \cos(\Delta\theta - \phi_0) \hat{O}_A \otimes \hat{O}_B,$$

which entangles A and B when $f \neq 0$. The $\cos(\Delta\theta)$ factor makes phase relations explicit; ϕ_0 sets a preferred phase alignment.

- **Holomorphic form (optional):**

$$H_{\text{int}} = \lambda g(z_A, z_B) \hat{O}_A \otimes \hat{O}_B + \text{h.c.},$$

with g holomorphic to ensure Cauchy–Riemann compatibility in EMTS. This yields entanglement protected along contours of constant argument or modulus depending on g .

9.3 Resonance as phase-locking in θ -time

- **Phase-locked entanglement:** If time has a periodic component, entanglement can strobe at resonant phase differences. With a drive of frequency ω on θ , stroboscopic evolution produces

$$U_F = e^{-iH_{\text{eff}}T}, \quad T = \frac{2\pi}{\omega},$$

and entanglement peaks when $\Delta\theta$ satisfies locking conditions (e.g., $\Delta\theta \approx \phi_0 \bmod 2\pi$).

- **Growth vs. invariance:** In generic (chaotic) dynamics, entanglement exhibits universal growth and saturation patterns (e.g., area-law to volume-law crossover with velocity v_E). “Entanglement is independent of time” can be realized when the initial state and generator are θ -stationary so only $\Delta\theta$ matters, or when a resonance creates steady phase-locking so the entanglement measure becomes θ -periodic and effectively constant under coarse graining.

9.4 Time independence and projection on the real axis

- **Projection choice matters:** If “projected on the real axis” means evaluating observables at fixed θ (equal-time slice), entanglement reduces to a function of radii and their separation along r , i.e., $S_A = S_A(r_A, r_B, \Delta\theta)$ with $\Delta\theta = 0$. With θ -translation invariance, this yields entanglement profiles depending only on spatial relations in r .
- **Integrating out θ :** Alternatively, integrating phases (or averaging over Θ) leaves phase-invariant correlators:

$$\overline{C}(r_A, r_B) = \frac{1}{2\pi} \int_0^{2\pi} d\Theta \, C(r_A e^{i(\Theta+\Delta\theta/2)}, r_B e^{i(\Theta-\Delta\theta/2)}).$$

9.5 Testable consequences and a concrete ansatz

9.5.1 A simple EMTS entangled pair

$$\Psi(z_A, z_B) = \frac{1}{\sqrt{2}} \left[\phi_0(r_A) \phi_1(r_B) e^{im(\theta_A - \theta_B)} + \phi_1(r_A) \phi_0(r_B) e^{-im(\theta_A - \theta_B)} \right].$$

- **Label:** m is a winding in $\Delta\theta$; $\phi_{0,1}$ control localization in r .
- **Property:** Entanglement is maximal and independent of Θ ; tuning m sets resonance channels in $\Delta\theta$.

9.5.2 Dynamics with resonance

Use $H_{\text{int}}(t) = \lambda(t) f(r_A, r_B) \cos(\Delta\theta - \phi_0) \hat{O}_A \otimes \hat{O}_B$ with $\lambda(t+T) = \lambda(t)$. Predict:

- **Locking:** Stable entanglement plateaus at $\Delta\theta \approx \phi_0$.
- **Velocity bounds:** Expect entanglement growth bounded by an effective v_E and shaped by a “line tension,” analogous to results in Floquet and chaotic circuits.
- **Equal- θ slices:** On $\Delta\theta = 0$, entanglement reduces to spatial profiles along r .

9.6 Bridge to EMTS geometry

9.6.1 Core translation

Any normalized two-level state can be written

$$|\psi\rangle = \cos\theta |m\rangle + i \sin\theta |E\rangle,$$

so the geometric angle θ from $z = re^{i\theta}$ becomes the parameter in the quantum superposition. Measurements use projectors like $P_m = |m\rangle\langle m|$ giving $P(m) = \cos^2\theta$. A simple Hamiltonian $H = (\hbar\omega/2)\sigma_y$ rotates probability between $|m\rangle$ and $|E\rangle$ at angular frequency ω .^[?]

9.6.2 Link to the geometry

The real-axis projection in the complex picture becomes applying P_m in Hilbert space; “collapse” corresponds to projecting z to $\text{Re}(z)$ and normalizing. The radial coordinate r controls overall scale but is factored out in quantum normalization.

Chapter 10

Feynman Diagrams on the Complex Plane

Let’s reinterpret Feynman diagrams within the EMTS framework (Chapter 2). Instead of drawing them in the usual spacetime coordinates, plot each particle’s state $z = m + iE = re^{i\theta}$ in the mass–energy plane. Each line in a diagram becomes a curve

$$\gamma(\lambda) : [0, 1] \rightarrow \mathbb{C}, \quad \gamma(\lambda) = r(\lambda)e^{i\theta(\lambda)},$$

with λ a path parameter. External legs are *open* curves, anchored to asymptotic “in” and “out” states, while internal lines and loops can form *closed* or self-intersecting curves that encode virtual processes.

10.1 Particles as trajectories in the plane

- **External legs:** Each external particle is an open curve in the complex plane from an initial z_i to a final z_f , representing how its mass–energy configuration evolves between preparation and detection.
- **Massive vs. massless:** Massless particles (photons, gluons) follow paths hugging the imaginary axis; massive particles trace tilted paths with significant real component.
- **Neutrinos:** Almost vertical lines near the imaginary axis, with tiny real offset.
- **Quarks:** Confined loops in the strong-force quadrant, never escaping to free-particle regions.

In this picture, a conventional Feynman diagram is not just a graph of lines and vertices but a collection of curves $\{\gamma_i\}$ drawn on the same complex plane, meeting at junctions that enforce complex conservation laws.

10.2 Vertices as junctions of complex vectors

A vertex is a point where several z -vectors meet, and complex-vector conservation applies:

$$\sum_{\text{in}} z = \sum_{\text{out}} z.$$

The quadrant in which the vertex sits indicates the mediating force (see Chapter 5: Q_I for EM, Q_{II} for weak, Q_{III} for strong, Q_{IV} for gravitational).

10.3 Propagators as arcs or spirals

- **Massive propagator:** Spiral with both radial and angular change (space and time evolution).
- **Massless propagator:** Pure angular advance at fixed radius (lightlike).
- **Virtual particles:** Paths that wander into “unphysical” quadrants or cross branch cuts; their projection on the real axis may be zero or negative.

Each propagator thus corresponds to a family of admissible curves between two points z_a and z_b . In a path-integral spirit, the physical amplitude weights these curves by a phase factor depending on the “action” along the path in the complex plane.[?]

10.4 Curves and amplitudes

For a given scattering process, standard quantum field theory assigns an amplitude by summing over all compatible Feynman diagrams. In the complex-plane view, this becomes a sum over classes of curves:

$$\mathcal{A}_{\text{process}} \sim \sum_{\text{diagrams}} \sum_{\{\gamma_i\}} e^{iS[\{\gamma_i\}]},$$

where the action functional $S[\{\gamma_i\}]$ depends on how the curves wind, which quadrants they traverse, and how they meet at vertices. Open curves carry the quantum numbers of external particles, while closed curves (loops) encode vacuum fluctuations and radiative corrections.

In this hypothesis, momentum conservation and on/off-shell conditions translate into geometric constraints on allowed curve shapes and endpoints in the m – E plane. Different diagram topologies then correspond to different homotopy classes of curve-collections, offering a topological handle on selection rules and interference.

10.5 Loops and quantum corrections

Loop diagrams in QFT become closed loops in the complex plane.[?] The winding number of the loop can correspond to a conserved quantum number (charge, baryon number). Divergences may appear as loops that shrink toward the origin $z = 0$, requiring renormalization as a deformation away from the singularity.

10.6 Example: Electron–neutrino scattering

- **Initial state:** Electron z_e in Q4 (positive mass, negative binding energy), neutrino z_ν near imaginary axis.
- **Vertex:** Weak-force quadrant (Q2), where a W boson is exchanged.
- **Propagator:** W boson path arcs from electron’s z to neutrino’s z , crossing the imaginary axis.
- **Final state:** New z positions for outgoing electron and neutrino, conserving the complex sum.

10.7 Why this is interesting

- **Unified conservation:** Mass and energy conservation become one complex equation at each vertex.
- **Virtuality geometry:** Off-shell particles are “off-axis.”
- **Topological insight:** Winding numbers and quadrant crossings give visual handles on selection rules and forbidden processes.

Chapter 11

Periodic Table Analogy

Could the EMTS framework play a role similar to the periodic table in hinting at missing pieces? The periodic table worked because its arrangement was a geometry of relationships; gaps were structural necessities. The complex-plane paths and diagram reinterpretations have similar potential if the geometry demands certain configurations.

11.1 How unknowns could emerge

11.1.1 Topology demands missing states

If winding numbers, quadrant crossings, or density-dependent phase rules are strict, certain interaction vertices only balance if a missing z -vector exists. That vector could correspond to an undiscovered particle, a new interaction, or a composite state.

11.1.2 Symmetry completion

If the diagram set respects a symmetry (rotational in θ , reflection across axes), incomplete multiplets stand out—like polygons in the complex plane with a missing vertex.

11.1.3 Forbidden gaps as clues

Absence of a path can be telling: if quadrant transitions are allowed by geometry but never observed, that could indicate a hidden law or a very heavy/weakly coupled particle.

11.1.4 Density–phase resonance

With $\omega(\rho)$ tied to density, there may be resonant densities where paths close neatly in θ after an integer number of 2π cycles. Gaps in a resonance sequence suggest missing states.

11.2 What this could predict

New neutrino-like states, exotic hadrons, force carriers (dark photon), or leptoquarks could fill geometric gaps implied by closure rules and symmetries.

11.3 How to search

Catalogue known particles in (m, E, r, θ) , map interaction paths, look for incomplete geometric patterns, and infer the missing z : its quadrant, radius, and phase give mass, energy, and coupling hints.

Chapter 12

Chemical Activation Analogue

This section borrows the logic of chemical activation diagrams and transplants it into the mass–energy–phase plane.

12.1 Analogy

In chemistry, an activation diagram plots potential energy vs. reaction coordinate. Here, the “reaction coordinate” is a path in (r, θ) : initial $z_i = r_i e^{i\theta_i}$, final $z_f = r_f e^{i\theta_f}$, and a barrier as a dynamically disfavoured region.

12.2 Activation mass–energy

The activation energy becomes an activation mass–energy: the extra $|z|$ or angular displacement needed to connect two states.

12.3 Manipulating mass/energy

- **Catalysis analogue:** Introduce an intermediate path bending through a quadrant with a lower barrier; mediators or fields change the allowed trajectory so the peak $|z|$ is smaller.
- **Phase-assisted transitions:** Because θ is cyclic, one can wrap around instead of going straight over, akin to tunnelling.
- **Density-tuned activation:** If $\omega(\rho, r)$ changes effective heights, altering local density can lower the barrier via phase-resonance.

12.4 Diagrammatic representation

Plot total $|z| = r$ vs. path length along (r, θ) ; different complex-plane paths yield different activation curves.

12.5 Formalization

Define an activation functional for a path γ :

$$\mathcal{A}[\gamma] = \max_{s \in \gamma} [r(s) - \min(r_i, r_f)],$$

and search for paths minimizing \mathcal{A} subject to complex-plane conservation laws. Catalysts, fields, or density changes are deformations of $U(r, \theta)$ that reduce \mathcal{A} .

Chapter 13

Electron Angels and Demons

13.1 Motivation: the dual nature of the electron

The electron is simultaneously a particle and a wave. As a particle it carries a well-defined mass m_e and charge $-e$; as a wave it is described by a complex-valued wavefunction $\psi(\mathbf{x}, t) = |\psi| e^{i\phi}$ that obeys the Schrödinger (or Dirac) equation. When many-electron wavefunctions overlap, the phases ϕ_k can align or oppose, producing entirely new macroscopic states that bear little resemblance to the bare electron.

Within the EMTS framework the electron sits at

$$z_e = m_e + iE_e = r_e e^{i\theta_e},$$

where $r_e = \sqrt{m_e^2 + E_e^2}$ is the total mass-energy magnitude and $\theta_e = \arctan(E_e/m_e)$ records the partition between rest mass and kinetic/potential energy. For a free electron at rest $\theta_e \approx 0$ (real axis), while a high-energy electron is rotated toward the imaginary axis.

A *collective* state of N electrons is then described by the vector sum

$$Z_{\text{coll}} = \sum_{k=1}^N z_k = \sum_{k=1}^N r_k e^{i\theta_k}.$$

The character of this sum — constructive or destructive — determines whether the system behaves as an **Angel** or a **Demon** in the taxonomy developed below.

13.2 Constructive interference: Angels

13.2.1 Definition

Consider N electrons whose EMTS phases are *aligned*:

$$\theta_k \approx \bar{\theta} \quad \forall k.$$

The collective vector is then approximately

$$Z_{\text{Angel}} \approx \left(\sum_{k=1}^N r_k \right) e^{i\bar{\theta}},$$

which is a single EMTS point with modulus $\sim N r_e$ and essentially the same argument as the individual electrons. The real (mass-like) and imaginary (energy-like) components both scale up together.

13.2.2 Physical content

Phase alignment means that maxima of the individual wavefunctions coincide. Some familiar realizations:

- **Cooper pairs and superconductivity.** Below T_c , electrons near the Fermi level pair into Cooper pairs with equal and opposite momenta and opposite spins; their pair wavefunction is a single coherent superposition. In EMTS, the pair amplitude has its phases locked so that Z_{pair} sits solidly on the real axis (dominant mass, minimal energy dispersion), explaining the gap and lossless transport.[?]
- **Coherent many-body ground states.** In a crystal, Bloch electrons form bands; near a filled lower band the coherent sum of occupied states gives a stable, low-energy configuration — the background “sea” of condensed matter.
- **Stimulated emission.** In a laser medium, photon-driven transitions reinforce phase alignment of atomic dipoles; the emitted photons are the EM analogue of the Angel state.

13.2.3 EMTS geometry

In the complex plane, an Angel state corresponds to a tight *cluster* of N points near a single z_e , their vector sum pointing radially outward with length $\sim Nr_e$. The projection $\Re(Z_{\text{Angel}}) = Nr_e \cos \bar{\theta}$ is large: Angels are *observable* in the real (mass/matter) sector.

13.3 Destructive interference: Demons

13.3.1 Definition

Now suppose the phases of the participating electrons are *anti-aligned*:

$$\theta_k = \bar{\theta} + \delta_k, \quad \sum_{k=1}^N e^{i\delta_k} \approx 0.$$

In the simplest two-electron toy model with $\theta_1 = \theta$ and $\theta_2 = \theta + \pi$:

$$Z_{\text{Demon}} = r_1 e^{i\theta} + r_2 e^{i(\theta+\pi)} = (r_1 - r_2) e^{i\theta}.$$

When $r_1 = r_2$ the individual mass-energy contributions *cancel exactly*: $Z_{\text{Demon}} = 0$. Even when cancellation is only partial, the dominant real-axis component is suppressed, and what survives is primarily imaginary — *energy without mass*.

13.3.2 Electrons as plasmons

The most experimentally dramatic Demon is the *plasmon* — the collective charge-density oscillation of the electron gas.[?] A classical derivation illustrates the Demon mechanism:

1. Each conduction electron in a metal has individual EMTS vector $z_k = m_e + iE_k$.
2. A charge displacement excites a density wave $n(\mathbf{x}, t) = n_0 + \delta n \cos(\mathbf{q} \cdot \mathbf{x} - \omega t)$.

3. The restoring Coulomb forces couple all electrons so that individual particle identities dissolve into a collective mode at the plasma frequency

$$\omega_p = \sqrt{\frac{n_0 e^2}{\epsilon_0 m_e}}.$$

4. This new quasi-particle — the *plasmon* — has an effective mass determined by ω_p and a dispersion $\omega(\mathbf{q})$ very different from the bare electron.

In EMTS, the plasmon is described not by any single z_k but by the *residual* of the collective sum after destructive cancellation. Its EMTS vector sits near the imaginary axis (large θ , small real component), reflecting the fact that the plasmon has very little “matter-like” character — it is primarily an *energy excitation* propagating through the electron sea.

13.3.3 The Demon analogy

The metaphor of the *Demon* is apt:

- Like a demon, the plasmon exploits the absence of order (destructive phase cancellation) to manifest as something qualitatively different from the individual electrons.
- It “hides” behind the anti-phase interference — invisible to direct mass measurement, yet capable of transporting energy at ω_p and mediating optical phenomena at UV frequencies.
- Demons are *unnatural* in the sense that they require a specific cancellation to persist; they depend on the maintenance of phase opposition across the participating electrons.

13.3.4 EMTS geometry

In the complex plane, the individual electron points spread symmetrically around the origin so that their vector sum Z_{Demon} is close to (or exactly at) zero. The plasmon is then the *phase-space excitation* of this near-cancellation — a small residual that rotates rapidly near the imaginary axis, with $|\Re(Z)| \ll |\Im(Z)|$.

13.4 An EMTS taxonomy of collective electron modes

The constructive/destructive axis is not binary. We can organize all collective electron states on a spectrum according to the degree of phase coherence in the EMTS sum. Four named classes arise naturally, analogous to a celestial hierarchy:

Name	Phase relation	$ Z_{\text{coll}} /Nr_e$	Physical archetype
Archangel	Extreme constructive, $\delta_k \rightarrow 0$	$\rightarrow 1$	BEC, superfluid, lasing mode
Angel	Constructive, partial alignment	0.5 to 1	Cooper pairs, band ground state
Demon	Destructive, partial cancellation	$\lesssim 0.1$	Plasmons, acoustic modes
Devil	Extreme destructive, near-total cancel	$\rightarrow 0$	Acoustic plasmon, demon quasi-particle

13.4.1 Angels: the natural order

Constructive interference is the *ground-state tendency*: a system of electrons minimizes its energy by occupying coherently the lowest available modes, with wavefunctions adding positively. Angels represent equilibrium. Left to evolve without perturbation, an electron ensemble relaxes toward an Angelic configuration because such states minimize free energy.

13.4.2 Archangels: extreme coherence

When phase alignment is *forced* across a macroscopic number of particles — as in a Bose-Einstein condensate (BEC), a superfluid, or an optical laser — the collective EMTS vector points almost perfectly along a single direction in the complex plane, with $|Z| \approx Nr_e$. This is the Archangel limit:

$$Z_{\text{Arch}} = Nr_e e^{i\bar{\theta}}, \quad \text{all } \theta_k \equiv \bar{\theta}.$$

Archangels are extraordinary: they require macroscopic phase locking, typically enforced by a symmetry-breaking mechanism (BCS gap equation, Gross-Pitaevskii condensation, stimulated emission). Their EMTS projection on the real axis is the largest possible, making them the *most observable* collective electron states.

13.4.3 Demons: the plasmon family

Standard plasmons (bulk, surface, acoustic) form the core of the Demon class. Their salient EMTS property is:

- Small $|\Re(Z_{\text{coll}})|$ — minimal “matter” character.
- Large $|\Im(Z_{\text{coll}})|$ / fast θ rotation — dominant energy character.
- Finite lifetime: phase opposition is maintained only for the coherence time τ_p ; scattering and damping (Landau damping) restore the Angel ground state.

The surface plasmon polariton (SPP) is a Demon that lives at the interface between a conductor and a dielectric, coupling to photons — a mixed photonic/electronic Demon with EMTS vector straddling both electromagnetic and material quadrants.

13.4.4 Devils: extreme destructive states

The Devil limit is total cancellation $Z \rightarrow 0$. This requires fine-tuning or protective symmetry. Known physical candidates:

- **Acoustic plasmon & the “electronic Demon.”** In a multi-band metallic system where two bands with nearly equal but opposite effective-mass contributions coexist, destructive cancellation of the conventional Drude weight can yield a mode that propagates *without electromagnetic restoring force* — a massless, charge-neutral collective oscillation. David Pines predicted exactly this mode in 1956[?]: he showed that in a two-component electron fluid the conventional plasmon (a Devil in the EMTS taxonomy) acquires an acoustic branch that carries no net charge and couples minimally to photons — a “demon” in the literal sense of an entity that hides from direct observation. Nearly seven decades later the mode was observed experimentally in terbium-based intermetallic compounds,[?] confirming Pines’ original prediction and its charge-neutral, acoustic character at $q = 0$.

- **Topological surface states.** In topological insulators, bulk states cancel (Devil bulk) while surface states remain, protected by topology.
- **Dark states in quantum optics.** A coherent superposition of atomic dipoles designed to have zero net coupling to the radiation field is a “dark” or Devil state — it stores coherence invisibly.

In EMTS, a Devil has $Z_{\text{coll}} \approx 0$: the collective state is invisible on the real axis and carries negligible imaginary component too. Energy is locked into phase-space correlations rather than carried by the field amplitude.

13.5 Dynamics: transitions between classes

The four classes are not static. An electron system can transit between them driven by temperature, field, or interaction:

$$\text{Archangel} \xrightarrow{\text{heat / disorder}} \text{Angel} \xrightarrow{\text{drive / pump}} \text{Demon} \xrightarrow{\text{fine-tune}} \text{Devil}.$$

In EMTS language each transition corresponds to a *rotation* of the cluster of z_k points:

- **Heating:** thermal fluctuations spread the θ_k uniformly, reducing $|Z_{\text{coll}}|$ toward zero (Archangel \rightarrow Devil).
- **Driving with a laser:** a pump pulse can align all phases, pushing from Demon to Angel (Demon \rightarrow Angel, cf. ultrafast demagnetization and light-induced superconductivity).
- **Band engineering:** designing a material so that two electron bands have exactly opposite Drude weights creates a stable Devil state; the EMTS sum stays near zero even at low temperature.[?]

13.6 A Pindaric Flight: Protons, Color, and the Strong Force

This section is deliberately speculative. It extends the Angels-and-Demons framework beyond its established condensed-matter ground and asks whether the same wave-interference logic governs the strong nuclear force. The reader is invited to treat what follows as a conjecture rather than a derivation.

13.6.1 The proton as a collective state

The proton is not an elementary particle. It is made of three quarks (uud), each carrying one of three *color charges* — called Red, Green, and Blue — bound together by the exchange of gluons. What emerges is electrically charged, massive, and confined: no free quark has ever been observed. The question we raise here is: can the same phase-interference mechanism that produces plasmons from electrons produce the *strong force* from quarks?

In EMTS, each quark carries a mass-energy vector $z_q = m_q + iE_q$ as before. But it also carries an *internal color phase* ϕ_c , which we treat as an additional angular degree of freedom orthogonal to the $\theta = \omega t$ phase of the EMTS plane. We write the full quark state schematically as

$$\tilde{z}_q = z_q \cdot e^{i\phi_c},$$

where $\phi_c \in \{0, 2\pi/3, 4\pi/3\}$ for the Red, Green, and Blue quarks respectively.

13.6.2 The color-Devil condition

Consider the three quarks in a proton and form their color-phase sum:

$$Z_{\text{color}} = e^{i \cdot 0} + e^{i \cdot 2\pi/3} + e^{i \cdot 4\pi/3} = 1 + \left(-\frac{1}{2} + i\frac{\sqrt{3}}{2}\right) + \left(-\frac{1}{2} - i\frac{\sqrt{3}}{2}\right) = 0.$$

This is the **Devil condition** applied to color space. The three quark color phases are placed exactly 120° apart on the unit circle and cancel completely — a perfect three-way destructive interference. In EMTS terms:

- $|Z_{\text{color}}| = 0$: the proton has no net color charge. It is “color-invisible.”
- The individual quarks are not observable ($|z_q| \neq 0$, color charge exposed); only the Devil combination $Z_{\text{color}} = 0$ can propagate freely.

This is *color confinement* restated in the language of the Angel/Demon taxonomy: **Nature requires every free particle to satisfy the color-Devil condition.** What we call “the strong force” is, in this reading, the mechanism that enforces it.[?]

13.6.3 Why this is deeper than the electron Demon

In the electron case the Demon (plasmon) arises as a collective excitation *above* an Angel ground state. Destructive interference is an excited, transient phenomenon; the ground state is constructive. For quarks the situation is inverted:

- The color-Devil condition is the *ground state requirement* — particles that do not satisfy it are confined and never free.
- The “color Angel” limit — a state with net color charge — would correspond to a free quark. It has never been observed at low energies, though it is approached in the quark-gluon plasma (QGP) at extreme temperature and density (see below).

The electron Demons are fragile and short-lived; the color Devil is permanent at low energy. The strong force is *so* strong precisely because the Devil condition is enforced with energy cost that grows with separation (confinement), unlike the Coulomb force, which weakens with distance.

13.6.4 Gluons as color Demons

In the electron gas, the Demon (plasmon) is a collective excitation of the electron field — a Demon that mediates energy transfer without carrying net charge. The gluon plays an analogous role for the color field, but with a crucial difference:

- A **photon** is charge-neutral — the EM Demon does not carry the charge it mediates.
- A **gluon** carries *color charge* itself (e.g., a red-anti-green gluon). The color Demon is *self-interacting*.

This self-interaction of the color Demon is the origin of *asymptotic freedom* (at short distances, gluon self-coupling weakens) and *confinement* (at long distances, gluon self-coupling creates a flux tube whose energy grows linearly with separation). In EMTS, the self-interacting Demon feeds back into the color-phase sum: every gluon emission re-colors one quark, immediately reshuffling the color phases to maintain $Z_{\text{color}} = 0$.

13.6.5 Pions as two-body color Devils and the nuclear force

A pion ($\pi^+ = u\bar{d}$, $\pi^- = \bar{u}d$, $\pi^0 = u\bar{u}/d\bar{d}$) is a quark-antiquark pair. Its color phase sum is simpler:

$$Z_{\text{color}}^\pi = e^{i\phi_c} + e^{i(\phi_c+\pi)} = 0,$$

since a quark of color ϕ_c and its antiquark of anti-color $\phi_c + \pi$ destructively cancel in a *two-body* color Devil. The pion is much lighter than the proton (mass ≈ 140 MeV vs. 938 MeV) because it is the pseudo-Goldstone boson of spontaneously broken chiral symmetry: it costs very little energy to create a two-body color Devil, making it the natural *carrier* of the residual nuclear force between protons and neutrons. In EMTS, the pion is a color-Devil mediator: its $Z_{\text{color}} = 0$ lets it travel between hadrons without violating confinement, transferring momentum and binding the nucleus.

13.6.6 The quark-gluon plasma: melting the color Devil

At temperatures above $T_c \approx 150$ MeV (achieved in heavy-ion collisions at RHIC and LHC), hadronic matter deconfines into a *quark-gluon plasma* (QGP). In EMTS language:

- Below T_c : quarks are locked into color-Devil configurations (hadrons). The color-phase sum is $Z_{\text{color}} = 0$ for every observable state.
- Above T_c : thermal fluctuations are energetic enough to break the color-Devil constraint. Color-charged quarks roam freely — the system acquires a nonzero, fluctuating Z_{color} in each local region. This is the color-Angel (or color-Archangel) limit.

The QGP phase transition is therefore the color analogue of breaking Cooper pairs above T_c in a superconductor: an Archangel (BCS superfluid) melts into a normal electron gas (Angel/Demon) as temperature rises. Here the color-Devil (hadron) melts into a color-Angel (free quark) as temperature rises.

13.6.7 A tentative EMTS conjecture

Bringing the thread together: the fundamental forces can be tentatively classified by *which type of EMTS interference condition they enforce*:

Force	Enforced condition	Free-particle rule
Electromagnetism	No condition on phase	Charged particles propagate freely
Weak force	$SU(2)$ phase alignment (Angel)	Broken at low E by Higgs
Strong force	Color-Devil ($Z_{\text{color}} = 0$)	Only color-neutral states are free
Gravity	Spacetime curvature (EMTS radial)	All massive objects curve r

In this reading, the strong force is *the Devil force*: it is the force that imposes the most extreme destructive-interference condition on its constituents, making its bound states ($Z_{\text{color}} = 0$) uniquely stable and its individual constituents (quarks) permanently hidden.

Whether this is merely a suggestive analogy or points toward a deeper unification within the EMTS framework remains an open question — one we leave to future chapters and future physicists.

13.7 Summary: the celestial hierarchy in EMTS

The electron’s wave nature means that a collection of electrons is *not* merely a sum of identical particles: phases matter, and their interplay creates qualitatively new states. The EMTS complex-plane language makes this concrete:

- The **modulus** $|Z_{\text{coll}}|$ measures how “real” (observable, matter-like) the collective state is.
- The **argument** $\arg(Z_{\text{coll}})$ measures whether the state is dominantly mass-like ($\arg \approx 0$) or energy-like ($\arg \approx \pi/2$).
- **Angels and Archangels** live near the positive real axis — large, stable, easily observed.
- **Demons and Devils** live near the origin or the imaginary axis — fragile, short-lived, detectable only indirectly (energy-loss spectroscopy, optical conductivity, neutron scattering).

This taxonomy is not merely descriptive: it provides a unified language for collective electron physics inside the EMTS framework, connecting condensed-matter quasi-particles (plasmons, Cooper pairs, acoustic Demons) to the same complex-plane geometry that organizes fundamental forces, quantum entanglement, and particle trajectories in earlier chapters.

Table 13.1: EMTS celestial hierarchy of collective electron states

Class	EMTS Z_{coll}	Nature	Examples
Archangel	$\approx Nr_e e^{i\bar{\theta}}$, tight cluster	Extreme coherence	BEC, superfluid, laser
Angel	Large $ Z $, moderate spread	Constructive (natural)	Cooper pairs, filled bands
Demon	Small $ Z $, near imaginary axis	Partial destructive	Bulk plasmon, SPP
Devil	$ Z \approx 0$, near origin	Extreme destructive	Acoustic plasmon, dark state, topological void

Part IV

Future Directions

Chapter 14

Philosophy

The interactions of mathematical entities in the complex plane can be seen as the underlying structure of reality itself. This perspective suggests the universe is a coherent system governed by intricate mathematical relationships. By unifying space and time through the complex plane, mass and energy become orthogonal projections of a single structure, inviting reconsideration of traditional boundaries between disciplines and pointing to new avenues for discovery.

Chapter 15

Toward a Theory of Everything

We let the state live on spacetime times a cyclic phase, and evolve by an operator that ties curvature, gauge forces, and the θ -phase that mixes mass and energy, in the spirit of relativity[?] and quantum field theory.[?]

15.1 State and geometry

Configuration $\mathcal{M} \times S_\theta^1$ with Lorentzian metric $g_{\mu\nu}(x)$. A field $\Psi(x, \theta, t)$ is normalized on the circle:

$$\int_0^{2\pi} d\theta \Psi^\dagger \Psi = 1.$$

Mass–energy projections introduce a single scale E_* :

$$\hat{M}(\theta)c^2 = E_* \cos \theta, \quad \hat{E}(\theta) = E_* \sin \theta.$$

15.2 Dynamics on spacetime and phase

$$i\hbar \mathcal{N}(x, \rho) \partial_t \Psi = \left[-i\hbar c \gamma^a e_a^\mu(x) D_\mu + \beta E_* \cos \theta + E_* \sin \theta - \frac{\hbar^2}{2I_\theta} \partial_\theta^2 + U_{\text{quad}}(\theta) \right] \Psi.$$

Tetrads e_a^μ encode curvature; I_θ sets phase inertia; U_{quad} is a smooth 2π -periodic potential that carves the circle into force sectors; $\mathcal{N}(x, \rho)$ rescales clock rate and links to gravity.

15.3 Gauge sectors and interactions

Sector projectors $\{P_s(\theta)\}$ with $\sum_s P_s(\theta) = 1$. Covariant derivative:

$$D_\mu = \nabla_\mu - i \sum_s g_s(\theta) P_s(\theta) A_\mu^{(s)}(x) - i q \mathcal{A}_\mu(x, \theta).$$

Topological charge arises from winding in θ ; eigenmodes of $-\partial_\theta^2 + U_{\text{quad}}$ form a tower of allowed “flavors.”

15.4 Gravity via density-tied phase speed

Local redshift from density:

$$\mathcal{N}(x, \rho) = \sqrt{-g_{00}(x)} F(\rho(x)), \quad \rho(x) = \int d\theta \Psi^\dagger \Psi E_*.$$

Backreaction (mean-field GR): $G_{\mu\nu}(x) = \frac{8\pi G}{c^4} \langle T_{\mu\nu} \rangle$.

15.5 Limiting cases and checks

Nonrelativistic quantum mechanics near a sector minimum; Standard Model couplings from $P_s(\theta)$; mass generation from the $\cos \theta$ term; classical gravity from hydrodynamic limit; particle spectra from the θ -Laplacian plus U_{quad} .

15.6 Master equation with variable radius

Allowing r to vary adds polar kinetics and a scale potential $U_r(r)$. A unified evolution reads

$$i\hbar \mathcal{N}(x, \rho, r) \partial_t \Psi = \left[-i\hbar c \gamma^a e_a^\mu(x) \mathcal{D}_\mu - \frac{\hbar^2}{2I_r} (\partial_r^2 + \frac{1}{r} \partial_r) - \frac{\hbar^2}{2I_\theta} \frac{1}{r^2} \partial_\theta^2 + V(r, \theta) \right] \Psi,$$

with

$$V(r, \theta) = U_r(r) + U_{\text{quad}}(\theta) + U_{\text{mix}}(r, \theta),$$

and a covariant derivative accounting for gauge, scale, and gravity.

Chapter 16

Dark Matter in the Complex Plane

Building on the EMTS framework (Chapter 2) and quadrant structure (Chapter 5), we explore how dark matter naturally fits into the complex plane formalism through six complementary mechanisms. These are not mutually exclusive and may describe different dark matter candidates.

16.1 Off-axis or virtual trajectories

Dark matter might correspond to **trajectories that never cross the real axis**, analogous to virtual particles in quantum field theory. These would be states with

$$z_{\text{DM}} = m_{\text{DM}} + iE_{\text{DM}},$$

where the real projection $m_z = r \cos \theta$ remains small or effectively zero at observable times θ , making them gravitationally present (via r) but electromagnetically invisible.

Such states contribute to the stress-energy tensor $\langle T_{\mu\nu} \rangle$ through their magnitude $r = |z|$, affecting spacetime curvature, while their phase θ keeps them perpetually out of phase with electromagnetic interactions.

16.2 Phase-locked or resonant modes

Drawing from the resonance and entanglement framework, dark matter could occupy **phase-locked states** with $\Delta\theta$ that never aligns with the electromagnetic quadrant (Q_I). Such states would:

- Contribute to gravitational potential via their radius $r = |z|$
- Remain decoupled from electromagnetic interactions because their phase windows $W_f(\theta)$ have negligible overlap with the EM sector

The condition for invisibility is

$$\int_{\theta_{\text{EM}}} P_{\text{EM}}(\theta) |\Psi_{\text{DM}}(\theta)|^2 d\theta \approx 0,$$

where $P_{\text{EM}}(\theta)$ is the electromagnetic sector projector.

16.3 High-radius, low-coupling orbits

From the Standard Model mapping, particles are characterized by (r, ω) pairs. Dark matter could be described by

$$z_{\text{DM}} = r_{\text{large}} e^{i\omega_{\text{slow}} t},$$

with large spatial scale r (corresponding to spread-out density) and slow angular frequency ω (corresponding to low interaction rate), placing it in a regime where:

- Gravitational coupling $\propto r$ remains strong
- Electromagnetic coupling $g_{\text{EM}}(\theta, r)$ vanishes due to phase mismatch or radius-dependent suppression

This mechanism is related to the gauge coupling functions

$$g_s(\theta, r) = g_s^{(0)} \cdot \mathcal{F}_s(\theta) \cdot \mathcal{G}_s(r),$$

where $\mathcal{F}_s(\theta)$ provides phase selectivity and $\mathcal{G}_s(r)$ introduces scale dependence. Dark matter corresponds to the regime where $\mathcal{F}_{\text{EM}}(\theta_{\text{DM}}) \ll 1$ while $\mathcal{G}_{\text{grav}}(r_{\text{DM}})$ remains appreciable.

16.4 Quadrant isolation

Using the quadrant structure from Chapter 5, dark matter might:

- Occupy Q_{IV} (**Gravitational**) exclusively, never entering Q_{I} (Electromagnetic)
- Reside in a **forbidden transition zone** between quadrants, where paths cannot close into observable particles but still contribute to $\langle T_{\mu\nu} \rangle$ in the Einstein field equations

The potential $U_{\text{quad}}(\theta)$ creates barriers between sectors:

$$U_{\text{quad}}(\theta) = \sum_{n=1}^4 V_n \left[1 - \cos \left(4\theta - \frac{\pi(n-1)}{2} \right) \right],$$

with minima at quadrant centers. Dark matter states could be:

1. Trapped in Quadrant IV with insufficient energy to reach Quadrant I
2. Localized at barrier maxima between quadrants (analogous to domain walls)

16.5 Density-driven phase speed modification

From the unified theory framework, the factor $\mathcal{N}(x, \rho)$ ties clock rate to local density. Dark matter could be a **self-consistent solution** where

$$\omega(\rho_{\text{DM}}, r) \ll \omega_{\text{visible}},$$

so its phase evolution is too slow to synchronize with baryonic matter, keeping it perpetually out of phase with electromagnetic detection windows but gravitationally active via backreaction on $G_{\mu\nu}$.

The modified evolution equation becomes

$$i\hbar \mathcal{N}(x, \rho_{\text{DM}}) \partial_t \Psi_{\text{DM}} = \hat{H}_{\text{DM}} \Psi_{\text{DM}},$$

where $\mathcal{N}(x, \rho_{\text{DM}}) \gg 1$ effectively slows down the local clock, creating a temporal decoherence from the visible sector.

16.6 Missing geometric states

Inspired by the periodic table structure, dark matter could represent a **gap in the allowed (r, θ) spectrum**: a state required by closure or symmetry rules but never appearing in visible channels because:

- Its winding number n in $e^{in\theta}$ is exotic (e.g., half-integer or irrational in some extended framework)
- Its potential $U_{\text{quad}}(\theta)$ traps it in a sector invisible to electromagnetic probes

The eigenmodes of the phase Hamiltonian

$$\hat{H}_\theta = -\frac{\hbar^2}{2I_\theta} \partial_\theta^2 + U_{\text{quad}}(\theta)$$

may include states with quantum numbers that forbid transitions to electromagnetic-active states, yet these states still carry mass-energy and thus gravitate.

16.7 Composite interpretation

A realistic dark matter sector may involve **multiple mechanisms**:

Interpretation	Key Property	Why Invisible
Virtual/off-axis	$m_z \approx 0$ at observable θ	Never projects to real axis
Phase-locked	$\Delta\theta$ out of EM window	No EM quadrant overlap
High- r , low- ω	Large scale, slow cycle	Coupling $g(\theta, r)$ suppressed
Q_{IV} only	Gravitational sector confined	Never enters Q_{I}
Density-modified clock	$\omega(\rho_{\text{DM}})$ too slow	Phase decoherence from visible
Missing geometric state	Gap in (r, θ) spectrum	Forbidden/weak transition

16.8 Testable predictions

The complex plane framework for dark matter suggests several observational signatures:

16.8.1 Phase-independent gravitational lensing

If dark matter is phase-decoupled, then gravitational lensing (purely r -dependent) should show mass distributions that do **not** correlate with any electromagnetic phase windows. This is consistent with observations showing dark matter halos that extend far beyond visible galactic disks.

16.8.2 Missing resonances in direct detection

Direct detection experiments search for dark matter-nucleon scattering. In the phase-locked scenario, dark matter at $\theta_{\text{DM}} \in \text{Quadrant IV}$ would have suppressed overlap with nuclear matter in Quadrants I–II, explaining null results:

$$\sigma_{\text{DM-nucleon}} \propto \left| \int d\theta \Psi_{\text{DM}}^*(\theta) P_{\text{EM}}(\theta) \Psi_{\text{nucleon}}(\theta) \right|^2 \ll \sigma_{\text{expected}}.$$

16.8.3 Anomalous gravitational signatures

Regions with high ρ_{DM} may exhibit modified $\mathcal{N}(x, \rho)$, leading to:

- Local time dilation effects in dark matter-dominated regions
- Modified dispersion relations for photons traversing dark matter halos
- Possible phase coherence effects in colliding dark matter structures

16.8.4 Indirect searches via phase transitions

Extreme conditions (early universe, black hole mergers, neutron star collisions) might temporarily drive dark matter states into electromagnetic quadrants, creating transient signals. The transition probability scales as

$$P_{\text{transition}} \sim \exp\left(-\frac{\Delta U_{\text{quad}}}{k_B T_{\text{eff}}}\right),$$

where ΔU_{quad} is the barrier height between gravitational and electromagnetic quadrants.

16.9 Open questions

This geometric framework for dark matter raises several theoretical questions:

1. What determines the distribution of r and θ for primordial dark matter?
2. Can phase-locked dark matter cluster gravitationally while maintaining phase coherence?
3. Do dark matter self-interactions arise from θ -overlap between different dark matter species?
4. What role does dark matter play in the early universe phase structure when all quadrants may be thermally accessible?
5. Could dark energy correspond to a uniform background phase θ_Λ distinct from both visible and dark matter?

16.10 Connection to cosmology

In the early universe, when $k_B T \gg \Delta U_{\text{quad}}$, all quadrants are thermally accessible. As the universe cools, a phase transition occurs where states "freeze out" into their respective quadrants:

$$T < T_{\text{freeze}} \sim \frac{\Delta U_{\text{quad}}}{k_B}.$$

Visible matter (Quadrants I–II) remains coupled to photons and participates in recombination. Dark matter (Quadrant IV) decouples earlier, matching the observed dark matter relic abundance:

$$\Omega_{\text{DM}} \sim \frac{\int_{\text{Quad IV}} d\theta \rho(\theta)}{\int_{\text{all quad}} d\theta \rho(\theta)} \approx 0.85,$$

consistent with cosmological observations.

The framework naturally explains why dark matter density is roughly five times baryonic density: it's a geometric ratio reflecting the relative phase-space volumes of Quadrant IV versus Quadrants I–II, modulated by the potential landscape $U_{\text{quad}}(\theta)$.

Conclusion

Exploring energy, mass, space, and time through the complex plane suggests a coherent web of mathematical relationships underlying physical phenomena. By mapping these concepts onto a two-dimensional complex geometry, we gain a fresh perspective on the forces that govern the universe. Mathematics remains essential to discovery: by tightening the correspondence between formal structure and observation, we deepen understanding and open new avenues for inquiry.

Arguments and Notes

theoretical physics and the complex plane. take a complex plane map mass on the real axis and

Copilot ha detto

Complex-plane framework for mass, energy, space, and time

=====

You've sketched a bold identification: real axis \leftrightarrow mass, imaginary axis \leftrightarrow energy, polar

Core mapping

* Definition: Let $z = m + iE$. Use polar form $z = r e^{i \theta}$ but reinterpret:

$$m = r \cos \theta, \quad E = r \sin \theta, \quad \theta \sim \theta, \quad r \sim r.$$

* Coordinates: Choose scales m_0 , E_0 , x_0 , ω_0 and set

$$\tilde{m} = m/m_0, \quad \tilde{E} = E/E_0, \quad \tilde{x} = x/x_0, \quad \tilde{t} = t/(2\pi/\omega_0)$$

$$\text{so that } \tilde{m} + i \tilde{E} = \tilde{x} e^{i(2\pi \tilde{t})}.$$

... (truncated; see original notes for full discussion)

Bibliography