



Non-target impacts of forest defoliator management options: Decision for no spraying may have worse impacts on non-target Lepidoptera than *Bacillus thuringiensis* insecticides

J. Mark Scriber

Department of Entomology, Michigan State University, East Lansing, MI 48824 (e-mail: Scriber@msu.edu; phone: +1-517-432-1975; fax: +1-517-353-4354)

Received 18 December 2003; accepted in revised form 12 May 2004

Key words: *Bacillus thuringiensis*, Butterflies, Disease, Moths, Parasites, Predators

Abstract

Management programs for major forest defoliators such as gypsy moths or forest tent caterpillars, and crop pests such as the European corn borer have shifted from broad-spectrum insecticides to more environmentally benign microbial pesticides such as *Bacillus thuringiensis* (foliage sprays and transgenic toxin expression in plant tissues). Phytochemically resistant host plants and natural enemies have been used as alternative pest management strategies (including generalist tachinid flies such as *Comptosia*, viruses, microsporidians, and fungi), but all of these have some non-target impacts, as described from literature review. A sequence of lab and field studies were conducted to determine non-target impacts on native Lepidoptera in North America. The conclusions reached are that a decision not to spray *Bt* pesticides (i.e. to allow defoliation and natural pest outbreaks to run their course) could be as bad or worse for non-target Lepidoptera as the microbial insecticides would be. The important concept that must be maintained is that all pest management programs have some risk of negative non-target impacts, but it is the magnitude and relative importance that will remain the most critical issue for environmental impacts and pest management.

Controlling phytophagous insect outbreak defoliations

Management of catastrophic forest defoliators and agricultural/ornamental plants with broad-spectrum insecticides has had short-term successes, but also resulted in serious and extensive non-target impacts. Alternatives to such insecticides have been sought with more specific and environmentally benign microbially derived insecticides such as *Bacillus thuringiensis*, Laird et al. 1990 *Bt*; (Faust and Bulla 1982; Beegle and Yamamoto 1992; Leong et al. 1992; Lambert and Peferoen

1992; Reardon et al. 1994). However, drift of *Bt* insecticides with aerial application has been reported as far as 3 km downwind, depending on method of application (Barry et al. 1993; Swadener 1994; Whaley et al. 1998). On foliage, viable spores have also been recovered weeks (or years) after application (Reardon and Haissig 1984; Huang et al. 1990). Such undesirable environmental distribution of *Bt* sprays have led to even safer genetically engineered plants being developed with tissue-specific expression of *Bt* endotoxins (or other toxins and growth inhibitors such as plant lectins, or protease- and amylase-inhibitors; Babu

et al. 2003). The hopes were to achieve acceptable pest control via 'host plant resistance' (Maxwell and Jennings 1980; Fritz and Simms 1992) using modern transgenic techniques and minimizing the dangers to non-target organisms of all types (Reardon 1994; Robison and Raffa 1994; Wang et al. 1996; Babu et al. 2003; Nap et al. 2003; Conner et al. 2003; Hancock 2003). Nonetheless, undesirable non-target impacts of *Bt* insecticides (whether externally on leaf surfaces or tissue-specific toxin expression) will occur, to varying degrees, depending on many environmental (abiotic) and biotic variables (some known, but most unknown: Smith and Couche 1991; Altmann 1992; Addison 1993; Martin 1994; Moldenke et al. 1994; Ferber et al. 1999; van Emden 1999; Hails 2000; Scriber 2001; Shelton et al. 2002).

Sprays of *Btk* (variety *kurstaki* used against Lepidoptera) employed against European and Asian gypsy moths (*Lymantria* spp), forest tent caterpillars (*Malacosoma* spp), spruce budworms (*Choristoneura* spp.) had negative impacts on many beneficial (biological controls) and non-target invertebrates (Muck et al. 1981; Flexner et al. 1986; Dreistadt and Dahlsten 1989; Melin and Cozzi 1990; Miller 1990; Miller 1992; Chapman and Hoy 1991; James et al. 1993; Addison 1993; Swadener 1994; Johnson et al. 1995; Papp-Herms 1996; Sample et al. 1996; Chenot and Raffa 1998; Peacock et al. 1993 and 1998; Scriber 1998; Whalley et al. 1998; Boulton 1999; Glare and O'Callaghan 2000; Rastall et al. 2003). Based upon phenological overlap with the gypsy moth, host-sharing, and larval habitat (canopy/understory), it was determined that 92–98% of the locally recorded 223 species of Lepidoptera (from 22 families) had measurable vulnerability to *Bt* sprays in the National Parks of Washington, DC (Venables 1990). In a different study of 42 species of Lepidoptera, all four species of butterflies were extremely sensitive to *Btk*, in contrast to only 10 of the 38 moth species (geometrids, lymantriids, and noctuids were least sensitive; Peacock et al. 1998). It seems that butterfly larvae are generally more susceptible than moth larvae, and therefore may be useful as an 'early warning' signal for environmental danger in different habitats or ecosystems (see also Scriber and Gage 1995; Brown 1997; New 1997).

While the direct measurable impacts of *Bt* on non-target invertebrates have been noted above,

there are other ways that *Bt* might indirectly impact non-targets. Although not adequately measured, some of these include the sub-lethal effects of *Bt* on food consumption, growth rates, and increased vulnerability of herbivorous insects to parasites or predators (Johnson and Gould 1992; Johnson 1997). The behavioral preference or avoidance of *Bt* on leaf surfaces has not been thoroughly evaluated for phytophagous insects, but such adaptations to plant protectants may evolve (Gould 1991). Similarly, the sub-lethal impacts of *Bt* (with regard to pathogen susceptibility) remains largely unknown. The relative competitive abilities of *Bt*-impacted phytophagous larvae have not been thoroughly assessed in field or even lab situations.

The fate of *Bt* crystals, spores, and the ecological interactions with related microbes remains basically unknown (Doane 1970; Feitelson et al. 1992; Addison 1993; Watanabe et al. 1998). The persistence of toxic and chronic effects of *Bt* applied on leaf surfaces (or genetically engineered into crop plants and trees) remains variable, with little understanding why (Smith and Couche 1991; Martin 1994; Bauer 1997; Scriber 2001). Sprays are differentially effective (Krieg and Langenbruch 1981; Wagner et al. 1996) because of the varied Lepidopteran guilds with different feeding behavior/ecology. Some species are protected/shielded (e.g. stem borers, bud feeders, leaf-miners, leaf-rollers, leaf-tiers, etc) versus exposed/unprotected external feeders (Slansky and Scriber 1985; Scoble 1992; Stamp and Casey 1993; Sheehan 1994) and the bulk of Lepidoptera are living on leaf surfaces (as with the Papilionidae and Saturniidae; Scriber et al. 1995b; Tuskes et al. 1996). Early instars tend to be more susceptible to *Bt* than later instars, but this is not always the case (Navon et al. 1990; Wagner et al. 1996; Peacock et al. 1998). The environmental persistence, genetic exchanges via hybridization, and ecological interactions of *Bacillus* used in control programs with other microbes surprisingly remain largely unknown (Hails 2000).

Bacillus thuringiensis

Bacillus thuringiensis and related crystalliferous bacteria have been produced in many countries

with widely varying methods and variable results. Before 1996, there were at least 137 insect species (Lepidoptera, Coleoptera, Diptera, and Hymenoptera) known to be susceptible (Heimpel 1967). Since then, the production has been more standardized and contamination of Btk with other microbes or different *Bt* toxins such as *Bti* has been reduced (Martin and Trivers 1989; Reardon et al. 1994; Swadener 1994). Within the species *B. thuringiensis*, at least 16,000 (and perhaps as many as 40,000) strains had been isolated (Lambert and Peferoen 1992; Addison 1993) with at least 34 different subspecies or serotypes (de Barjac and Frachon 1990). Smith and Couche (1991) isolated *Bt* from unsprayed leaf surfaces of hardwood and conifer species. These bacteria were naturally present throughout the year and may have symbiotically functioned to protect leaves from phytophagous insects. It has been shown that *Bt* readily germinates, produces vegetative cells, and can grow and reproduce successfully in soils (Addison 1993). Ultraviolet light kills most of the phylloplane spores of *Bt*, but whether germination, growth, and reproduction occur is still undetermined, but suspected, especially in shaded areas and on the undersides of leaves (Addison 1993; Scriber 1998). The lack of understanding of the natural ecology of *B. thuringiensis* has been clearly recognized but not addressed (Martin and Travers 1989; Lambert and Peferoen 1992; Addison 1993). We do not know the functional role *B. thuringiensis* in the soil (or on leaf phylloplanes; Andrews and Hirano 1991), whether it has any 'natural' hosts, what feeds on it, or how it interacts with other microorganisms (including the closely related natural *B. anthrax* and *B. cereus*, which are capable of exchanging plasmids; Helgason et al. 2000; Kanda et al. 2000).

When *Bt* cell replication is complete, an endospore and a protein crystal are formed within the vegetative cell, called a sporangium. When the cell wall breaks down the spore and crystals are released. The crystal is a large molecule that is not toxic to insects until solubilized by digestive processes in the gut, which releases smaller delta-endotoxin crystals. The spore coat can also contain proteins that act as toxins (Reardon et al. 1994). The crystal types of *Bt* have been classified by Hofte and Whitely (1989) into Cry I and Cry II (both with subgroups) primarily toxic to

Lepidoptera, Cry III toxic to Coleoptera and Cry IV toxic to Diptera.

Persistence of *Bt*

The toxic activity of *Btk* toward target Lepidoptera has been reported to disappear within a few days after application to agricultural crops (Ignoffo et al. 1974; Leong et al. 1980; Beegle et al. 1981). *Bt* disappears within a week or two in the soil (West and Burgess 1985; or longer, Petras and Cassida 1985). However, in forest ecosystems, the results have been quite variable, ranging from 1 day for gypsy moths on oak trees (Sundaram and Sundarum 1992) or 3–4 days in conifers for western spruce budworm (Beckwith and Stelzer 1987). In other studies activity against gypsy moths persisted up to 60 days (Miller and West 1987). Toxicity of Btk 30–40 days post-spray was shown for *Papilio* (Johnson et al. 1995). Nonetheless the long-held belief (misconception) persists that 'within hours, or more conservatively, within a few days, Bt toxin levels are expected to be ineffective in reducing most lepidopteran larval populations' (Rastall et al. 2003).

Variability in persistence of *Btk* toxicity at a given dose/density (Bryant and Yendol 1988; Radcliffe and Yendol 1993) may be due to the type of foliage (and its natural phytochemical composition; Felton and Dahlman 1984; Meade and Hare 1993, 1994; Krischik et al. 1988; Reichelderfer 1991; Arteel and Lindroth 1992; Appel 1994; Huang et al. 1994), environmental conditions (rain, sun/shade, temperatures etc.; Reardon et al. 1994), and the relative susceptibility of the insect species/genotypes being targeted (Ignoffo 1992; Gould et al. 1992; Lambert and Peferoen 1992; Bauer 1995; Scriber and Haas 1995; Gould 1998; Scriber 1998; Gahan et al. 2001; Ferre and van Rie 2002). The variable susceptibility of different Lepidopteran species may be related to the presence of spores in the *Bt* spray (Pinnock et al. 1971; Burgess et al. 1976), or caused by the different toxicity of various mixtures of crystal proteins (van Frankenhuyzen et al. 1991), or due to synergized toxic effects of *Bt* in the presence of Bt spores or even other bacteria on host foliage (Dubois and Dean 1995).

Might a decision not to spray forests for gypsy moths may be as bad or worse for non-target Lepidoptera (and other organisms)?

Management of defoliating insects such as gypsy moths via insecticidal sprays such as Btk may result in serious non-target effects as described above. However, a management decision not to spray, allowing the defoliator outbreak to run its course, may have equally undesirable non-target impacts. The desiccation of the forest understory impacts many members of delicately balanced communities some with threatened/endangered wildflowers, the high levels of nitrogenous runoff into streams and lakes can pollute the water, and the repeated defoliation of forest trees can create stress-vulnerability and tree mortality (especially in drought years). Perhaps the most significantly negative for phytophagous Lepidoptera, the large and rapid increases of natural enemies of all types associated with outbreak defoliator insect populations can have profound negative effects that may be worse than the *Bt* sprays. An example from North America illustrates some of the issues that may face non-target Lepidoptera elsewhere.

The gypsy moth, *Lymantria dispar* (L.), has caused serious broadleaf forest defoliations in eastern North America and continues to spread southward in the Appalachian mountains and westward through the Great Lakes region (Gage et al. 1990; Gage and Pijanowski 1993). The highly accelerated spread through Michigan's northern aspen/birch forests in 1990–1993 (Sharov et al. 1999) was especially alarming for foresters because of the widespread defoliation damage, mixed with two late defoliating spring freezes in 1992 (Scriber and Gage 1995; Gage 1996).

Cyclic periods (8–10 year intervals) of outbreaks follow the initial wave of gypsy moth outbreak and defoliation (Elkinton and Liebhold, 1990; Liebhold et al. 1997; Dwyer et al. 2004). These population crashes may occur only after 3 to 4 years depending on many factors, including parasitoids (e.g. *Compsilura concinnata*, Scriber 1998), nuclear polyhedrosis viruses (Kukan and Myers 1997), the fungus *Entomophaga maimaiga* (Hajek et al. 1995; Hajek et al. 1996), how much *Bacillus thuringiensis* is applied in suppression management efforts (Johnson et al. 1995; Scriber

1998), as well as time-delayed maternal effects (Rossiter 1994). Also, in 1994 and 1996, extremely cold periods and winter mortality of gypsy moth egg masses may have contributed directly to population crashes throughout northern Michigan (Smitley et al. 1998).

Severe defoliations by one herbivore species such as the gypsy moth were known to affect the quality of the host for other species (Faeth 1987; Hunter 1992; Denno et al. 1995; Stewart 1996; Karban and Baldwin 1997). Trees may respond to herbivore damage by increasing the allelochemical defenses or decreasing nutrient concentrations (Wallner and Walton 1979; Schultz 1988; Karban and Myers 1989; Bryant et al. 1987; Neuvonen et al. 1987; Tallamy and Raupp 1991; Koricheva et al. 1998). Rapid induced resistance (RIR) can occur within hours or days, whereas delayed induced resistance (DIR) occurs in the year following defoliation and can last for several years (Haukioja et al. 1985; Baldwin 1994; Kaitaniemi et al. 1999; Parry et al. 2003; Nykanen and Koricheva 2004). Induced defenses in plants illustrate adaptive plasticity because the induced phenotype can exhibit higher fitness in environments with strong (defoliating) herbivory, whereas the uninduced phenotypes can show higher fitness in environments not having intense herbivory (Karbon et al. 1999; Heil et al. 2004). Recent studies have shown that tiger swallowtail butterflies are significantly slowed in growth rate due to rapid induced resistance from forest tent caterpillar defoliation of quaking aspen, *Populus tremuloides* (Dankert et al. 1997). The growth reduction was proportional to the severity of the defoliation. Additional long-term-ecological-research (LTER) sites were established to experimentally manipulate the (genetically identical clonal) hybrid poplar ecosystem by severe defoliation using nearly 10-million (virus-treated) gypsy moth egg masses (see below).

What follows is a summary of selected studies undertaken in our Lepidoptera research program to address *Bt* non-target impact issues as well as impacts of decisions not to spray *Bt* (and impact of transgenic *Bt* in crops/forests). The experiments below represent a research overview to illustrate some major remaining gaps in our knowledge of non-target impacts and intended to highlight important future needs for additional research;

Direct impacts of aerially applied Btk sprays on non-target Lepidoptera (experimental results with giant silkmoths of the Saturniidae and butterflies of the Papilionidae)

Different host plant species

At four different sites in Michigan, we placed first and second instar larvae on Btk-sprayed and unsprayed control tree species at 1-days post-spray, including: 1) the silkmoth, *Callosamia promethea*, on black cherry, BC of the Rosaceae; white ash, WA of the Oleaceae; tulip tree, TT of the Magnoliaceae; and spicebush, SB of the Lauraceae) 2) the Eastern tiger swallowtail butterfly (*P. glaucus*, on BC, WA, TT, and hop-tree or HT of the Rutaceae), and 3) the Canadian swallowtail butterfly (*P. canadensis*, on BC, WA, Amelanchier, AM of the Rosaceae, Quaking aspen, QA, and Balsam poplar, BP, both of the Salicaceae). Sprays of Foray 48B were applied by backpack sprayer to our trees at the rate calibrated to equal to that achieved in aerial sprays of forest canopies (30 BIU/ha with 40 IU/cm²; BIU = billion international units/IU = infective units; see Johnson et al. 1995). Early instar larvae are most useful and ecologically meaningful in survival bioassays (Zalucki et al. 2002), and at 5-days post-spray (short-term impacts) we found near-total mortality of larvae on leaves sprayed with *Bt*, regardless of the host plant (94% of *P. canadensis*; 93% of *P. glaucus*; and 89% of *C. promethea*; Johnson et al. 1995), and essentially all of the larvae on *Bt* sprayed foliage were dead or gone by day 8.

The impact on larvae placed on BC tree leaves at an 'intermediate' interval of 12 days post-spray ($n = 202$ larvae on 15 sprayed/15 unsprayed trees for *P. glaucus* and $n = 120$ larvae on 10 sprayed/10 unsprayed trees for *P. canadensis* after an 8-day exposure) was as great as the 1-day ('short term') post-spray. For *P. glaucus*, 90% of the larvae died or disappeared compared with only 41 % of the control larvae under natural field conditions and predation; and nearly identical results were seen for *P. canadensis* larvae (Johnson et al. 1995).

Two consecutive years of 'long term' experiments on non-target impact assessments (at time intervals of 0-d 10-d 20-d, 30-d, and 40-d post-spray placement of larvae) consistently showed significantly more daily mortality on the *Bt*-sprayed trees than on the control trees. This

statistically significant mortality on *Bt*-sprayed trees (compared to paired unsprayed controls) persisted through 30-days post-spray (at 40 days the difference persisted, but was not statistically significant; $p = 0.06$). While the *Bt*-toxicity was not diminished significantly in the no-canopy sun and rain treatments of these field studies (Johnson et al. 1995), some attenuation of *Bt*-toxicity was evident after 30-days post-spray in the direct sun/rain treatment compared to the below-canopy shaded treatment in both years of this experiment (Figure 1 in Scriber and Gage 1995).

Different insect species

There were no (host \times *Bt* spray) interactions in the ANOVA for *C. promethea*, *P. glaucus*, or *P. canadensis* after 5 days of feeding (see above, and Johnson et al. 1995). The concentrations of *Bt* on the foliage were similar to aerial application rates, and there were no significant (detectable) differences as a result of phytochemical differences between tree species.

The reasons for this extended period of toxicity for these non-target *promethea* silk moths and swallowtail butterflies remained unknown. Based on toxicity to the targeted gypsy moth caterpillars, the conventional wisdom and impression given by the USDA Forest Service was that *Bt* sprays were basically safe for Lepidoptera after 1–2 or a few days since they putatively degraded or were washed away by rain. This clearly was incorrect, but the explanation of these differences in impact still eludes researchers.

Different dosage sensitivities as an explanation of long-term toxicity (?)

In order to see if our non-target Lepidoptera species were simply much more dose-sensitive than gypsy moth larvae to *Bt* formulations, we conducted some basic laboratory studies. In experiments with standard Btk formulations (Foray 48B and Dipel), we determined that the LD-50 for *P. canadensis* was 0.008 IU/cm²; *P. glaucus* was 0.0014–0.0030 IU/cm²; *P. palamedes* was 0.004 IU/cm²; and *P. troilus* was 0.007 IU/cm².




Figure 1. The relative growth rates ($\text{mg mg}^{-1} \text{d}^{-1}$) of final instar ($n = 9$ larvae per treatment/plot, for two field plots) *Papilio canadensis* larvae (TOP); and final instar hybrid backcross larvae (of a hybrid female, *P.g.* x *P.c.* mated to a *P. glaucus* male; BOTTOM) on *Populus* hybrid leaves subjected to four experimental field treatments gypsy moth defoliated (D) and non-defoliated (N), with fertilization (F) and unfertilized (UF). For the hybrid backcross larvae, the two defoliated plots were significantly poorer for larval growth than the two non-defoliated plots ($F = 42.6$, $p = 0.003$) but treatments were not significantly different not for the *P. canadensis*. On the Defoliated/Unfertilized leaves, 67% of the hybrid larvae died before pupation, although mortality was observed in any of the three other treatments. These bioassays were conducted in 1997, approximately 2 months after a severe experimental defoliation of *Populus* plots using nearly 10 million gypsy moth caterpillars (Scriber et al. 1999).

(Scriber 1998). These lethal doses (LD-50s) are all considerably less than the 30–43 IU/cm^2 actually deposited on forests in aerial sprays of *Bt* for gypsy moths and the average LD-50 of 2.7 IU/cm^2 (Radcliffe and Yendol 1993).

Although low-dose toxicity prevailed on black cherry (*Prunus serotina*) and tulip tree (*Liriodendron tulipifera*) leaves, we were able to detect different host plant-LD-50 dosage responses for second instar *P. glaucus* larvae. On tulip tree, 87–100% of all larvae died on leaves with all *Bt* doses (including; 0.0004, 0.0017, 0.0023, 0.0030, 0.004,

0.0070, 0.0170, 0.050 IU/cm^2), compared to only 14.6% mortality on the tulip tree leaf controls (96 larvae/dose x 8 doses; larvae from 24 different families; Scriber and Haas 1996). In contrast to this highly enhanced mortality on tulip tree leaves, larvae in the same experiment on black cherry (Foray 48B leaves dipped and maintained in individual petri dishes with a larva at 24 °C; 18:6 photoperiod; for 4-day survival) were much less sensitive (LD-50 was between 0.002 and 0.004 IU/cm^2 ; with only 19% mortality at 0.0005 and 9% on the control; Scriber and Haas 1996).

Potential synergism from phyllosphere (phylloplane) microbes as an explanation of long-term toxicity (?)

Toxicity for *Papilio* caused by such low doses, might partially explain the long-term persistence of *Bt* toxicity in our field studies. We examined the possibility that these low doses were also partially from a synergism resulting from other bacteria that were found naturally present on leaf surfaces used as natural food in our *Papilio* (e.g. *Serratia marcessans*, Leah Bauer, pers. comm.) during both in our field studies and in some lab studies. Previous research (Miyasono et al. 1994; Dubois and Dean 1995) had shown that at least 14 different bacterial strains (including those of *Klebsiella* sp., *Erwinia* sp., *Pseudomonas* sp., *Xanthomonas* sp., *Actinomyces* sp., *Corynebacterium* sp., *Flavobacterium* sp., and *Escherichia coli*) were of effective synergizers of at least one of the Cry IA *Bt* toxins. Unlike *Bt* spores which contain some protein toxins, in the absence of the CryIA toxins, none of these bacterial cell or spore synergists exhibited any toxicity or growth inhibition (Miyasono et al. 1999; Dubois and Dean 1995). Ulceration of the caterpillar midgut resulting from cell lysis and loss of peritrophic membrane integrity from *Bt* toxins are hypothesized to have facilitated this invasion of the hemolymph by bacterial opportunists. This type of interaction may, to varying extents, explain some of the laboratory mortality in insect culturing (whether on leaves or on synthetic diets, especially butterflies; Cohen 2003; see also Dillon and Dillon 2004). In our lab, it appears that hybrids between *P. canadensis* and *P. glaucus* may be better able than the parental species to survive the unavoidable lab-pathogens while we were using late summer tree leaves for larval rearing (Scriber et al. unpublished). Leaf sterilization with 5% bleach solutions gave some improvement in our rearing success, but this technique was also tedious and variable.

In an attempt to determine the possible synergistic effect of leaf surface microbes with different *Btk* application doses, Haas and Scriber (1998) used 4 commercial *Bt* (Foray 48B) doses (0.268, 0.034, 0.008, and 0.004 IU/cm²) and control leaves of tulip tree and black cherry with tiger swallowtail (*P. glaucus*) butterfly larvae. Each experiment included a pre-treatment leaf dip of 5% Clorox bleach followed by a 10-minute water rinse (with a

no-dip control leaf). The *Btk* dosage effects were clearly observed for all four treatments (cherry leaves with and without bleach treatment and tulip tree leaves with and without bleach). However, removal of the microbial community from the leaf surface did not decrease the *Bt* toxicity dose required to kill *Papilio*, as predicted if microbes synergize *Bt* toxicity (Dubois and Dean 1995).

In fact, the growth and survival of larvae were slightly poorer for the bleached leaves, perhaps reflecting inadvertent leaf-wax removal from the bleach treatment and subsequently faster leaf desiccation or an unintentional leached/reduced nutritional value. The actual reasons for poorer performance on our experimentally bleached leaves remains unknown, but we verified that very low doses of *Btk* will kill *Papilio*, even when not synergized by any leaf surface microbes. The LD-50 doses were between 0.0006 and 0.0025 IU/cm² for these four treatments (860 larvae from 58 different females evenly distributed across treatments; Haas and Scriber 1998). This extremely low dose corresponds to dosage results obtained in previous lab studies of four *Papilio* species (Scriber and Haas 1996) and are much lower (1000×) than the 2.7 IU dose required for gypsy moth larvae (Radcliffe and Yendol 1993; Reardon et al. 1994).

Natural enemies and slowed growth with sub-lethal exposure (?)

Reduced growth rates of pathogen infected (or *Bt*-sprayed) larvae can result in enhanced parasitism rates (Andreadis et al. 1983; Reardon et al. 1994; Peacock et al. 1998). The interaction of pathogens, parasites, and competitors has not been comprehensively researched for many herbivores (but see Denno et al. 1995). However, the possibility of direct competition from severe defoliators such as gypsy moths (or indirect competition from such defoliators via impacts on 'enemy-free space' or induced resistance in host trees), as a result of outbreak defoliations must be seriously considered for non-target impacts from non-spray management decisions (Scriber 1992; Redman and Scriber 2000). Some of the mortality in field studies above may have been partially related to some kind of different behavior of larvae feeding on *Bt*-sprayed trees that may have rendered them more vulnerable to predators or parasites because of slowed

growth rates (Reardon et al. 1994; Wraight et al. 2000; Zangerl et al. 2001).

Larval performance of Salicaceae-‘adapted’ (*P. canadensis*; Hwang and Lindroth 1995) and unadapted (*P. glaucus*; Lindroth et al. 1988) swallowtail butterflies is related to levels of phenolic glycosides in *Populus* leaves. One of these species (*Papilio canadensis* R & J) showed little reduction in larval growth rates or survival on any *Populus*, while the other (*P. glaucus* L.) found Salicaceae basically toxic for all larvae; Scriber et al. 1995a). In feeding studies using leaves sampled from these four treatments of defoliated/fertilized hybrid poplars we did see these all-or-none survival and growth patterns for larvae of each of the parental species (i.e. all *P. glaucus* died). However, significant leaf suitability differences were detected among the four treatments (defoliated, fertilized) and reflected in growth rates of hybrid and back-cross larvae (Scriber et al. 1999, Figure 1). As with other field studies of defoliated and re-leafed trees (Redman and Scriber 2000), we found no differ-

ences in ability of adult females of either species to detect these phytochemically and nutritionally induced leaf treatment differences (Figure 2a, b, below).

Other possibilities remain unknown (vegetative growth and resproutation on leaves?)

The extent to which *B. thuringiensis* reproduces and increases on leaf surfaces remains basically unknown, although it is known to persist on leaves for long periods of time (Leong et al. 1980; Smith and Couche 1991) and may have impacts even if it doesn’t increase vegetatively with resproutation and increased concentrations of toxins. The undersides of leaves may protect some spores from sun light and rain. The pollen, dust, insect frass, and aphid honeydew exudates tend to accumulate on tree leaves later in the summer, and might provide abundant nutrients for leaf surface microbes such as *Bacillus* and others.




Figure 2. The percentage of *Papilio canadensis* eggs (TOP) laid on leaves of four different hybrid poplar treatment leaves (gypsy moth defoliated/ fertilized, GM+ F+; and GM+ F-, GM- F+, GM- F-, respectively) in early September 1997. No significant differences were observed for *P. canadensis* female choices (Chi-square $p > 0.90$; 16 females). There were no significant differences in the very small percentages of *P. glaucus* eggs (BOTTOM) on leaves from 4 *Populus* treatments with a normal tulip tree host in 5-choice arena (no eggs were laid by any female in arenas with only the 4 *Populus* leaves; $n = 28$ females in 1997, 44 females in 1998).

Indirect impacts causing non-target effects (Decision for no *Btk* spray)

Biological control of gypsy moths with natural and introduced parasites, predators and pathogens (fungi, bacteria, viruses, and microsporidians) have been shown to have variable non-target impacts on Lepidopteran species. When gypsy moth populations increase to outbreak levels, associated natural enemies abound that can attack native non-target Lepidoptera (Elkinton and Liebold 1990; Elkinton et al. 1990; Gould et al. 1990; Ferguson et al. 1994; Redman and Scriber 2000). This is especially true of generalist parasitoids such as the tachinid fly *Compsilura concinnata* (With at least three generations/year) that kills more than 200 species of butterflies and moths (Culver 1919; Webber and Schaffner 1926; Williams et al. 1992; Scriber 1998). It has been estimated that 68%–81% of two large silkworm species (*Callosamia promethea* and *Hyalophora cecropia*; Boettner et al. 2000) and up to 62% of *Actias luna* moth larvae (Kellogg et al. 2003) are killed by *Compsilura*. However, while these species have persisted even after 100 years of parasitism by the introduced *Compsilura*, other Lepidoptera have apparently been extirpated in New England (Schweitzer 1988;

Louda et al. 2003). In addition to these direct attacks by introduced and natural parasites, the vulnerability to introduced and natural pathogens (fungi, microsporidians, and viruses) is an issue for non-target Lepidopterans in the same community as defoliating herbivores (Elkinton et al. 1991; Hajec et al. 1995; Wesoloh and Andreadis 1997; Redman and Scriber 2000). Direct competition may also occur between phytophagous insects and these targeted defoliator pests (Scriber 1992, 1998; Stewart 1996). The costs of evaluating non-target impacts of potential biological control introductions (which can evolve and change in time; Louda et al. 1977) would be huge and unrealistic (Hawkins and Marino 1997), yet seem important enough to deserve serious study (Stiling and Simberloff 2000). An excellent way to start might include sentinel species that are large, well-known to the public, and common (Stamp 1990; Bossart and Carlton 2002) much as described by Boettner et al. (2000) and (Kellogg et al. 2003)

Direct competition of forest defoliators with native non-target species of Lepidoptera

Northern Michigan suffered severe and extensive defoliations from gypsy moths and forest tent caterpillars during the 1980's and early 1990's (Gage et al. 1990; Gage 1996; Sharov et al. 1998 and 1999). Few tree leaves (other than Maples, *Acer*, or ashes, *Fraxinus*) could be found in the heavily defoliated outbreak areas. In addition to these outbreak caterpillars eating leaves with *Papilio canadensis* eggs (as a direct mortality factor; Scriber 1996), there was a serious lack of host leaves of *Papilio* hosts. Trees such as paper birch and other birch/alders (*Betula* spp./*Alnus* spp.), quaking aspen, balsam poplars, big-toothed aspens, and willows (*Populus* spp./*Salix* spp.), shadbush (*Amelanchier* spp.), pin cherry, choke cherry, black cherry (*Prunus* spp.), basswood (*Tilia americana*), and others were extensively, if not completely, defoliated (Scriber 1992). Such severe and widespread defoliation (millions of hectares) certainly posed potential competitive threats for *Papilio* populations and other species of Lepidoptera in these forests (Scriber and Gage 1995). However, determining the precise role of direct competition for phytophagous insect herbivores is difficult for generalist species such as these

P. glaucus and *P. canadensis* (local specialists on a gypsy moth devoured tree species may have been more seriously devastated).

Indirect competition with native non-targets (enemy-free space)

Increased densities of gypsy moths attract natural enemies of all sorts, which impact other Lepidoptera locally in an 'indirect competition' (Elkinton and Liebold 1990; Elkinton et al. 1990; Gould et al. 1990; Ferguson et al. 1994). Similar patterns of natural enemy increases have been observed in the wake of forest tent caterpillar outbreaks as well (Parry et al. 1997). It was determined that swallowtail butterflies in northern Michigan were negatively affected by gypsy moths in several ways, including loss of usable host foliage (see above; Arteel and Lindroth 1992), depressed growth rates and survival on regrowth leaves (see below), and increased rates of parasitism when placed near gypsy moth infestations (Williams et al. 1992; Redman and Scriber 2000), or even when artificially high numbers were created (Boettner et al. 2000). Some evidence was observed that suggests that gypsy moth body fluids with NPV virus naturally or experimentally painted on plant leaves could also reduce growth and survival of *Papilio canadensis* larvae (Redman and Scriber 2000).

It is unknown whether insect hybrids might survive disease better than parental species but this may be the case with our *Papilio*. Much less is known about parasites or diseases of hybrid animals than parasites (e.g. phytophagous insects) and disease of hybrid plants (Moullia 1999; Whitham et al. 1994, 1999). However, increased resistance of animal hybrids have been reported (e.g. hybrid pocket gopher resistance to lice; Heany and Timm 1985; Hafner et al. 1998). In some cases F-1 primary hybrids are more resistant (express a hybrid vigor) than the parental species, but the backcrossed and recombinant genotypes may be more susceptible due to the breakup of co-adapted gene complexes of the immune response system (Moullia et al. 1995, 1996). Few studies of hybrid insect resistance to disease or parasites exist, but hybrid honey bees (of European and African subspecies) may have greater resistance to some parasites such as *Varroa* or tracheal mites (Nasr 1997; Erickson 1998; Sammataro et al.

2000). However, no difference in fungal pathogen susceptibility/resistance was found in the hybridizing cicada species (Duke et al. 2002).

Our hybrids between *P. canadensis* and *P. glaucus* show higher survival and growth rates than parental species (Donovan 2001; Scriber et al. 2003). We do not rule out the possibility that this was perhaps due to better survival of hybrids on pathogen-laden field-collected leaves in our lab rearing process. Extensive interspecific natural gene introgression and recombination has been extensive across the historical hybrid zone between these *Papilio* since the 1998 regional climate warming in North America (Scriber 2002a, 2002b). Differentially sensitive genotypes spawned by introgression might survive in local pockets of high pathogen concentrations. The differential susceptibility/resistance of non-target species and their hybrids has not been investigated for any insect pathogen (bacterial, fungal, microsporidian, nematode, or virus). Surprisingly, the ability of female insects to avoid mating with infected males has been reported for some insects (English 2001) as in mammals (Kavaliers and Colwell 1995). Such differential mating behavior across the *Papilio* hybrid zone (see Deering and Scriber 2002) could indirectly affect the survival of local populations (or particular genotypes) with exposure to pathogens/parasites in defoliation outbreak areas (Dwyer et al. 2004).

*Indirect competition with native non-targets
(induced phytochemical resistance in previously
defoliated plants)*

In recent years, we have seen the natural spread of gypsy moths into southwestern Michigan Long Term Ecological Research (LTER) poplar plots. This was an excellent position from which to document effects on tree physiology, allelochemicals, insect population dynamics and nutrient cycling under natural infestations with simultaneous experimental manipulations of the hybrid poplar plantations of the Kellogg Biological Station. In 8 Lepidopteran species of *Populus*-feeders on experimentally defoliated (with and without Nitrogen fertilization) and undefoliated (with and without Nitrogen), it was found that catastrophic (near-total) experimental defoliations did result in

altered leaf quality and slowed growth or smaller pupae or reduced fecundity for subsequent phytophagous insects in the same year, and in 2–3 years post-defoliation (Scriber et al. 1999; Parry 2000). The variation we observed between defoliation and fertilization treatments may have related to the structure of the particular tannins (Ayres et al. 1997) or the biochemistry of different herbivore guts (Barbehenn and Martin 1994) as well as the availability of carbon and/or nitrogen to the plants (Herms and Mattson 1992; Arnold and Schultz 2002; Lovett et al. 2002). In *Populus tremuloides*, experimentally defoliated using *Malacosoma disstria*, previous years of defoliations did not affect the parasitoids (Parry et al. 2003) although other herbivores were affected by experimental defoliation of this aspen species (Dankert et al. 1997). Experimental defoliation of birch and sugar maples showed interactions of sunlight/shade and nitrogen fertilizer on leaf quality for other Lepidoptera (Govenor 1998). Meta-analyses of 68 studies of damage-induced changes in woody plants show that herbivore responses to defoliation depend upon the plant species, the type of damage, and the timing of damage (Nykanen and Koricheva 2004).

In order to determine the extent of potential impact of *Populus* defoliations and potential phytochemical induction (Lindroth 1992; Bryant et al. 1993; Hunter and Schultz 1995; Lindroth and Hwang 1996; Dankert et al. 1997; Wait et al. 1998; Peters and Constabel 2002) on genotypes of generalist herbivores, we created interspecific *Papilio* hybrids and backcrosses between adapted and unadapted species using hand-paired lab crosses (Scriber et al. 1999). In attempts to simulate a defoliating insect outbreak in experimental forests of genetically identical hybrid poplar trees (540/plot within larger plots at the LTER Long term 'agroecology' research site at Kellogg Biological Station in Michigan), two million gypsy moth eggs used in 1996 achieved 30–40% defoliation. Essentially total defoliation was achieved in 1997 plots by allowing nearly 10 million hatching gypsy moths to ascend from containers tied to the individual tree boles. All gypsy moth egg masses collected in the winter of 1996–1997 were surface sterilized (10% formalin) to kill virus. The highest feeding activity and defoliation rate coincided with the later instars and occurred in late June. Our feeding bioassay experiments with *Papilio* species,

hybrids, and backcrosses began July 3rd and ran through late August in 1997 and 1998 using the regrowth leaves from the four treatments of LTER treatment plots number 1 and 2. Oviposition studies were conducted on leaves from plots 1 and 2 in 1997 and 1998.

Female *Papilio glaucus* were, in both 1997 and 1998, directly obtained from one population in southeastern United States (Clarke Co. GA), and *P. canadensis* females from Northern Michigan (Emmet, Cheboygan, and Charlevoix counties). Females were individually placed into clear round plastic containers (10 cm high 25 cm diameter) with *Populus* leaves of respective treatments (+ +, + −, − +, − −) for defoliation and fertilization, respectively. These multi-choice oviposition arenas were stacked on rotating platforms aligned in front of a bank of 100 W incandescent bulbs on a 4 h on/off cycle. The rotation of containers was set so that each of the leaves (supported in a water-filled florists' aquapic) passed in front of the light 10 times per hour. Since the butterflies flutter and bounce along inside of the dishes at the side facing the lights, the multi-choice arena provided a slowly moving sequence of leaves to each female. The treatments were randomized each time they were replaced. Eggs were removed from each container and counted daily (see Scriber 1993 for details of the oviposition arenas). Females were fed daily with a 20% honey water solution.

The leaves bioassayed in this study came from the four treatments of two large plots in the LTER site at the Kellogg Biological Station. Upper crown branch samples were taken from 14 trees per plot with pruning poles above 2.5 m June 28 1997 and August 4 1997. Leaves were immediately flash frozen in liquid nitrogen. Methanol was used to extract phenolic compounds. Total phenolics were quantified using the Folin-Dennis assay on a Rapid-Flow autoanalyzer. Condensed tannins were quantified using a modified sulfuric acid procedure (Nitao et al. unpublished) on the same equipment. Leaf water content was determined for each treatment from subsamples taken at the time of experimental set-up of field-sampled leaves brought to the lab on ice in ziplock bags. Leaf nitrogen (1997) was determined on a dry weight basis with an autoanalyzer. Leaf water was taken at several additional sample dates. Difference between leaf treatment means for leaf water, nitrogen and total phenolics after ANOVA (using GLM procedures) were determined as Least Significant Differences (LSD; SAS 1989; Table 1).

The Salicaceae-adapted species *P. canadensis* did well on all treatment leaves (after defoliation and fertilization treatments) and the unadapted species, *P. glaucus*, did poorly on all leaves. Hybrids and backcrosses with intermediate levels of detoxification carboxylesterase enzymes (Scriber et al. 1989) were intermediate and showed reduced growth on the gypsy moth defoliated

Table 1. Leaf nutritional quality assessments (Mean of two plots)

Plot treatments	Leaf water ^A (%)			% Total nitrogen ^B (dry weight)		Total Foliar phenolics ^C (%)
	July 3	July 18	August 18	mid-June	mid-August	Mid-August
Defoliated/N+	80.5 a ± 0.8	73.0 ab ± 1.3	71.2 b ± 0.4	2.14 b ± 0.09	2.96 a ± 0.34	5.95 a ± 0.35
Defoliated/N−	79.6 a ± 2.8	73.5 ab ± 0.9	71.4 b ± 0.5	2.29 ab ± 0.07	2.40 a ± 0.12	5.65 a ± 0.05
Undefoliated/N+	71.4 b ± 0.1	72.7 b ± 0.5	70.2 b ± 0.4	2.75 a ± 0.03	2.89 a ± 0.10	7.10 b ± 0.20
Undefoliated/N−	70.5 b ± 2.7	72.2 b ± 1.6	70.5 b ± 0.0	2.54 ab ± 0.23	2.75 a ± 0.24	7.05 b ± 0.25

Budbreak in hybrid poplars late May-early April; experimental gypsy moth defoliation mid-June to late June; Releafing late-June to July 5th. Larvae of hybrid, backcross, and parental *Papilio* genotypes were bioassayed mid-July to mid-August. Females (second generation) were bioassayed for oviposition in mid- to late-August.

^A Leaf water means of 2 plots (24 individual trees) by date 1997.

^B Nitrogen means of 2 plots ($n = 55$ individual trees were sampled in Plot 1 and $n = 58$ trees from Plot 2) 1997.

^C Total phenolics six to seven weeks post-defoliation were determined on August 11 1997 samples (see Scriber et al. 1999).

ANOVA (GLM, SAS) conducted on plot means significant differences between means for: overall leaf water, June nitrogen and phenolics.

treatments, presumably due to the induced changes in allelochemicals such as specific phenolic glycosides (Lindroth et al. 1988) and nutrients such as lowered nitrogen concentrations (water content can be a very important nutrient, but was not lower in any of these 4 hybrid poplar treatment leaves; see Table 1). The total phenolics in both of the defoliated treatments were actually lower than the undefoliated treatments 2 months after the 1997 defoliation (Table 1) but rose in the subsequent years where they had minimal effect on herbivores (Parry 2000; Parry et al. 2003). It has been suggested that phenolics may perhaps be involved in other plant protection functions (e.g. from photodamage, not herbivores; Matsuki 1996; Close and McArthur 2002). It has also been found that with *Populus deltoides* Nitrogen fertilization changes leaf chemistry (including phenolics, leaf N, salicin, total soluble protein, etc.) via changes caused in leaf development (Wait et al. 1998).

Our hybrid/backcross *Papilio* genotypes had intermediate levels of esterase detoxification enzymes (Scriber et al. 1989) and were intermediately sensitive (compared to the parental species) to the phytochemically altered treatment differences in gypsy-moth-defoliated hybrid poplar regrowth leaves (*Populus* x *euoamericana* c.v. 'Eugenii'; *P. deltoides* x *P. nigra*), showing more growth rate and survival differences (Figure 1; 1999) than 5 other adapted species, including forest tent caterpillars, tussock moths, poplar tent maker, big poplar sphinx, and Fall webworms, as well as the gypsy moths themselves (Parry 2000). The phytochemically induced differences in *Populus* leaves that affected *Papilio* larvae did not differentially affect the ovipositing females of either *P. canadensis* or *P. glaucus* (Figure 2), which was also observed for poor quality aspen leaves in northern Michigan after gypsy moth outbreaks (Redman and Scriber 2000). Such subtle differences in the best and worse hosts were not detected or recognized yet for their physiological/ecological impacts, as is the case for other Lepidopteran 'oviposition mistakes' on toxic plants that may have been recently introduced or phytochemically altered (Scriber 1993; Renwick and Chew 1994; Mercader and Scriber 2004). Plant hybrids can serve as bridges or barriers to insect herbivores of various guilds (Whitham et al. 1999).

Genetically engineered (transgenic) crops and forest trees with Bt endotoxin expression in tissues

Transgenic crops occupy more than 40 million hectares on a global scale, and in the past 15 years, more than 100 species of genetically transformed plant species have been produced (Babu et al. 2003). Recently, publicity regarding transgenic (genetically modified) plants has raised levels of concern regarding non-target and other evolutionary impacts (Ferber 1999; van Emden 1999; Pimentel and Raven 2000; Hails 2000; Kanda et al. 2000; Brower 2001; Scriber 2001). The development of a new generation of *Bt*-crops with much higher toxin expression levels (Kota et al. 1999) may heighten concerns. The expression of *Bt* endotoxins in *Zea mays* (corn) pollen that killed monarch butterfly larvae (Losey et al. 1999; Hansen-Jesse and Obrycki 2000; Sears et al. 2001) led to a major environmental impact debate (Ferber 1999; Brower 2001). Subsequent research showed that other butterflies such as *Papilio* species were affected by Bt toxins in pollen only at very high concentrations, however sub-lethal effects were evident (Wraight et al. 2000; Scriber 2001; Zangerl et al. 2001). The results of a series of research studies presented in a special issue of the Proceedings of the National Academy of Sciences in the USA alleviated much of the concern about widespread movement and severe toxic impacts of transgenic corn pollen (which is heavy and thus primarily of only local concern; Pleasants et al. 2001; Zangerl et al. 2001). In addition to pollen, plant anther parts can disperse and contaminate hosts of non-target Lepidoptera such as monarch butterflies on milkweed plants (Asclepiadaceae; Hanson-Jesse and Obrycki 2000; Hellmich et al. 2001). The non-target impacts of *Bt*-corn pollen on the monarch butterfly in North America were generally low, but locally could be severe (Brower 2001; Scriber 2001).

Since most of the focus and research on non-target impacts of transgenic corn pollen was concentrated on the 'in-field' weed-feeding species of Lepidoptera (monarch butterflies and black swallowtails; Oberhauser et al. 2001; Zangerl et al. 2001), we felt it essential to evaluate the potential impact of transgenic corn pollen on two species of *Papilio* that utilize host plants along hedgerows and field edges throughout the Midwest region of North America (below). Similar non-target trans-

genic plant issues may emerge for trees (Raffa 1989; Bauer 1997), and the negative impacts may extend through herbivores that feed on these plants to include the natural enemies that eat them (Chapman and Hoy 1991; Hilbeck et al. 1998). Another concern is that some plant parts (e.g. fruits) expressing Cry3B toxins aimed at Coleoptera may nonetheless affect Lepidoptera (Arpaia 1997). Unlike *Bt* spores and crystals exposed on leaf surfaces from sprays, *Bt*-toxins when produced in plant cells (or in the midgut of herbivorous insects) are not exposed to sunlight and may remain active to subsequently affect herbivores or their entomophagous natural enemies (De Maagd et al. 2001; Groot and Dicke 2002). Toxins from *Bt*-plants (root exudates or other decaying biomass) also bind to the soil and may persist for weeks or months and remain toxic to insects (but not to many other invertebrates; Tapp and Stotzky 1998; Saxena and Stotzky 2001). Unfortunately, the natural ecological interactions of *B. thuringiensis* remains basically unknown (Addison 1993; Groot and Dicke 2002).

Transgenic Btk corn pollen does negatively impact North American non-target Lepidoptera (Papilionidae), but so does non-Btk corn pollen!

We determined that transgenic corn pollen that was genetically engineered to express the *Bacillus thuringiensis* var. *kurstaki* toxin against the European corn borer, *Ostrinia nubilalis*, suppressed growth and survival of two North American non-target species (*Papilio glaucus* and *P. troilus*) when dusted on their favorite host tree leaves (Tables 1 and 2). Corn pollen from the same maize genotype, but lacking the *Btk* toxin, also suppressed growth and survival of the two *Papilio* species at basically the same levels as the pollen with the *Btk* toxin (growth rates, RGR's, were approximately half that of larvae fed control leaves with no pollen). While these antibiotic effects in no-choice feeding arenas were clear with pollen dusted on tree leaves at approximately 10% fresh leaf weight equivalents; similar effects were observed at very light dustings (1% fresh leaf equivalents). While significant mortality of both *Papilio* species continued during days 3–7 on fresh pollen-free host leaves, survivors recovered and grew at increased rates that were still slightly less than the control

Table 2. Larval growth performances (2 day and 7 day) of *Papilio glaucus* fed untreated *Liriodendron* leaves and leaves dusted with transgenic (*Btk* toxins) or non-transgenic (non-*Btk*) corn pollen. Data are expressed as a mean (\pm SE)

	Initial larvae (<i>n</i>)	48-h survival (%)	Relative ^a growth rate (RGR)	Relative ^a consumpt. rate (RCR)	Approx. digest. (AD)	Efficiency (%) of food conversion		7-day surv. (%)	7-day RGR ^a
						Digested (ECD)	Ingested (ECI)		
Control leaves	(9)	100	0.297a \pm 0.056	2.545 \pm 0.309	51.4 \pm 5.7	23.6ab \pm 6.2	11.7a \pm 1.7	72.2	0.722a \pm 0.036
Btk pollen (10%)	(17)	83.1	0.152b \pm 0.027	2.123 \pm 0.398	45.8 \pm 6.3	43.3a \pm 9.5	7.2b \pm 1.1	37.5	0.400b \pm 0.066
Non-Btk pollen (10%)	(16)	88.2	0.150b \pm 0.036	2.985 \pm 0.282	54.9 \pm 4.5	11.6b \pm 3.1	4.9c \pm 1.0	23.5	0.420b \pm 0.089

^aRGR and RCR (mg mg⁻¹ d⁻¹). Significant differences indicated ($p = 0.05$).

larvae that were not exposed to either pollen type during the 48-h experimental period.

In addition to the recovery of some larvae that consumed leaves with pollen in no-choice studies (Table 3), we observed strong larval feeding deterrence in 2-choice 24-h preference studies in which half of each of the leaves was dusted with pollen (10% fresh weight equivalents Table 4). Larval feeding was basically confined to the non-dusted half of tulip tree leaves (*Liriodendron tulipifera*) for *P. glaucus*. Their growth rates and survival were significantly better than *P. troilus* larvae which apparently failed to discriminate between the treatments (they ate as much of the pollen-covered half of spicebush leaves, *Lindera benzoin*, as the untreated control half).

We conclude that leaves in forest edges or hedgerows adjacent to cornfields may become unsuitable for non-target Lepidoptera species if pollen were to accumulate on the leaf surfaces. However, survival capabilities of larger larvae with strong non-preference in larvae feeding of some species such as *P. glaucus* should allow survival via a change of larval feeding location on the tree. We still do not know what the impact of corn pollen would be for neonate larvae, nor whether ovipositing females would avoid pollen-dusted leaves.

In summary, no pest management tactic is without risk to non-target organisms (Wolfenbarger and Phifer 2000). Generally, the *Bt* pesticide sprays and transgenic plants are of relatively less risk than other insecticides. The natural and altered microbial communities on the leaf phylloplane remain as a black box whose interactions may account for much of the unknown variance in herbivore-plant interactions. Natural interactions among different bacteria synergize the toxicity of *Bacillus thuringiensis* against Lepidoptera 100-fold to 1000-fold (Dubois and Dean 1995), as is the case for other microbial interactions (e.g. virus and microsporidians; Bauer et al. 1998). In defoliator outbreak situations, not using pesticides may in many ways be even worse for some non-target Lepidoptera due to high levels of community or pest associated pathogens and/or parasitoids. With regard to *Bt*, even generalized butterflies may be more sensitive (than moths) and could serve as non-target impact monitoring or indicator species.

Table 3. Larval growth performances (2 day and 7 day) of *Papilio troilus* fed untreated leaves of spicebush (*Lindera benzoin*) and leaves dusted with corn pollen from transgenic (Btk toxins) and non-transgenic (non-Bt) plants

	Initial larvae (n)	48-h survival (%)	Relative ^a growth rate (RGR)	Relative ^a consumption rate (RCR)	Approx. digest. (AD)	Efficiency (%) of food conversion		7-day surv. (%)	7-day RGR ^a
						Digested (ECD)	Ingested (ECI)		
Control leaves	(5)	100.0	0.304 a ± 0.070	3.816 ± 0.491	61.4 ± 5.3	16.8 ± 6.7	9.9 ± 4.3	100.0	0.942 a ± 0.006
Btk pollen (10%)	(12)	81.1	0.148 b ± 0.034	4.381 ± 0.983	67.8 ± 4.4	8.7 ± 3.1	5.8 ± 2.1	45.5	0.763ab ± 0.053
Non-Btk pollen (10%)	(11)	83.3	0.136 b ± 0.024	3.869 ± 0.745	54.9 ± 8.8	18.5 ± 9.6	4.0 ± 0.8	58.3	0.588 b ± 0.056

Data are presented as a mean (±SE).

^aRGR and RCR (mg mg⁻¹ d⁻¹). Significant differences are indicated (*p* = 0.05).

Table 4. Larval growth rates (of survivors) at 24 h and 6 days^a for *P. glaucus* fed tulip tree leaves (and *P. troilus* fed spicebush) half of each leaf was dusted with pollen (Bt or non-Bt) at 10% fresh weight equivalents on one half of the leaf

	No. of initial larvae (n)	24 h surv. (%)	24 h RGR	6 day surv. (%)	Visual estimates pollen	Visual estimates control	Calculated leaf area pollen	Calculated leaf area control
<i>Glaucus</i>								
Bt pollen	(8)	87.5	0.179 ± 0.047	62.5	4.4 ± 2.6	23.8 ± 5.9	3.4 ± 2.2	23.5 ± 5.0
Non-Bt pollen	(7)	71.4	0.161 ± 0.050	42.9	4.6 ± 2.9	21.4 ± 8.6	3.8 ± 2.7	20.9 ± 7.9
<i>Troilus</i>								
Bt pollen	(5)	40.0	0.108 ± 0.018	0.0	18.2 ± 16.2	19.0 ± 4.0	21.4 ± 7.9	22.2 ± 4.6
Non-Bt pollen	(5)	40.0	.029 ± 0.009	20.0	9.0 ± 9.6	10.0 ± 4.5	11.8 ± 6.2	13.4 ± 4.7

At right, the relative amounts of leaf consumption for pollen dusted (at 10% leaf fresh weight equivalents) versus undusted control halves of tulip tree leaves by *Papilio glaucus* and spicebush leaves by *P. troilus*.

Acknowledgements

Research on non-target impacts of *Btk* and alternative decisions in forest and field was supported by the College of Natural Sciences, the Michigan Agricultural and Natural Resources Experiment Station (Project MICL#1644), the Michigan Dept. of Agriculture, the USDA Competitive Grants Program, the NC Regional Research Program (NC-212), and the National Science Foundation the National Science Foundation (LTER BSR-8702332; Ecology DEB-92201122; Ecosystems DEB-9510044; Population Biology DEB-9981608). NSF REU awards to Kirsten Weir, Will Sands, and Matt Lehnert helped with aspects of the gypsy moth defoliation and field work. The following people helped with the experiments and surveys in the lab and/or field; Leah Bauer, George Bird, Renee Brenner, Terry Davis, Mark Deering, Jessica Deering, Jen Donovan, Piera Giroux, Laura Haas, Holly Hereau, Dan Herms, Geoff Hoffa, Kelly Johnson, Keegan Keefover, Kevin Kosola, Bob Lederhouse, Matt Lehnert, James Maudsley, Ana de Morais, Jenny Muelhaus, James Nitao, Michelle Oberlin, Gabe Ording, Laura Palombi, Dylan Parry, Ahnya Redman, Christine Schoen, Dave Smitley, Doozie Snider, Lee Solter, Aram Stump, Suzanne Thiem, Patti Valella, and Wayne Wehling.

References

Addison J.A. 1993. Persistence and non-target effects of *Bacillus thuringiensis* in soil: a review. Can. J. For. Res. 23: 2329–2342.

- Altmann M. 1992. Biopesticides turning into new pests? Trends Ecol. Evol. 7: 65.
- Andreadis T.G., Dubois N.R., Moore R.E. B., Anderson J.F. and Lewis F.B. 1983. Single applications of high concentrations of *Bacillus thuringiensis* for control of gypsy moth populations and their impact on parasitism and disease. J. Econ. Entomol. 76: 1417–1422.
- Andrews J.H. and Hirano S.S. 1991. Phyllosphere Ecology: Microbial Ecology of Leaves. Springer, NY.
- Appel H. 1994. The chewing herbivore gut lumen: physiochemical conditions and their impact on plant nutrients, allelochemicals, and insect pathogens. In: Bernays E.A. (ed.), Insect–Plant Interactions, Vol. 5, CRC Press, Boca Raton, pp. 209–223.
- Arnold T.M. and Schultz J.C. 2002. Induced sink strength as a prerequisite for induced tannin biosynthesis in developing leaves of *Populus*. Oecologia 130: 585–593.
- Arpaia S. 1997. Ecological impact of Bt-transgenic plants: 1. Assessing possible effects of CryIIIB toxin on honey bees (*Apis mellifera* L.) colonies. J. Genet. Breed. 50: 315–319.
- Arteel G.F. and Lindroth R.L. 1992. Effects of aspen phenolic glycosides on gypsy moth (Lepidoptera: Lymantriidae) susceptibility to *Bacillus thuringiensis*. Gr. Lakes Entomol. 25: 239–244.
- Ayres M.P., Clausen T.P., MacLean S.F., Redman A.M. and Reichardt P.B. 1997. Diversity of structure and antiherbivore activity in condensed tannins. Ecology 78: 1696–1712.
- Babu R.M., Sajeena A., Seetharaman K. and Reddy M.S. 2003. Advances in Genetically Engineered (Transgenic) Plants in Pest Management—An Overview. Crop Protection 22: 1071–1086.
- Baldwin I.T. 1994. Chemical changes rapidly induced by folivory. In: Bernays E.A. (ed.), Insect–Plant Interactions. CRC Press, Boca Raton, FL, pp. 1–23.
- Barbehenn R.V. and Martin M.M. 1994. Tannin sensitivity of larvae of *Malacosoma disstria* (Lepidoptera): roles of the peritrophic envelope and midgut oxidation. J. Chem. Ecol. 8: 1985–2001.
- Barry J.W. et al. 1993. Predicting and measuring drift of *Bacillus thuringiensis* sprays. Environ. Toxicol. Chem. 12: 1977–1989.
- Bauer L.S. 1997. Fiber farming with insecticidal trees. J. Forestry 95: 20–23.

- Bauer L.S., Miller D.L., Maddox J.V. and McManus M.L. 1998. Interactions between a *Nosema* sp. (Microspora: Nosematidae) and nuclear polyhedrosis virus infecting the gypsy moth, *Lymantria dispar* (Lepidoptera: Lymantriidae). *J. Invertebr. Pathol.* 74: 147–153.
- Beckwith R.C. and Stelzer M.J. 1987. Persistence of *Bacillus thuringiensis* in two formulations applied by helicopter against the western spruce budworm (Lepidoptera: Tortricidae) in north central Oregon. *J. Econ. Entomol.* 80: 204–207.
- Beegle C.C. and Yamamoto T. 1992. Invitational paper (C.P. Alexander Fund): History of *Bacillus thuringiensis* Berliner research and development. *Canad. Entomol.* 124: 587–616.
- Beegle C.C., Kulmage H.T., Wolfenbarger D.A. and Martinez E. 1981. Persistence of *Bacillus thuringiensis* Berliner insecticidal activity on cotton foliage. *Environ. Entomol.* 10: 400–401.
- Boettner G.H., Elkinton J.S. and Boettner C.A. 2000. Effects of a biological control introduction on three nontarget native species of saturniid moths. *Conserv. Biol.* 14: 1798–1806.
- Bossart J.L. and Carlton C.E. 2002. Insect conservation in America: status and perspectives. *Am. Entomol.* 48: 82–93.
- Boulton T.J. 1999. Impacts of *Bacillus thuringiensis* subsp. *kurstaki* (Btk) on the population dynamics of non-target Lepidoptera on *Ribes cereum*. M.Sc. Thesis, Dept. Biol. Univ., Victoria, 291pp.
- Brower L.P. 2001. Canary in the cornfield: the monarch butterfly and the Bt corn controversy. *Orion* (Spring): 32–41.
- Brown K.S. 1997. Diversity, disturbance and sustainable use of neotropical forests: insects as indicators for conservation monitoring. *J. Insect Conserv.* 1: 25–42.
- Bryant J.E. and Yendol W.G. 1988. Evaluation of the influence of droplet size and density of Bt against gypsy moth larvae. *J. Econ. Entomol.* 81: 130–134.
- Bryant J.P., Clausen T.P., Reichardt P.B., McCarthy M.C. and Werner R.A. 1987. Effect of nitrogen fertilization upon the secondary chemistry and nutritional value of quaking aspen (*Populus tremuloides* Michx.) leaves for the large aspen tortrix (*Choristoneura conflictana* (Walker. *Oecologia* 73: 513–517.
- Bryant J.P., Reichardt P.B., Clausen T.P. and Werner R.A. 1993. Effects of mineral nutrition on delayed inducible resistance in Alaska paper birch. *Ecology* 74: 2072–2084.
- Burgess H.D., Thomson E.M. and Latchford R.A. 1976. Importance of spores and a-toxin protein crystals of *Bacillus thuringiensis* in *Galleria mellonella*. *J. Invert. Pathol.* 27: 87–94.
- Chapman M.H. and Hoy M.A. 1991. Relative toxicity of *Bacillus thuringiensis* var. *tenebrionis* to the two-spotted spider mite (*Tetranychus urticae* Koch) and its predator *Meta-seiulus occidentalis* (Nesbitt) (Acari: Tetranychidae and Phytoseiidae). *J. Appl. Entom.* 111: 147–154.
- Chenot A.B. and Raffa K.F. 1998. Effects of parasitoid strains and host instar on the interaction of *Bacillus thuringiensis* subspecies *kurstaki* with the gypsy moth (Lepidoptera: Lymantriidae) larval parasitoid *Cotesia melanoscela* (Hymenoptera: Braconidae). *Environ. Entomol.* 27: 137–147.
- Close D.C. and McArthur C. 2002. Rethinking the role of many plant phenolics—protection from photodamage not herbivores? *Oikos* 99: 166–172.
- Cohen A.C. 2003. *Insect Diets: Science and Technology*. CRC Press, Boca Raton, FL, pp. 324.
- Conner A.J., Glare T.R. and Nap, J.-P. 2003. The release of genetically modified crops into the environment. Part II. Overview of ecological risk assessment. *Plant J.* 33: 19–46.
- Culver J.J. 1919. A study of *Compsilura concinnata*, an imported tachinid parasite of the gypsy moth and the brown-tailed moth. USDA Bulletin #766.
- Dankert B.A., Herms D.A., Parry D., Scriber J.M. and Haas L.A. 1997. Mediation of competition between folivores through defoliated-induced changes in host quality. In: Raman A. (ed.), *Ecology and Evolution of Plant-Feeding Insects in Natural and Man-Made Environments*. Backhuys Publ, Netherlands, Leiden, pp. 71–88.
- de Barjac H. and Frachon E. 1990. Classification of *Bacillus thuringiensis* strains. *Entomophaga* 35: 233–240.
- De Maagd R.A., Bravo A. and Crickmore N. 2001. How *Bacillus thuringiensis* has evolved specific toxins to colonize the insect world. *Trend Genet.* 17: 193–197.
- Denno R.F., McClure M.S. and Ott J.R. 1995. Interspecific interactions in phytophagous insects: competition reexamined and resurrected. *Annu. Rev. Entomol.* 40: 297–331.
- Dillion R.J. and Dillon V.M. 2004. The gut bacteria of insects: nonpathogenic interactions. *Annu. Rev. Entomol.* 49: 71–92.
- Doane C.C. 1970. Primary pathogens and their role in the development of an epizootic in the gypsy moth. *J. Invertebr. Pathol.* 15: 21–33.
- Donovan J. 2001. Are hybrids more fit than their parental types A test using two swallowtail butterfly species *Papilio glaucus* and *P. canadensis* (Lepidoptera: Papilionidae). MS Thesis. Michigan State University, East Lansing, MI, pp. 79.
- Dreistadt S.H. and Dahlsten D.L. 1989. Gypsy moth eradication in Pacific Coast states: history and evaluation. *Bull. Entomol. Soc. Am.* 35: 13–19.
- Dubois N.R. and Dean D.H. 1995. Synergism between CryIA insecticidal crystal proteins and spores *Bacillus thuringiensis*, other bacterial spores, and vegetative cells against *Lymantria dispar* (Lepidoptera: Lymantriidae) larvae. *Environ. Entomol.* 24: 1741–1747.
- Duke L., Steinkraus D.C., English J.E. and Smith K.G. 2002. Infectivity of resting spores of *Massospora cicadina* (Entomophthorales: Entomophthoraceae), an entomopathogenic fungus of periodical cicadas (*Magicada* spp.) (Homoptera: Cicadidae). *J. Invert. Path.* 80: 1–6.
- Dwyer G., Dushoff J. and Yee S.H. 2004. The combined effects of pathogens and predators on insect outbreaks. *Nature* 430: 341–345.
- Elkinton J.S. and Liebold A.M. 1990. Population dynamics of gypsy moth in North America. *Annu. Rev. Entomol.* 35: 571–596.
- Elkinton J.S., Gould J.R., Ferguson C.S., Liebold A.M. and Wallner W.E. 1990. Experimental manipulation of gypsy moth density to assess impact of natural enemies. In: Watt A., Hunter M., Leather S. and Kidd N. (eds), *Population Dynamics of Forest Insects*. Hampshire, UK, Intercept, pp. 275–287.
- Elkinton J.S., Hajec A.E., Boettner G.H. and Simons E.E. 1991. Distribution and apparent spread of *Entomophaga maimaiga* (Zygomycetes: Entomophthorales) in gypsy moth (Lepidoptera: Lymantriidae) populations in North America. *Environ. Entomol.* 20: 1601–1605.
- English J.E. 2001. Sexual selection in the lek-mating system of periodical cicadas *Magicada cassini* and *M. decim*: selection

- for male body size, dominant courtship call and pathogen avoidance. Doctoral dissertation, Univ. Arkansas, Fayetteville.
- Erickson E.H., Atmowidijoyo A.H. and Hines L. 1998. Can we produce varroa-tolerant honey bees in the United States? *Am. Bee J.* 138: 828–832.
- Faeth S.H. 1987. Community structure and folivorous insect outbreaks: the roles of vertical and horizontal interactions. In: Barbosa P. and Schultz J.C. (eds), *Insect Outbreaks*. Academic Press, NY, pp. 135–171.
- Faust R.M. and Bulla L.A. 1982. Bacteria and their toxins as insecticides. In: Kurstak E. (ed.), *Microbial and Viral Pesticides*. Marcel Dekker, NY, pp. 75–208.
- Felton G.W. and Dahlman D.L. 1984. Allelochemical induced stress: effects of L-canavanine on the pathogenicity of *Bacillus thuringiensis* in *Manduca sexta*. *J. Invertebr. Pathol.* 44: 187–191.
- Ferber D. 1999. GM crops in the cross hairs. *Science* 286: 1662–1666.
- Ferguson C.S., Elkinton J.S., Gould J.R. and Wallner W.E. 1994. Population regulation of the gypsy moth (Lepidoptera: Lymantriidae) by parasitoids: does spatial density dependence lead to temporal density dependence? *Environ. Entomol.* 23: 1155–1164.
- Ferre J. and van Rie J. 2002. Biochemistry and genetics of resistance to *Bacillus thuringiensis*. *Annu. Rev. Entomol.* 47: 501–533.
- Feitelson J.S., Payne J. and Kim L. 1992. *Bacillus thuringiensis*: Insects and beyond. *Bio/Technology* 10: 271–276.
- Flexner J.L., Lighthart B. and Croft B.A. 1986. The effects of microbial pesticides on non-target beneficial arthropods. *Agric. Ecosyst. Environ.* 16: 203–254.
- Fritz R.S. and Simms E.L. 1992. *Plant Resistance to Herbivores and Pathogens*. Univ. Chicago Press, IL.
- Gage S. 1996. Ten years of data reveal gypsy moth patterns. *Gather/Scatter Natl. Lab. Comput. Sci. Eng.* 12: 20–21.
- Gahan L.J., Gould F. and Heckel D.C. 2001. Identification of a gene associated with *Bt* resistance in *Heliothis virescens*. *Science* 293: 857–860.
- Gage S. and Pijanowski B. 1993. Application of remote sensing and analysis of digital landscape maps to assess ecological risk from pest populations. In: 25th Intern. Symp. on remote sensing and global environmental change tools for sustainable development. Graz, Austria, ERIM 2: 390–402.
- Gage S., Wirth T.M. and Simmons G.A. 1990. Predicting regional gypsy moth (Lymantriidae) population trends in an expanding population using pheromone trap catch and spatial analysis. *Environ. Entomol.* 19: 370–377.
- Glare T.R. and O'Callaghan M. 2000. *Bacillus thuringiensis*: Biology, Ecology, and Safety. Wiley, NY.
- Gould F. 1991. Arthropod behavior and the efficacy of plant protectants. *Annu. Rev. Entomol.* 36: 305–330.
- Gould F. 1998. Sustainability of transgenic insecticidal cultivars: Integrating pest genetics and ecology. *Annu. Rev. Entomol.* 43: 701–726.
- Gould J.R., Elkinton J.S. and Wallner W.E. 1990. Density-dependent suppression of experimentally created gypsy moth, *Lymantria dispar* (Lepidoptera: Lymantriidae) populations by natural enemies. *J. Anim. Ecol.* 59: 213–223.
- Gould F., Martinez-Ramirez A., Ferre J., Silva F.J. and Moar W.J. 1992. Broad spectrum resistance to *Bacillus thuringiensis* toxins in *Heliothis virescens*. *PNAS (USA)* 89: 7986–7990.
- Governor H.L. 1998. Light, nutrient availability, and defoliation effects on resource acquisition, resource allocation, and herbivore resistance of paper birch and sugar maple. M.S. Thesis. Michigan State University, Dept. Entomology, East Lansing, MI 169 pp.
- Groot A.T. and Dicke M. 2002. Insect-resistant transgenic plants in a multi-trophic context. *Plant J.* 31: 387–406.
- Haas L. and Scriber J.M. 1998. Phylloplane sterilization with bleach does not reduce *Btk* toxicity for *Papilio glaucus* larvae. *Great Lakes Entomol.* 31: 49–57.
- Hails R.S. 2000. Genetically modified plants – the debate continues. *Trends Ecol. Evol.* 15: 14–18.
- Hafner M.S., Demastes J.W., Hafner D.J., Spradling T.A., Sudman P.D. and Nadler S.A. 1998. Age and movement of a hybrid zone: implications for dispersal distance in pocket gophers and their chewing lice. *Evolution* 52: 278–282.
- Hajek A.E., Butler L., Walsh S., Silver J.C., Hain F.P., Hastings F.L., Odell T.M. and Smitley D.R. 1996. Host range of the gypsy moth (Lepidoptera: Lymantriidae) pathogen *Entomophaga maimaiga* (Zygomycetes: Entomophthorales) in field versus laboratory. *Environ. Entomol.* 25: 709–721.
- Hajek A.E., Humber R.A. and Elkinton J.S. 1995. Mysterious origin of *Entomophaga maimaiga* in North America. *Am. Entomol.* 41: 31–42.
- Hancock J.A. 2003. A framework for assessing the risk of transgenic crops. *BioScience* 53: 512–519.
- Hanson-Jesse L.C. and Obrycki J.J. 2000. Field deposition of *Bt* transgenic corn pollen: lethal effects on the monarch butterfly. *Oecologia* 125: 241–248.
- Hawkins B.A. and Marino P.C. 1997. The colonization of native phytophagous insects in North America by exotic parasitoids. *Oecologia* 112: 566–571.
- Heaney L.R. and Timm R.M. 1985. Morphology, genetics, and ecology of pocket gophers (genus *Geomys*) in a narrow hybrid zone. *Biol. J. Linn. Soc.* 25: 301–317.
- Heil M., Greiner S., Meimberg H., Kruger R., Noyer J-L., Heubl G., Linsenmair K.E. and Boland W. 2004. Evolutionary change from induced to constitutive expression of an indirect plant resistance. *Nature* 430: 205–208.
- Heimpel A.M. 1967. A critical review of *Bacillus thuringiensis* var. *thuringiensis* Berliner and other crystalliferous bacteria. *Annu. Rev. Entomol.* 12: 287–322.
- Helgason E., Andreas O., Stadt O.K., Caugant D., Johannsen H., Fouet A., Mock M., Hegna I. and Kolsto A. 2000. *Bacillus anthracis*, *Bacillus cereus*, and *Bacillus thuringiensis*, one species on the basis of evidence. *Appl. Environ. Microbiol.* 66: 2627–2630.
- Hellmich R.L., Siegfried B.D., Sears M.K., Stanley-Horn D.E., Matilla H.R., Foster J.E., Clark T.L. and Jones G.D. 2001. Monarch larvae sensitivity to *Bacillus thuringiensis*-purified proteins and pollen. *PNAS (USA)* 98: 11925–11930.
- Hermes D.A. and Mattson W.J. 1992. The dilemma of plants: to grow or defend. *Quart. Rev. Biol.* 67: 283–335.
- Hilbeck A., Moar W.J., Pusztai-Carey M., Filippini A. and Bigler F. 1998. Toxicity of *Bacillus thuringiensis* Cry1Ab toxin to the predator *Chrysopa carnea* (Neuroptera: Chrysopidae). *Environ. Entomol.* 27: 1255–1263.

- Hofte H. and Whitely H.R. 1989. Insecticidal crystal proteins of *Bacillus thuringiensis* Microbiol. Review 53: 242–255.
- Huang Y., Huang R. and Li K. 1990. A field study of the persisting effect of *Bacillus thuringiensis* in Citrus groves. Chin. J. Biol. Control 6: 131–133.
- Hwang S.Y. and Lindroth R.L. 1995. Aspen leaf quality affects gypsy moth (Lepidoptera: Lymantriidae) susceptibility to *Bacillus thuringiensis*. J. Econ. Entomol. 88: 278–282.
- Hunter M.D. 1992. Interactions within herbivore communities mediated by the hostplant: the keystone herbivore concept. In: Hunter M.D., Ohgushi T. and Price P.W. (eds), Effects of Resource Distribution on Animal–Plant Interactions. Academic Press, NY, pp. 287–325.
- Hunter M.D. and Schultz J.C. 1995. Fertilization mitigates chemical induction and herbivore responses within damaged oak trees. Ecology 76: 1226–1232.
- Ignoffo C.M. 1992. Environmental factors affecting persistence of entomopathogens. Florida Entomol. 75: 516–525.
- Ignoffo C.M., Hostetter D.L. and Pinnell R.E. 1974. Stability of *Bacillus thuringiensis* and *Baculovirus heliothis* on soybean foliage. Environ. Entomol. 3: 117–119.
- James R.R., Miller J.C. and Lighthart B. 1993. *Bacillus thuringiensis* var. *kurstaki* affects a beneficial insect, the cinnabar moth (Lepidoptera: Arctiidae). J. Econ. Entomol. 86: 334–339.
- Johnson K.S., Scriber J.M., Nitao J.K. and Smitley D.R. 1995. Toxicity of *Bacillus thuringiensis* var. *kurstaki* to three non-target Lepidoptera in field studies. Environ. Entomol. 24: 288–297.
- Johnson M.T. 1997. Interaction of resistant plants and wasp parasitoids of tobacco budworm (Lepidoptera: Noctuidae). Environ. Entomol. 26: 207–214.
- Johnson M.T. and Gould F. 1992. Interaction of genetically-engineered host plant resistance and natural enemies of *Heliothis virescens* (Lepidoptera: Noctuidae) in tobacco. Environ. Entomol. 21: 586–597.
- Kaitaniemi P., Neuvonen S. and Nyssönen T. 1999. Effects of accumulative defoliations on growth reproduction, and insect resistance in maintaining birch. Ecology 80: 524–532.
- Kanda K., Takada Y., Kawasaki F., Kato F. and Murata M. 2000. Mating in *Bacillus thuringiensis* can induce plasmid integrative prophage J7W-1. Acta Virology 44: 189–192.
- Karban R. and Baldwin I.T. 1997. Induced responses to herbivory. University of Chicago Press, Chicago, IL.
- Karban R. and Myers J.H. 1989. Induced plant responses to herbivory. Ann. Rev. Ecol. Syst. 20: 331–348.
- Karban R., Agrawal A.A., Thalen J.S. and Adler L.S. 1999. Induced plant response and information content about risk of herbivory. Trends Ecol. Evol. 14: 443–447.
- Kavaliers M. and Colwell D.D. 1995. Discrimination by female mice between the odours of parasitized and non-parasitized males. Proc. Roy. Soc. London 261: 31–35.
- Kellogg S.K., Fink L.S. and Brower L.P. 2003. Parasitism of native luna moths, *Actias luna* (L.) (Lepidoptera: Saturniidae) by the introduced *Compsilura concinnata* (Meigen) (Diptera: Tachinidae) in central Virginia, and their hyperparasitism by Trigonid wasps (Hymenoptera: Trigonidae). Environ. Entomol. 32: 1019–1027.
- Koricheva J., Larsson S. and Haukioja E. 1998. Insect performance on experimentally stressed woody plants: a meta-analysis. Ann. Rev. Entomol. 43: 195–216.
- Lindroth R.L., Scriber J.M. and Hsia M.T.S. 1988. Chemical ecology of the tiger swallowtail: mediation of host use by phenolic glycosides. Ecology 69: 814–822.
- Louda S.M., Kendall D., Conner J. and Simberloff D. 1977. Ecological effects of an insect introduced for the biological control of weeds. Science 277: 181–194.
- Lovett G.M., Christenson L.M., Groffman P.M., Jones C.G., Hart J.E. and Mitchell M.J. 2002. Insect defoliation and nitrogen cycling in forests. BioScience 52: 335–341.
- Kota M., Daniell H., Varma S., Garczynski S.F., Gould F. and Moar W.J. 1999. Overexpression of the *Bacillus thuringiensis* (Bt) Cry2Aa2 protein in chloroplasts confers resistance against susceptible and Bt-resistant insects. PNAS (USA) 96: 1840–1845.
- Krieg A. and Langenbruch G.A. 1981. Susceptibility of arthropod species to *Bacillus thuringiensis*. In: Burgess H.D. (ed.), Microbial Control of Pests and Plant Diseases. Academic Press, NY, pp. 837–896.
- Krischik V.A., Barbosa P. and Reichelderfer C. 1988. Three tropic level interactions: allelochemicals, *Manduca sexta*, and *Bacillus thuringiensis* var. *kurstaki*. Environ. Entomol. 17: 476–482.
- Kukan B. and Myers J.H. 1997. Prevalence and persistence of nuclear polyhedrosis virus in fluctuating populations of forest tent caterpillars (Lepidoptera: Lasiocampidae) in the area of Prince George, British Columbia. Environ. Entomol. 26: 882–887.
- Laird M., Lacey L.A. and Davidson E.W. 1990. Safety of Microbial Insecticides. CRC Press, Boca Raton, FL.
- Lambert B. and Peferoen M. 1992. Insecticidal promise of *Bacillus thuringiensis*. BioScience 42: 112–122.
- Leong K.L., Cano R.J. and Kubinski A.M. 1980. Factors affecting *Bacillus thuringiensis* total field persistence. Environ. Entomol. 9: 593–599.
- Leong K.L.H., Yoshimura M.A. and Kaya H.K. 1992. Low susceptibility of overwintering monarch butterflies to *Bacillus thuringiensis* Berliner. Pan-Pacific Entomol. 68: 66–88.
- Liebold A.M., Gottschalk K.W., Luzader E.R., Mason D.A., Bush R. and Twardas D.B. 1997. Gypsy moth in the United States: An Atlas. USDA Forest Service, Radnor, PA.
- Lindroth R.L. 1992. Biochemical ecology of aspen-Lepidoptera interactions. J. Kansas Entomol. Soc. 64: 372–380.
- Lindroth R.L. and Hwang S. 1996. Diversity, redundancy, and multiplicity in chemical defense systems of aspen. In: Romeo J.T., Sanders J.A. and Barbosa P. (eds), Phytochemical Diversity Redundancy Ecological Interactions. Plenum, NY, pp. 25–56.
- Losey J.E., Raynor L.S. and Carter M.E. 2002. Transgenic pollen harms monarch larvae. Nature 388: 214.
- Louda S.M., Pemberton R.W., Johnson M.T. and Follett P.A. 2003. Nontarget effects – the Achilles' heel of biological control? Retrospective analyses to reduce risk associated with biological control introductions Annu. Rev. Entomol. 48: 365–393.
- Martin P.A.W. 1994. An iconoclastic view of *Bacillus thuringiensis* ecology. Am. Entomol. 40: 85–90.
- Martin P.A.W. and Trivers R.S. 1989. Worldwide abundance and distribution of *Bacillus thuringiensis* isolates. Appl. Environ. Microbiol. 55: 2437–2442.

- Matsuki M. 1996. Regulation of plant phenolic synthesis: from biochemistry to ecology and evolution. *Aust. J. Bot.* 44: 613–634.
- Maxwell F. and Jennings P. 1980. *Breeding Plants Resistant to Insects*. Wiley, NY.
- Meade T. and Hare J.D. 1993. Effects of differential host plant consumption by *Spodoptera exigua* (Lepidoptera: Noctuidae) on *Bacillus thuringiensis* efficacy. *Environ. Entomol.* 22: 432–437.
- Meade T. and Hare J.D. 1994. Effects of genetic and environmental host plant variation on the susceptibility of two noctuids to *Bacillus thuringiensis*. *Entomol. Expt. Appl.* 70: 165–178.
- Melin B.E. and Cozzi E.M. 1990. Safety to non-target invertebrates of lepidopteran strains of *Bacillus thuringiensis* and their B-exotoxins. In: Laird M., Lacey L. and Davidson E. (eds), *Safety of Microbial Insecticides*, CRC Press, Boca Raton, FL, pp. 150–167.
- Mercador R. and Scriber J.M. 2004. Phenotypic plasticity in host selection in adult tiger swallowtail butterflies, *Papilio glaucus* (Lepidoptera: Papilionidae). In: Ananthakrishnan T.N. (ed.), *"Insects and Phenotypic Plasticity"*. Oxford & IHB Publ.
- Miller J.C. 1990. Field assessment of the effects of a microbial pest control agent on non-target Lepidoptera. *Am. Entomol.* 36: 135–139.
- Miller J.C. 1992. Effects of a microbial insecticide, *Bacillus thuringiensis kurstaki*, on non-target Lepidoptera in a spruce budworm-infested forest. *J. Res. Lepidopt.* 29: 267–276.
- Miller J.C. and West K.J. 1987. Efficacy of *Bacillus thuringiensis* and diflubenzuron on Douglas fir and oak for gypsy moth in Oregon. *J. Arboric.* 13: 240–242.
- Miyasono M., Yamamoto M., Inagaki S., Ohba S., Ishiguru T., Hayashi Y. and Takeda R. 1994. Effects of several microorganisms on the insecticidal activity of delta endotoxin of *Bacillus thuringiensis* serovar *kurstaki* HD-1. *Jpn. J. Appl. Entomol. Zool.* 38: 101–108.
- Moldenke A.F., Berry R.E., Müller J.C., Wernz J.G. and Li X.H. 1994. Toxicity of *Bacillus thuringiensis* subsp. *kurstaki* to gypsy moth, *Lymantria dispar*, fed alder or Douglas fir. *J. Insect Pathol.* 64: 143–145.
- Moullia C., Le Brun N., Loubes C., Marin R. and Renaud E. 1995. Hybrid vigor against parasites in interspecific crosses between two mice species. *Heredity* 74: 48–52.
- Moullia C., Le Brun N. and Renaud F. 1996. Mouse-parasite interactions: from gene to population. *Advances in Parasitology* 38: 119–167.
- Moullia C. 1999. Parasitism of plant and animal hybrids: Are facts and fates the same? *Ecology* 80: 392–406.
- Muck O., Hassan S., Huger A.M. and Krieg A. 1981. The effect of *Bacillus thuringiensis* Berliner on the parasitic hymenopterans *Apanteles glomeratus* L. (Braconidae) and *Pimpla thurionella* (L.) (Ichneumonidae). *Z. Angew. Entomol.* 92: 303–314.
- Nap, J.-P., Metz P.L.J., Escaler M. and Conner A.J. 2003. The release of genetically modified crops into the environment. Part I. Overview of current status and regulations. *Plant J.* 33: 1–18.
- Nasr M.E. 1997. Tracheal mite resistant and hygienic honey bee stocks in Ontario. *Can. Beekeep.* 20: 63–64.
- Navon A., Klein M. and Braun S. 1990. *Bacillus thuringiensis* potency bioassays against *Heliothis armigera*, *Earias insulana*, and *Spodoptera littoralis* larvae based on standardized diets. *J. Invertebr. Pathol.* 55: 387–393.
- Neuvonen S., Haukioja E. and Molarius A. 1987. Delayed inducible resistance against a leaf-chewing insect in four deciduous tree species. *Oecologia* 74: 363–369.
- New T.R. 1997. *Butterfly Conservation*. 2nd ed. Oxford Univ. Press, Melbourne.
- Nykanen H. and Koricheva J. 2004. Damage-induced changes in woody plants and their effects on insect herbivore performances: a meta-analysis. *Oikos* 104: 247–268.
- Oberhauser K.S., Prysby M.D., Mattila H.R., Stanley-Horn D.E., Sears M.K., Dively G.P., Olson E., Pleasants J.M., Lam, W.-K.F. and Hellmich R.L. 2001. Temporal and spatial overlap between monarch larvae and corn pollen. *PNAS (USA)* 98: 11913–11918.
- Papp-Herms C.P. 1996. The endangered Karner Blue butterfly (Lepidoptera: Lycaenidae) in Michigan: habitat suitability, potential impacts of gypsy moth (Lepidoptera: Lymantriidae) suppression, and laboratory rearing. M.Sc. thesis. Dept. Entomology, Michigan State Univ., East Lansing, MI.
- Parry D. 2000. Induced responses of poplars to defoliation and their effects on leaf-feeding Lepidoptera. Dissertation. Michigan State University, East Lansing, 232 pp.
- Parry D., Herms D.A. and Mattson W. 2003. Responses of an insect folivore and its parasitoids to multiyear experimental defoliation of aspen. *Ecology* 84: 1768–1783.
- Parry D., Spence J.R. and Volney W.J.A. 1997. Responses of natural enemies to experimentally increased populations of the forest tent caterpillar, *Malacosoma disstria*. *Ecol. Entomol.* 22: 97–108.
- Peacock J.W., Wagner D.L. and Schweitzer D.E. 1993. Impacts of *Bt* on non-target Lepidoptera. *News. Mich. Entom. Soc.* 38: 1–5.
- Peacock J.W., Schweitzer D.F., Dale F., Carter J.L. and Dubois N.R. 1998. Laboratory assessment of the effects of *Bacillus thuringiensis* on native Lepidoptera. *Environ. Entomol.* 27: 450–457.
- Peters D.J. and Constabel C.P. 2002. Molecular analysis of herbivore-induced condensed tannin synthesis: cloning and expression of dihydroflavonol reductase from trembling aspen (*Populus tremuloides*). *Plant J.* 32: 701–712.
- Petrus S.F. and Cassida L.E. 1985. Survival of *Bacillus thuringiensis* spores in soil. *Appl. Environ. Microbiol.* 50: 1496–1501.
- Pimentel D.S. and Raven P.H. 2000. Bt corn impacts on non-target Lepidoptera: assessment of effects in nature. *PNAS (USA)* 97: 8198–8199.
- Pinnock D.E., Brand R.J. and Milstead J.E. 1971. The field persistence of *Bacillus thuringiensis* spores. *J. Invert. Pathol.* 18: 405–411.
- Pleasants J.M., Hellmich R.L., Dively G.P., Sears M.K., Stanley-Horn D., Mattila D.E., Foster J.E., Clark T.L. and Jones G.D. 2001. Corn pollen deposition on milkweeds in and near cornfields. *PNAS (USA)* 98: 11919–11924.
- Radcliffe S.L. and Yendol W.G. 1993. Lethal dose and associated effects of *Bacillus thuringiensis* in sprayed droplets against gypsy moth. *J. Environ. Sci. Health* 28: 91–104.
- Rastall K., Kondo V., Strazanac J.S. and Butler L. 2003. Lethal effects of biological insecticide applications on nontarget

- lepidopterans in two Appalachian forests. *Environ. Entomol.* 32: 1364–1369.
- Reardon R., Dubois N. and McLane W. 1994. *Bacillus thuringiensis* for managing gypsy moth: a review. USDA Forest Service FHM-NC-01-94, Washington, DC.
- Reardon R.C. and Haissig K. 1984. Efficacy and field persistence of *Bacillus thuringiensis* after ground application to Balsam fir and White spruce in Wisconsin. *Canad. Entomol.* 116: 1523–1526.
- Redman A. and Scriber J.M. 2000. Competition between gypsy moth, *Lymantria dispar*, and the northern tiger swallowtail, *Papilio canadensis*: interactions mediated by host plant chemistry, pathogens, and parasitoids. *Oecologia* 125(2): 218–228.
- Reichelderfer C.F. 1991. Interactions among allelochemicals, some Lepidoptera, and *Bacillus thuringiensis* Berliner. In: Barbosa P., Krischik V.A. and Jones C.G. (eds), *Microbial Mediation of Plant–Herbivore Interactions*. Wiley, NY, pp. 507–524.
- Renwick J.A.A. and Chew F. 1994. Oviposition behavior in Lepidoptera. *Annu. Rev. Entomol.* 39: 377–400.
- Robison D.J. and Raffa K.F. 1994. Characterization of hybrid aspen clones for resistance to the forest tent caterpillar. *Forest Sci.* 40: 686–714.
- Rossiter M.C. 1994. Maternal effects hypothesis of herbivore outbreak. *Bioscience* 44: 752–763.
- Sammataro D., Gerson U. and Needham G. 2000. Parasitic mites of honey bees: life history, implications, and impact. *Annu. Rev. Entomol.* 45: 519–548.
- Sample B.E., Butler L., Zivkovich C., Whitmore R.C. and Reardon R. 1996. Effects of *Bacillus thuringiensis* var. *kurstaki* and defoliation by gypsy moth on native arthropods in West Virginia. *Canad. Entomol.* 128: 573–592.
- Saxena D. and Stotsky G. 2001. *Bacillus thuringiensis* (Bt) toxin released from root exudates and biomass of Bt corn has no apparent effect on earthworms, nematodes, protozoa, bacteria, and fungi in the soil. *Soil Biol. Biochem.* 33: 1225–1230.
- Schweitzer D.E. 1988. Status of the Saturniidae in the north-eastern USA: a quick review. *New Lepidop. Soc.* 1: 4–5.
- Scoble M.J. 1992. *The Lepidoptera: Form, Function, and Diversity*. Oxford Univ. Press, UK.
- Scriber J.M. 1992. Latitudinal trends in oviposition preferences: ecological and genetic influences. In: Menken S.B.J., Visser J.H. and Harrewijn P. (eds), *Proc. 8th Intern. Symp. Insect-Plant Relationships*. Kluwer Academic Publishers, B.V. Dordrecht, Netherlands, pp. 212–214.
- Scriber J.M. 1993. Absence of behavioral induction in oviposition preference of *Papilio glaucus* (Lepidoptera: Papilionidae). *Great Lakes Entomol.* 26: 81–95.
- Scriber J.M. 1996. Tiger tales: natural history of native North American swallowtails. *Am. Entomol.* 42: 19–32.
- Scriber J.M. 1998. Non-target Lepidoptera impacted by Btk pesticide sprays, induced phytochemical defenses, and generalized parasitoids and predators of *Lymantria dispar*. In: Zaluki M.P., Drew R.A.I. and White G.G. (eds), *Pest Management – Future Challenges*, Vol. 2, Univ. Queensland, Brisbane, Australia, pp. 236–246.
- Scriber J.M. 2001. *Bt* or not *Bt*? Is that the question (Invited Commentary on 6 Bt corn pollen articles). *Natl. Acad. Sci. (USA)* 98: 12328–12330.
- Scriber J.M. 2002a. The evolution of insect-plant relationships; chemical constraints, coadaptation and concordance of insect/plant traits. *Entomologia Exp. Appl.* 104: 217–235.
- Scriber J.M. 2002b. Latitudinal and local geographic mosaics in host plant preferences as shaped by thermal units and voltinism. *Eur. J. Entomol.* 99: 225–239.
- Scriber J.M., Lederhouse R.C. and Dowell R. 1995. Hybridization studies with North American swallowtails. In: Scriber J.M., Tsubaki Y. and Lederhouse R.C. (eds), *The Swallowtail Butterflies: Their Ecology and Evolutionary Biology*. Scientific Publishers Inc., Gainesville, FL, pp. 269–282.
- Scriber J.M. and Gage S. 1995. Pollution and global climate change: Plant ecotones, butterfly hybrid zones, and biodiversity. In: Scriber J.M., Tsubaki Y. and Lederhouse R.C. (eds), *Swallowtail Butterflies: Their Ecology and Evolutionary Biology*. Scientific Publishers, Inc., Gainesville, FL, pp. 319–344.
- Scriber J.M. and Haas L. 1996. Long term effects of *B.t.k.* on non-target Lepidoptera species and potential causal mechanisms. *Proceedings 1995 National Gypsy Moth Review*, MI, Traverse City, pp. 132–137.
- Scriber J.M., Lindroth R.L. and Nitao J. 1989. Differential toxicity of a phenolic glycoside from quaking aspen leaves by *Papilio glaucus* subspecies, their hybrids, and backcrosses. *Oecologia* 81: 186–191.
- Scriber J.M., Weir K., Parry D. and Deering J. 1999. Using hybrid and backcross larvae of *Papilio canadensis* and *P. glaucus* to detect induced chemical resistance in hybrid poplars experimentally defoliated by gypsy moths. *Entomologia experimentalis et applicata* 91: 233–236.
- Scriber J.M., Stump A. and Deering M. 2003. Hybrid zone ecology and tiger swallowtail trait clines in North America. In: Boggs C., Watt W. and Ehrlich P. (eds), *Ecology and Evolution Taking Flight: Butterflies as Model Study Systems*. University of Chicago Press, Chicago, IL, pp. 367–391.
- Scriber J.M., Tsubaki Y. and Lederhouse R.C. 1995. *Swallowtail Butterflies: Their Ecology and Evolutionary Biology*. Scientific Publ., Gainesville, FL.
- Sears M.K., Hellmich R.L., Stanley-Horn D.E., Oberhauser K.S., Pleasants J.M., Mattila H.R., Siegfried B.D. and Dively G.P. 2001. Impact of Bt corn pollen on monarch butterfly populations: a risk assessment. *PNAS (USA)* 98: 11937–11942.
- Sharov A.A., Liebold A.M. and Roberts E.A. 1998. Optimizing the use of barrier zones to slow the spread of gypsy moth (Lepidoptera: Lymantriidae) in North America. *J. Econ. Entomol.* 91: 165–174.
- Sharov A.A., Pijanowski B.C., Liebold A.M. and Gage S.H. 1999. What affects the rate of gypsy moth (Lepidoptera: Lymantriidae) spread: winter temperature or forest susceptibility. *Agr. Forest Entomol.* 1: 37–45.
- Sheehan W. 1994. Parasitoid community studies: effects of host abundance, phylogeny, and ecology. In: Hawkins B.A. and Sheehan W. (eds), *Parasitoid Community Ecology*. Oxford Univ. Press, NY, pp. 90–107.
- Shelton A.M., Zhao J.-Z., and Roush R.T. 2002. Economic, ecological, food safety, and social consequences of the deployment of Bt transgenic plants. *Annu. Rev. Entomol.* 47: 845–881.
- Slansky F. and Scriber J.M. 1985. Food consumption and utilization. In: Kerkut G.A. and Gilbert L.I. (eds), *Compre-*

- hensive Insect Physiology, Biochemistry, and Pharmacology, Vol. 4, Pergamon Press, Oxford, pp. 87–163.
- Smith R.A. and Couche G.A. 1991. The phylloplane as a source of *Bacillus thuringiensis* variants. Appl. Environ. Microbiol. 57: 311–315.
- Smitley D., Andresen J., Priest R., Mech R. and McCullough D. 1998. Winter mortality of gypsy moth (Lepidoptera: Lymantriidae) eggs in Michigan. Environ. Entomol. 27: 700–708.
- Stamp N.E. 1990. Parasitism of New England buck moth caterpillars (*Hemilieuca lucina*: Saturniidae) tachinid flies. J. Lepid. Soc. 44: 199.
- Stamp N.E. and Casey T.M. 1993. Caterpillars: Ecological and Evolutionary Constraints on Foraging. Chapman and Hall, NY.
- Stewart A.J.A. 1996. Interspecific competition reinstated as an important force structuring insect herbivore communities. Trends Ecol. Evol. 11: 233–234.
- Stiling P. and Simberloff D. 2000. The frequency and strength of nontarget effects of invertebrate biological control agents of plant pests and weeds. In: Follett P.A. and Duan J.J. (eds), Nontarget Effects of Biological Control. Kluwer Academic, Boston, pp. 31–43.
- Sundarum K.M.S. and Sundarum A. 1992. An insect bioassay method to determine persistence of *Bacillus thuringiensis* var. *kurstaki* (B.t.k.) protein in oak foliage, following application of a commercial formulation under field and laboratory conditions. J. Environ. Health Part B. Pest. Food Contain. Agric Wastes 27: 73–112.
- Swadener C. 1994. *Bacillus thuringiensis* (B.t.) J. Pesticide Reform 14: 13–20.
- Tallamy D.W. and Raupp M.J. 1991. Phytochemical Induction by Herbivores. Wiley, NY.
- Tapp H. and Stotzky G. 1998. Persistence of the insecticidal toxin from *Bacillus thuringiensis* subsp. *kurstaki* in soil. Soil Biochem. 30: 471–476.
- Tuskes P.M., Tuttle J.P. and Collins M.M. 1996. The Wild Silk Moths of North America. Cornell Univ. Press, Ithaca, NY.
- Van Emden H.F. 1999. Transgenic host plant resistance to insects—some reservations. Ann. Entomol. Soc. Am. 92: 788–797.
- Van Frankenhuyzen K., Gringorten J.L., Milne R.E., Gauthier D., Pusztai M., Brousseau R. and Masson L. 1991. Specificity of activated CryIA proteins from *Bacillus thuringiensis* subsp. *kurstaki* HD-1 for defoliating forest Lepidoptera. Appl. Environ. Microbiol. 57: 1650–1655.
- Venable B.A.B. 1990. Preliminary assessment of the susceptibility of non-target Lepidopteran species to *Bacillus thuringiensis* (B.t.) and dimilin for gypsy moth suppression. Rept. For USDA Dept. Interior Nat. Parks Serv. National Capital Region Urban Ecology Program, Wash. DC.
- Wagner D.L., Peacock J.W., Carter J.L. and Talley S.E. 1996. Field assessment of *Bacillus thuringiensis* on non-target Lepidoptera. Environ. Entomol. 25: 1444–1454.
- Wait D.A., Jones C.L. and Coleman J.S. 1998. Effects of nitrogen fertilization on leaf chemistry and beetle feeding are mediated by leaf development. Oikos 82: 502–514.
- Wang G., Castiglione S., Chen Y., Li L., Han Y., Gabriel D.W., Han Y., Mang K. and Scakla F. 1996. Poplar (*Populus nigra*) plants transformed with a *Bacillus thuringiensis* toxin gene: insecticidal activity and genomic analysis. Transgenic Res. 5: 289–301.
- Watanabe, K., Kara W. and Sato M. 1998. Evidence for growth of strains of plant epiphytic bacterium *Erwinia herbicola* and transconjugation among bacterial strains in guts of the silkworm, *Bombyx mori*. J. Invertebr. Pathol. 72: 104–111.
- Webber R.T. and Schaffner J.V. 1926. Host relations of *Compsilura concinnata* Meigen, an important tachinid parasite of the gypsy moth and brown-tail moth. USDA Bulletin 1363.
- Weseloh R.M. and Andreadis T.G. 1997. Persistence of resting spores of *Entomophaga maimaiga*, a fungal pathogen of the gypsy moth, *Lymantria dispar*. J. Invertebr. Pathol. 69: 195–196.
- West A.W. and Burgess H.D. 1985. Persistence of *Bacillus thuringiensis* and *Bacillus cereus* in soil supplemented with grass or manure. Plant Soil 83: 389–398.
- Whaley W.H., Arnold J. and Schaalje B. 1998. Canyon drift and dispersion of *Bacillus thuringiensis* and its effects on select non-target lepidopterans in Utah. Environ. Entomol. 27: 539–548.
- Whitham T.G., Morrow P.A. and Potts B.M. 1994. Plant hybrid zones as centers of biodiversity: the herbivore community of two endemic Tasmanian eucalypts. Oecologia 97: 481–490.
- Whitham T.G., Martinsen G.D., Floate K.D., Dungey H.S., Potts B.M. and Keim P. 1999. Plant hybrid zones affect biodiversity: tools for a genetically-based understanding of community structure. Ecology 80: 416–428.
- Williams D.W., Fuester R.W., Metterhouse W.W., Balaam R.J., Bullock R.H., Chianese R.J. and Reardon R.C. 1992. Incidence and ecological relationships of parasitism in larval populations of *Lymantria dispar* (Lepidoptera: Lymantriidae). Biol. Control 2: 35–43.
- Wolfenbarger L.L. and Phifer P.R. 2000. The ecological risks and benefits of genetically engineered plants. Science 290: 2088–2093.
- Wraight C.L., Zangerl A.R., Carroll M.J. and Berenbaum M.R. 2000. Absence of toxicity of *Bacillus thuringiensis* pollen to black swallowtails under field conditions. PNAS (USA) 97: 7700–7703.
- Zalucki M.P., Clarke A.R. and Malcolm S.B. 2002. Ecology and behavior of first instar larval Lepidoptera. Annu. Rev. Entomol. 47: 361–393.
- Zangerl A.R., McKenna D., Wraight C.L., Carroll M., Ficarello P., Warner R. and Berenbaum M.R. 2001. Effects of exposure to event 176 *Bacillus thuringiensis* corn pollen on monarch and black swallowtail caterpillars under field conditions. PNAS (USA) 98: 11908–11912.