

Elsevier L^AT_EX template[☆]

Elsevier¹

Radarweg 29, Amsterdam

Elsevier Inc^{a,b}, Global Customer Service^{b,}*

^a1600 John F Kennedy Boulevard, Philadelphia

^b360 Park Avenue South, New York

Abstract

This template helps you to create a properly formatted L^AT_EX manuscript.

Keywords: `elsarticle.cls`, L^AT_EX, Elsevier, template

2010 MSC: 00-01, 99-00

1. GLRT

Suppose $\{X_{i1}, \dots, X_{in_i}\}$ are i.i.d. distributed as $N(\mu_i, \Sigma)$ for $1 \leq i \leq K$. Let $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$ for $i = 1, \dots, k$. The k samples are independent. μ_i , $i = 1, \dots, k$ and $\Sigma > 0$ are unknown. An interesting problem in multivariate analysis is to test the hypotheses

$$H : \mu_1 = \mu_2 = \dots = \mu_k \quad v.s. \quad K : \mu_i \neq \mu_j \text{ for some } i \neq j. \quad (1)$$

Let $\mathbf{Z} = (X_1, \dots, X_k)$.

$$f(Z; \mu_1, \dots, \mu_k, \Sigma) = \prod_{i=1}^k \left[(2\pi)^{-n_i p/2} |\Sigma|^{-n_i/2} \exp\left(-\frac{1}{2} \text{tr} \Sigma^{-1} \sum_{j=1}^{n_i} (x_{ij} - \mu_i)(x_{ij} - \mu_i)^T\right) \right].$$

Assume $n = \sum_{i=1}^p n_i < p$. Let $a \in \mathbb{R}^p$ be a vector satisfying $a^T a = 1$. Then

$$f_a(a^T Z; \mu_1, \dots, \mu_k, \Sigma) = (2\pi)^{-n/2} |a^T \Sigma a|^{-n/2} \exp\left(-\frac{1}{2a^T \Sigma a} \sum_{i=1}^k \sum_{j=1}^{n_i} (a^T x_{ij} - a^T \mu_i)^2\right)$$

[☆]Fully documented templates are available in the elsarticle package on CTAN.

^{*}Corresponding author

Email address: support@elsevier.com (Global Customer Service)

URL: www.elsevier.com (Elsevier Inc)

¹Since 1880.

$$\max_{\mu_1, \dots, \mu_k, \Sigma} f_a(a^T Z, \mu_1, \dots, \mu_k, \Sigma) = (2\pi)^{-n/2} \left(\sum_{i=1}^k \sum_{j=1}^{n_i} (a^T x_{ij} - a^T \bar{\mathbf{X}}_i)^2 \right)^{-n/2} e^{-n/2} \quad (2)$$

Let $S_i = \sum_{j=1}^{n_i} (x_{ij} - \bar{\mathbf{X}}_i)(x_{ij} - \bar{\mathbf{X}}_i)^T$ and $S = \sum_{i=1}^k S_i$.

Under H , we have

$$\max_{\mu, \Sigma} f_a(a^T Z, \mu, \dots, \mu, \Sigma) = (2\pi)^{-n/2} \left(\sum_{i=1}^k \sum_{j=1}^{n_i} (a^T x_{ij} - a^T \bar{\mathbf{X}})^2 \right)^{-n/2} e^{-n/2} \quad (3)$$

The generalized likelihood ratio test statistic is defined as

$$T(Z) = \max_{a^T a=1, a^T S a=0} a^T \sum_{i=1}^k n_i (\bar{\mathbf{X}}_i - \bar{\mathbf{X}})(\bar{\mathbf{X}}_i - \bar{\mathbf{X}})^T a \quad (4)$$

Let $J = \text{diag}(n_1^{-1/2} \mathbf{1}_{n_1}, \dots, n_k^{-1/2} \mathbf{1}_{n_k})$. Then $S = Z(I_n - JJ^T)Z^T$ and

$$\sum_{i=1}^k n_i (\bar{\mathbf{X}}_i - \bar{\mathbf{X}})(\bar{\mathbf{X}}_i - \bar{\mathbf{X}})^T = Z(JJ^T - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T)Z^T. \quad (5)$$

The matrix $I_n - JJ^T$, $JJ^T - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T$ and $\frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T$ are all projection matrix and pairwise orthogonal with rank $n - k$, $k - 1$ and 1.

Let \tilde{J} be a $n \times (n - k)$ matrix satisfied $\tilde{J}\tilde{J}^T = I - JJ^T$. Then $S = Z\tilde{J}\tilde{J}^T Z^T$ and Note that

$$Z(JJ^T - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T)Z^T = ZJ(I_k - \frac{1}{n} J^T \mathbf{1}_n \mathbf{1}_n^T J)J^T Z^T.$$

5 Note that $I_k - \frac{1}{n} J^T \mathbf{1}_n \mathbf{1}_n^T J$ is a projection matrix with rank $k - 1$. Let C be a $k \times (k - 1)$ matrix satisfied $CC^T = I_k - \frac{1}{n} J^T \mathbf{1}_n \mathbf{1}_n^T J$.

In Proposition 1, letting $A = Z\tilde{J}$ and $B = ZJCC^T J^T Z^T$ yields

$$T(Z) = \lambda_{\max}((I_p - H_A)ZJCC^T J^T Z^T(I_p - H_A)) = \lambda_{\max}(C^T J^T Z^T(I_p - H_A)ZJC),$$

where $H_A = Z\tilde{J}(\tilde{J}Z^T Z\tilde{J})^{-1} \tilde{J}^T Z^T$. Note that

$$\begin{aligned} & \left(\begin{pmatrix} J^T \\ \tilde{J}^T \end{pmatrix} Z^T Z \begin{pmatrix} J & \tilde{J} \end{pmatrix} \right)^{-1} \\ &= \begin{pmatrix} J^T Z^T Z J & J^T Z^T Z \tilde{J} \\ \tilde{J}^T Z^T Z J & \tilde{J}^T Z^T Z \tilde{J} \end{pmatrix}^{-1} = \begin{pmatrix} J^T (Z^T Z)^{-1} J & J^T (Z^T Z)^{-1} \tilde{J} \\ \tilde{J}^T (Z^T Z)^{-1} J & \tilde{J}^T (Z^T Z)^{-1} \tilde{J} \end{pmatrix}. \end{aligned} \quad (6)$$

It follows that

$$\begin{aligned}
& (J^T(Z^T Z)^{-1}J)^{-1} \\
&= J^T Z^T Z J - J^T Z^T Z \tilde{J}(\tilde{J}^T Z^T Z \tilde{J})^{-1} \tilde{J}^T Z^T Z J \\
&= J^T Z^T (I_p - Z \tilde{J}(\tilde{J}^T Z^T Z \tilde{J})^{-1} \tilde{J}^T Z^T) Z J
\end{aligned} \tag{7}$$

It follows that

$$T(Z) = \lambda_{\max}\left(C^T (J^T(Z^T Z)^{-1}J)^{-1}C\right) \tag{8}$$

Proposition 1. *Suppose A is a $p \times r$ matrix with rank r and B is a $p \times p$ non-zero semi-definite matrix. Let $H_A = A(A^T A)^{-1}A^T$. Then*

$$\max_{a^T a=1, a^T A A^T a=0} a^T B a = \lambda_{\max}((I_p - H_A)B(I_p - H_A)). \tag{9}$$

Proof. Note that $a^T A A^T a = 0$ is equivalent to $A^T a = 0$ and is in turn equivalent to $H_A a = 0$. In this circumstance, $a = (I_p - H_A)a$. Then

$$\begin{aligned}
\max_{a^T a=1, a^T A A^T a=0} a^T B a &= \max_{a^T a=1, H_A a=0} a^T B a \\
&= \max_{a^T a=1, H_A a=0} a^T (I_p - H_A)B(I_p - H_A)a.
\end{aligned} \tag{10}$$

It's obvious that $(10) \leq \lambda_{\max}((I - H_A)B(I - H_A))$. On the other hand, let α_1 be one eigenvector corresponding to the largest eigenvalue of $(I - H_A)B(I - H_A)$. Note that the row of H_A are all eigenvectors of $(I - H_A)B(I - H_A)$ corresponding to eigenvalue 0. It follows that $H_A \alpha_1 = 0$. Now that α_1 satisfies the constraint of (10), (10) is maximized when $a = \alpha_1$. □

2. The Elsevier article class

Installation. If the document class `elsarticle` is not available on your computer, you can download and install the system package `texlive-publishers` (Linux) or install the L^AT_EX package `elsarticle` using the package manager of your T_EX installation, which is typically T_EX Live or MikT_EX.

Usage. Once the package is properly installed, you can use the document class *elsarticle* to create a manuscript. Please make sure that your manuscript follows the guidelines in the Guide for Authors of the relevant journal. It is not necessary to typeset your manuscript in exactly the same way as an article, unless you are submitting to a camera-ready copy (CRC) journal.

Functionality. The Elsevier article class is based on the standard article class and supports almost all of the functionality of that class. In addition, it features commands and options to format the

- document style
- baselineskip
- front matter
- keywords and MSC codes
- theorems, definitions and proofs
- lables of enumerations
- citation style and labeling.

3. Front matter

The author names and affiliations could be formatted in two ways:

- (1) Group the authors per affiliation.
- (2) Use footnotes to indicate the affiliations.

See the front matter of this document for examples. You are recommended to conform your choice to the journal you are submitting to.

4. Bibliography styles

40 There are various bibliography styles available. You can select the style of
your choice in the preamble of this document. These styles are Elsevier styles
based on standard styles like Harvard and Vancouver. Please use BibTeX to
generate your bibliography and include DOIs whenever available.

Here are two sample references: [? ?].

45 **References**