

On the Wilks phenomenon of Bayes factors

Rui Wang¹ and Xingzhong Xu^{*1,2}

¹ School of Mathematics and Statistics, Beijing Institute of Technology, Beijing
100081, China

² Beijing Key Laboratory on MCAACI, Beijing Institute of Technology, Beijing
100081, China

Friday 21st June, 2019

Abstract

Likelihood ratio test play a prominent role in parametric hypotheses testing. A key property of the likelihood ratio test statistic is the Wilks phenomenon, that is, the asymptotic null distribution is free of nuisance parameters, which allows to determine the critical value from the asymptotic distribution. However, the likelihood ratio test can not be used for some moderately complex models, for example, the models with unbounded likelihood functions. Bayes factor and its variants have been extensively studied in Bayesian hypotheses testing framework and have been shown to have good performance in a wealth of testing problems. The Bayes factor can be applied to complex problems, even if the likelihood functions are unbounded. The good performance of Bayes factor in complex models motivates us to use Bayes factors as frequentist test statistics. We investigate the asymptotic distribution of Bayes factor and two of its variants, i.e. posterior Bayes factor and fractional Bayes factor. These asymptotic results do not require the likelihood is bounded. It shows that the classical Bayes factor has Wilks phenomenon only for a restricted class of prior distributions, and some popular objective priors do not yield Bayes factors with Wilks phenomenon. On the other hand, posterior Bayes factor and fractional Bayes factor have Wilks phenomenon for general priors. Frequentist tests based on Bayes factors are constructed using the Wilks phenomenon. For regular models, the proposed tests have the same asymptotic local power as the likelihood ratio test. We apply the proposed test to two submodels of the normal mixture model. The likelihood ratio test cannot be defined for the first submodel and has undesirable local power behavior for the second submodel. In contrast, we show that the proposed test has good asymptotic power behavior for both submodels.

Key words: Bayes consistency, fractional posterior, integrated likelihood ratio, mixture model, posterior Bayes factor.

*Corresponding author

Email address: xuxz@bit.edu.cn

1 Introduction

Likelihood inference plays a dominant role in parametric statistic inference. On the one hand, the maximum likelihood estimation is asymptotically optimal in a great variety of problems. On the other hand, the fundamental lemma of Neyman and Pearson tells us that the likelihood ratio test (LRT) is the most powerful test if the null and alternative hypotheses are both simple. For testing composite hypotheses

$$H : \theta \in \Theta_0 \quad \text{vs.} \quad K : \theta \in \Theta_1, \quad (1)$$

where $\Theta_0 \cap \Theta_1 = \emptyset$, $\Theta_0 \cup \Theta_1 = \Theta$, Θ is an open subset of \mathbb{R}^p and Θ_0 is a p_0 -dimensional subset of \mathbb{R}^p , the LRT statistic is defined as

$$\Lambda_{\text{LRT}} = \frac{\max_{\theta \in \Theta} L(\theta)}{\max_{\theta \in \Theta_0} L(\theta)} = \frac{L(\hat{\theta}_{\text{MLE}})}{L(\hat{\theta}_{\text{MLE}}^{(0)})},$$

where $L(\theta)$ is the likelihood function, $\hat{\theta}_{\text{MLE}}$ and $\hat{\theta}_{\text{MLE}}^{(0)}$ are the MLE of θ in Θ and Θ_0 , respectively. In a celebrated paper, (Wilks, 1938) proved that for regular models, the asymptotic null distribution of the LRT statistic is free of nuisance parameters. With this important property, one can determine the critical value of the LRT statistic using only its asymptotic null distribution and do not need to estimate the nuisance parameters. In this paper, we say a test statistic has Wilks phenomenon if its asymptotic null distribution does not depend on the nuisance parameters. The LRT has been very successful in many specific problems. However, for some moderately complex problems, some difficulties may arise when using the LRT. The maximization of $L(\theta)$ may be difficult if the likelihood function is not concave and has multiple local maxima. Worse still, in some problems the likelihood functions are unbounded and hence the LRT is not defined; see, e.g., Le Cam (1990). Notice that the unbounded likelihood occurs not only in artificial models, but also in some widely used models, such as the mixture models with unknown component location and scale parameters (Chen, 2017).

In goodness of fit test, there are two common types of tests: extreme value type (Kolmogorov-Smirnov test, e.g.) and integral type (Cramér-von Mises test, e.g.). In classical parametric hypothesis testing, however, no attention has been paid to the integrated likelihood functions. A natural integral type test statistic for hypothesis (1) is

$$\frac{\int_{\Theta} L(\theta) d\Pi(\theta)}{\int_{\Theta_0} L(\theta) d\Pi^{(0)}(\theta)}, \quad (2)$$

where Π and $\Pi^{(0)}$ are some probability measures on Θ and Θ_0 , respectively. If Π and $\Pi^{(0)}$ are independent of data, then the statistic (2) is exactly the Bayes factor (Jeffreys, 1931) with the prior distributions Π and $\Pi^{(0)}$. The Bayes factor is the conventional tool for Bayesian hypothesis testing and has been widely used by practitioners; see Kass and Raftery (1995) for a review. However, Bayes factor is sensitive to the choice of prior distribution. In fact, the asymptotic distribution of Bayes factor depends on the prior density at the true parameter; see, e.g., Clarke and Barron

(1990). As a result, the Bayes factor cannot be treated as a frequentist test statistic. Thus, the measures Π and $\Pi^{(0)}$ considered in this paper will depend on data.

If $\Pi(\theta = \hat{\theta}_{\text{MLE}}) = 1$ and $\Pi^{(0)}(\theta = \hat{\theta}_{\text{MLE}}^{(0)}) = 1$, then the statistic (2) becomes the LRT statistic. In this case, the measure Π and $\Pi^{(0)}$ both concentrate on one point and hence are highly nonsmooth. For many models where the LRT fails, the likelihood function $L(\theta)$ still has good properties for most θ and the MLE is unfortunately trapped in a fairly small area of θ where $L(\theta)$ has bad behavior. Figure 1 exhibits this phenomenon. Intuitively, if Π and $\Pi^{(0)}$ are smooth and have small tail probability, then the defeat of the likelihood function in a small area will not introduce much effect on the integrated likelihood. Following this idea, a natural choice is to take Π and $\Pi^{(0)}$ as the posterior distribution (with certain prior distributions) of θ in Θ and Θ_0 , respectively. In this case, the statistic (2) becomes the posterior Bayes factor (PBF) proposed by Aitkin (1991). Aitkin (1991) argued that if the likelihood concentrates around the MLE, the PBF should approximately equal to $2^{(p-p_0)/2}\Lambda_{\text{LRT}}$. This implies that PBF has the similar Wilks phenomenon as the LRT.

If we replace the likelihood function $L(\theta)$ by $L(\theta)^a$ for $a > 0$, then the LRT statistic becomes

$$\frac{\max_{\theta \in \Theta} L^a(\theta)}{\max_{\theta \in \Theta_0} L^a(\theta)} = \Lambda_{\text{LRT}}^a,$$

which is equivalent to the LRT statistic. In contrast, the statistic

$$\frac{\int_{\Theta} L^a(\theta) d\Pi(\theta)}{\int_{\Theta_0} L^a(\theta) d\Pi^{(0)}(\theta)} \quad (3)$$

is not equivalent to the statistic (2). We will also consider the test statistic (3) with $0 < a < 1$. Correspondingly, the measure Π and $\Pi^{(0)}$ can also take the fractional posterior (Bhattacharya et al., 2019). Raising the likelihood to a fractional power has several advantages; see, e.g., Walker and Hjort (2001) and Bhattacharya et al. (2019). In particular, the consistency of the fractional posterior requires less conditions than the consistency of the usual posterior. A special case of the statistic (3) is the fractional Bayes factor (FBF) proposed by O’Hagan (1995). We call the statistic (3) the generalized FBF if Π and $\Pi^{(0)}$ are fractional posterior distributions.

Under certain regular conditions, we rigorously prove the Wilks phenomenon of the generalized FBF. Based on the Wilks phenomenon, an asymptotically correct frequentist test procedure can be formulated. We also give the asymptotic local power of the resulting test procedure under contiguous alternative. It is shown that the generalized FBF has a similar asymptotic local power to the LRT. However, the generalized FBF can be applied to the cases where the likelihood is unbounded and thus has a wider application scope than the LRT.

The generalized FBF can be computed by sampling θ from the fractional posterior and calculate the sample mean of the fractional likelihood. For moderately complex model, however, sampling from the fractional posterior may be difficult and hence some approximation methods may be used in practice. Variational inference is a popular method for approximating intractable posterior distribution; see Blei et al. (2017) and the references therein. Such procedure produces a distribution

other than the fractional posterior distribution. To accommodate such cases, we also give a theorem (Theorem 2 in Section 2) for the general measure Π and $\Pi^{(0)}$ in (3).

To illustrate the behavior of the ILRT under moderately complex models, we consider testing the homogeneity in two submodels of the normal mixture model. For the first submodel, the likelihood is unbounded and thus the LRT is not be defined. For the second submodel, Hall and Stewart (2005) showed that the LRT has trivial power under $n^{-1/2}$ local alternative hypothesis. In contrast, we prove that the ILRT has good asymptotic power behavior for both submodels.

The paper is organized as follow. In Section 2, we prove the Wilks phenomenon of the ILRT statistic and gives the asymptotic local power of the corresponding test. In Section 3, we apply ILRT to testing the homogeneity in two submodels of normal mixture model. Section 4 concludes the paper. All technical proofs are in Appendix.

Uniformly most powerful Bayesian tests.

Clarke and Barron (1990) considered use Bayes factor for hypothesis testing. ? gave.

We avoid the use of the maximum likelihood estimator. This allows us to apply the methods to the case where the MLE does not exist.

Our work is not purely of theoretical interest. It also has implication on the choice of prior and Bayesian methods. As Clarke and Barron (1990) noted, Bayes' Consistency holds under a hypothesis that is much weaker than the conditions for the consistency of the MLE.

The most remarkable property of LRT is the Wilks phenomenon. That is, the asymptotic distribution of the LRT statistic is free of parameters.

One advantage of the Bayes factor over the LRT is that the Bayes factor is always well defined. sensitive to outliers as FBF paper discussed.

Motivations

- Applicable to moderately complex models, espially when the likelihood is unbounded.
- Give guidines to the choice of the threshold of the Bayes factors.
- Computation is easy for some cases.

They proposed three ways to set b : (a) $b = m_0/n$; (b) $b = n^{-1} \max \{m_0, \sqrt{n}\}$; (c) $b = n^{-1} \max \{m_0, \log n\}$. Here m_0 is the minimal training sample size.

Let

$$\pi_t(\theta|\mathbf{X}^n) = \frac{\exp\{-tR_n(\theta_0|\theta)\}\pi(\theta)}{\int_{\Theta} \exp\{-tR_n(\theta_0|\theta)\}\pi(\theta) d\theta}$$

be the fractional posterior density with exponent t .

\mathbf{I}_{p_0} denote the p_0 dimensional identity matrix

“ \rightsquigarrow ” means weak convergence.

For general models, the likelihood function may not be concave. This often makes it hard to implement the LRT. For some models, a more serious problem may occur, that is, the likelihood

may be unbounded and hence the LRT can not be defined. This problem may occur even if the likelihood function has good local analytical properties, such as location-scale mixture models. See Le Cam (1990) for more examples.

The Wilks phenomenon is desirable since it allows us to determine the critical value according to the asymptotic null distribution. The lack of Wilks phenomenon will often introduce additional complexity to estimate the nuisance parameters ν under the null hypothesis.

Difficulties:

- Threshold
- Computation
- Choice of Prior

2 Wilks phenomenon of Bayes factors

Let $\mathbf{X}^n = (X_1, \dots, X_n)$ be independent identically distributed (iid) observations taking values in a standard measurable space $(\mathcal{X}; \mathcal{A})$. Suppose that the possible distribution P_θ of X_i has a density $p(X|\theta)$ with respect to μ , a σ -finite measure on \mathcal{X} . Denote by P_θ^n the joint distribution of \mathbf{X}^n . Let $p_n(\mathbf{X}^n|\theta) = \prod_{i=1}^n p(X_i|\theta)$ denote the density of P_θ^n with respect to the n -fold product measure μ^n . The parameter θ takes its values in Θ , an open subset of \mathbb{R}^p . Suppose $\theta = (\nu^\top, \xi^\top)^\top$, where ν is a p_0 dimensional subvector and ξ is a $p - p_0$ dimensional subvector. We would like to test the hypotheses

$$H : \theta \in \Theta_0 \quad \text{v.s.} \quad K : \theta \in \Theta \setminus \Theta_0,$$

where the null space Θ_0 is a p_0 -dimensional subspace of Θ defined as

$$\Theta_0 = \{(\nu^\top, \xi^\top)^\top : (\nu^\top, \xi^\top)^\top \in \Theta, \xi = \xi_0\}.$$

If the null hypothesis is true, we denote by $\theta_0 = (\nu_0^\top, \xi_0^\top)^\top$ the true parameter which generates the data.

In Bayesian hypothesis testing framework, a fundamental tool is Bayes factor

$$\text{BF}(\mathbf{X}^n) = \frac{\int_{\Theta} p_n(\mathbf{X}^n|\theta) \pi(\theta) d\theta}{\int_{\tilde{\Theta}_0} p_n(\mathbf{X}^n|\nu, \xi_0) \pi_0(\nu) d\nu},$$

where $\tilde{\Theta}_0 = \{\nu : (\nu^\top, \xi_0^\top)^\top \in \Theta_0\}$ and $\pi(\theta)$ and $\pi_0(\nu)$ are the possibly improper prior densities of parameters under the alternative and the null hypotheses, respectively. Conventionally, the null hypothesis is rejected if $\text{BF}(\mathbf{X}^n)$ is larger than certain threshold. The choice of threshold is mostly empirical in the literature. For example, Jeffreys (1961) suggested that the evidence against the null hypothesis is *decisive* if $\text{BF}(\mathbf{X}^n) > 100$ while Kass and Raftery (1995) suggested that the evidence is *very strong* if $\text{BF}(\mathbf{X}^n) > 150$. Unfortunately, these choices of threshold are not theoretically justified. Worse still, for improper priors, these choices of threshold suffer from the

Lindley paradox, that is, the resulting test procedure largely depends on the arbitrary constants in priors densities; see, e.g., Shafer (1982).

In this paper, we treat the Bayes factor as a frequentist test statistic and the threshold is chosen to control the type I error rate. To achieve this, we need to investigate the asymptotic distribution of Bayes factor. We make the following assumption which is adapted from Kleijn and van der Vaart (2012).

Assumption 1. *The parameter spaces Θ and $\tilde{\Theta}_0$ are open subsets of \mathbb{R}^p and \mathbb{R}^{p_0} , respectively. The parameters θ_0 and ν_0 are inner points of Θ and $\tilde{\Theta}_0$, respectively. The derivative*

$$\dot{\ell}_{\theta_0}(X) = \frac{\partial}{\partial \theta} \log p(X|\theta) \Big|_{\theta=\theta_0}$$

exists P_{θ_0} -a.s. and satisfies $P_{\theta_0} \dot{\ell}_{\theta_0} = \mathbf{0}_p$, where Pf means the expectation of $f(X)$ when X has distribution P . The Fisher information matrix $I(\theta_0) = P_{\theta_0} \dot{\ell}_{\theta_0} \dot{\ell}_{\theta_0}^\top$ is positive-definite. For every $M > 0$,

$$\sup_{\|h\| \leq M} \left| R_n(\theta_0 \| \theta_0 + n^{-1/2}h) + h^\top I(\theta_0) \Delta_{n,\theta_0} - \frac{1}{2} h^\top I(\theta_0) h \right| \xrightarrow{P_{\theta_0}^n} 0,$$

where $R_n(\theta \| \theta') = \log \{p_n(\mathbf{X}^n | \theta) / p_n(\mathbf{X}^n | \theta')\}$ is the log-likelihood ratio between $p_n(\mathbf{X}^n | \theta)$ and $p_n(\mathbf{X}^n | \theta')$, and $\Delta_{n,\theta_0} = n^{-1/2} \sum_{i=1}^n I(\theta_0)^{-1} \dot{\ell}_{\theta_0}(X_i)$.

Assumption 1 assumes that the likelihood function $p_n(\mathbf{X}^n | \theta)$ has good local behavior when θ is close to θ_0 . Although Assumption 1 is about the full likelihood function, it also implies the analogous assumption for the null likelihood function $p_n(\mathbf{X}^n | \nu, \xi_0)$. To see this, let $\mathbf{J} = (\mathbf{I}_{p_0}, \mathbf{0}_{p_0 \times (p-p_0)})^\top$,

$$\dot{\ell}_{\theta_0}^{(0)}(X) = \mathbf{J}^\top \dot{\ell}_{\theta_0}(X), \quad I_\nu(\theta) = \mathbf{J}^\top I(\theta) \mathbf{J}, \quad \Delta_{n,\theta_0}^{(0)} = \frac{1}{\sqrt{n}} \sum_{i=1}^n I_\nu(\theta_0)^{-1} \dot{\ell}_{\theta_0}^{(0)}(X_i).$$

Then Assumption 1 implies that

$$\sup_{\|h\| \leq M} \left| R_n(\theta_0 \| \nu_0 + n^{-1/2}h, \xi_0) + h^\top I_\nu(\theta_0) \Delta_{n,\theta_0}^{(0)} - \frac{1}{2} h^\top I_\nu(\theta_0) h \right| \xrightarrow{P_{\theta_0}^n} 0.$$

Note that Assumption 1 imposes no condition on the likelihood function when θ is deviated from θ_0 . In the theory of LRT, it is assumed that the maximum likelihood estimator exists and is consistent; see, e.g., Wilks (1938) and van der Vaart (1998), Theorem 16.7. This condition essentially assumes that the likelihood value is negligible when θ is deviated from θ_0 , and is not true for some important models. In particular, this does not allow that the likelihood function is unbounded. Hence we will not assume this strong condition. Instead, we shall utilize the \sqrt{n} -consistency of the posterior density $\pi(\theta | \mathbf{X}^n)$. We say $\pi(\theta | \mathbf{X}^n)$ is \sqrt{n} -consistent if for every $M_n \rightarrow \infty$,

$$\int_{\{\theta: \|\theta - \theta_0\| > M_n / \sqrt{n}\}} \pi(\theta | \mathbf{X}^n) d\theta \xrightarrow{P_{\theta_0}^n} 0.$$

The \sqrt{n} -consistency of $\pi_0(\nu|\mathbf{X}^n)$ is similarly defined.

We shall derive the asymptotic distribution of $\text{BF}(\mathbf{X}^n)$ under the null hypothesis as well as the local alternative hypothesis. By local alternative hypothesis we means that the true parameter is θ_n and $\sqrt{n}(\theta_n - \theta_0) \rightarrow \eta$. Let

$$I_{\xi|\nu}(\theta) = \tilde{\mathbf{J}}^\top I(\theta) \tilde{\mathbf{J}} - \tilde{\mathbf{J}}^\top I(\theta) \mathbf{J} (\mathbf{J}^\top I(\theta) \mathbf{J})^{-1} \mathbf{J}^\top I(\theta) \tilde{\mathbf{J}},$$

where $\tilde{\mathbf{J}} = (\mathbf{0}_{(p-p_0) \times p_0}, \mathbf{I}_{(p-p_0)})^\top$. It can be seen that $|I(\theta)| = |I_\nu(\theta)| \cdot |I_{\xi|\nu}(\theta)|$. Let $\chi^2(p - p_0, \eta^\top \tilde{\mathbf{J}} I_{\xi|\nu}(\theta_0) \tilde{\mathbf{J}}^\top \eta)$ denote a noncentral chi-squared random variable with $p - p_0$ degrees of freedom and noncentrality parameter $\eta^\top \tilde{\mathbf{J}} I_{\xi|\nu}(\theta_0) \tilde{\mathbf{J}}^\top \eta$. The following proposition gives the asymptotic distribution of Bayes factor.

Proposition 1. *Suppose that Assumption 1 holds, $\pi(\theta)$ is continuous at θ_0 with $\pi(\theta_0) > 0$, $\pi_0(\nu)$ is continuous at ν_0 with $\pi_0(\nu_0) > 0$, $\pi(\theta|\mathbf{X}^n)$, $\pi_0(\nu|\mathbf{X}^n)$ are \sqrt{n} -consistent. Suppose $\{\theta_n\}$ satisfies $\sqrt{n}(\theta_n - \theta_0) \rightarrow \eta$. Then*

$$2 \log \text{BF}(\mathbf{X}^n) + (p - p_0) \log \left(\frac{n}{2\pi} \right) \overset{P_{\theta_n}^n}{\rightsquigarrow} \chi^2(p - p_0, \eta^\top \tilde{\mathbf{J}} I_{\xi|\nu}(\theta_0) \tilde{\mathbf{J}}^\top \eta) + 2 \log \frac{|I_{\xi|\nu}(\theta_0)|^{-\frac{1}{2}} \pi(\theta_0)}{\pi_0(\nu_0)}.$$

Proposition 1 implies that the asymptotic null distribution of the Bayes factor depend on the nuisance parameters and the priors. Hence the Bayes factor does not have Wilks phenomenon for general priors. Nevertheless, the dependence of the asymptotic null distribution on the nuisance parameters can be cancelled by a class of priors. In fact, Proposition 1 implies that the asymptotic distribution of $\text{BF}_t(\mathbf{X}^n)$ is free of the nuisance parameter ν if and only if

$$\frac{|I_{\xi|\nu}(\nu, \xi_0)|^{-\frac{1}{2}} \pi(\theta_0)}{\pi_0(\nu)} \equiv c \quad (4)$$

for some constant c and for any $\nu \in \tilde{\Theta}$. Unfortunately, simple examples show that (4) does not hold for many popular objective priors for bayes factor, including intrinsic priors (Berger and Pericchi, 1996), fractional intrinsic priors (De Santis and Spezzaferri, 1997), divergence-based priors (Bayarri and Garca-Donato, 2008), expected-posterior priors (Perez, 2002). Nevertheless, (4) holds for a large class of priors. For instance, the left hand side of (4) is equal to 1 if $\pi_0(\nu) = |I_\nu(\nu, \xi_0)|^{1/2}$ and $\pi(\theta) = |I(\theta)|^{1/2}$, that is, $\pi_0(\nu)$ and $\pi(\theta)$ are the Jeffrey's priors under the null and the alternative hypotheses, respectively. In general, (4) is constant provided $\pi(\theta) = \pi(\xi|\nu)\pi(\nu)$ satisfies

$$\pi(\nu) = \pi_0(\nu), \quad \pi(\xi|\nu) \propto |I_{\xi|\nu}(\nu, \xi_0)|^{1/2}. \quad (5)$$

A class of priors satisfying (5) is the unit information priors proposed by Kass and Wasserman (1995), which is defined as

$$\pi(\nu) = \pi_0(\nu), \quad \pi(\xi|\nu) = |\Sigma_\xi(\nu)|^{-1/2} f \left((\xi - \xi_0)^\top \Sigma_\xi(\nu)^{-1} (\xi - \xi_0) \right),$$

where $\Sigma_\xi(\nu)$ satisfies $|\Sigma_\xi(\nu)| = |I_{\xi|\nu}(\nu, \xi_0)|^{-1}$.

We have seen that with carefully chosen priors, Bayes factor has Wilks phenomenon. In this case, Bayes factor can be treated as a frequentist statistic and the critical value is determined by the asymptotic distribution. However, this approach still has some weaknesses. First, the Wilks phenomenon of Bayes factor holds only for a restricted class of priors. As we have already pointed out, many popular objective priors do not satisfy (5). Also, for some models, Fisher information matrix has a complicated form. In this case, priors satisfying (5) may be undesirable. Second, the computation of Bayes factor is a difficult problem in general; see Friel and Wyse (2012) for a review on the computation of Bayes factor.

We turn to the variants of Bayes factor. We denote by $L_t(\mathbf{X}^n) = \int_{\Theta} [p_n(\mathbf{X}^n|\theta)]^t \pi(\theta) d\theta$ the power marginal likelihood with power $t > 0$, and similarly define $L_t^{(0)}(\mathbf{X}^n)$. For $t > 0$, let $\text{BF}_t(\mathbf{X}^n) = L_t(\mathbf{X}^n)/L_t^{(0)}(\mathbf{X}^n)$. Then $\text{BF}_1(\mathbf{X}^n)$ is the usual Bayes factor. It can be expected that $\text{BF}_t(\mathbf{X}^n)$ depends on the nuisance parameters and the priors in the same way for all $t > 0$. Hence the ratio of $\text{BF}_t(\mathbf{X}^n)$ for two different t may cancel the dependence on the nuisance parameters and the priors. For $a > b > 0$, let $\Lambda_{a,b}(\mathbf{X}^n) = \text{BF}_a(\mathbf{X}^n)/\text{BF}_b(\mathbf{X}^n)$. This class of statistics includes two important variants of Bayes factor, namely, the posterior Bayes factor proposed by Aitkin (1991) and the fractional Bayes factor proposed by O'Hagan (1995). In fact, the posterior Bayes factor is equal to $\Lambda_{2,1}(\mathbf{X}^n)$ and the fractional Bayes factor is equal to $\Lambda_{1,b}(\mathbf{X}^n)$ for $b \in (0, 1)$. For this reason we shall call $\Lambda_{a,b}(\mathbf{X}^n)$ the generalized fractional Bayes factor throughout the paper.

Since we treat $\Lambda_{a,b}(\mathbf{X}^n)$ as a frequentist statistic, the Wilks phenomenon of $\Lambda_{a,b}(\mathbf{X}^n)$ should be examined. Note that the posterior Bayes factor uses fixed $a = 2, b = 1$ for all n . On the other hand, the fractional Bayes factor uses fixed $a = 1$ but a varying b as $n \rightarrow \infty$. In fact, O'Hagan (1995) proposed three ways to set b , the first one is $b = m_0/n$ for a fixed m_0 , the second one is $b = n^{-1/2}$ and the third one is $b = \log(n)/n$. In what follows, we shall derive the asymptotic distribution of $\Lambda_{a,b}(\mathbf{X}^n)$ in three settings. In all three settings, a is fixed. In the first setting, b is also fixed. In the second setting, $b \rightarrow 0$ and $nb \rightarrow \infty$. In the third setting, $nb \rightarrow b^* \in (0, +\infty)$.

Assumption 2. For any fixed $t \in (0, 1]$, we assume $D_t(\theta_0||\theta)$ satisfies the following conditions:

- $D_1(\theta_0||\theta)$ is finite for all $\theta \in \Theta$;
- for each $\delta > 0$, there exists a $\epsilon > 0$ such that $D_t(\theta_0||\theta) \geq \epsilon$ for $\|\theta - \theta_0\| \geq \delta$;
- as $\theta \rightarrow \theta_0$,

$$D_t(\theta_0||\theta) = (1 + o(1)) \frac{t}{2} (\theta - \theta_0)^\top I(\theta_0) (\theta - \theta_0), \quad V(\theta_0||\theta) = O(\|\theta - \theta_0\|^2).$$

See, e.g., van Erven and Harremoës (2014).

$$V(\theta_0||\theta) = P_{\theta_0}(\log(p(X|\theta_0)/p(X|\theta)) - D_1(\theta_0||\theta))^2.$$

Assumption 3. We assume $\pi(\theta)$ is continuous at θ_0 with $\pi(\theta_0) > 0$, $\pi_0(\nu)$ is continuous at ν_0 with $\pi_0(\nu_0) > 0$. For every $t^* \in (0, 1)$, there exist $c^*, c^\dagger > 0$, $t_0^* \in (0, 1)$ and $c_0^*, c_0^\dagger > 0$,

$$\int_{\Theta} \exp \{-c^* D_{1-t^*}(\theta_0 \parallel \theta)\} \pi(\theta) d\theta < \infty, \quad \int_{\Theta} V(\theta_0 \parallel \theta) \exp \{-c^\dagger D_1(\theta_0 \parallel \theta)\} \pi(\theta) d\theta < \infty,$$

$$\int_{\tilde{\Theta}_0} \exp \{-c_0^* D_{1-t_0^*}(\nu_0 \parallel \nu)\} \pi_0(\nu) d\nu < \infty, \quad \int_{\tilde{\Theta}_0} V(\nu_0 \parallel \nu) \exp \{-c_0^\dagger D_1(\nu_0 \parallel \nu)\} \pi_0(\nu) d\nu < \infty.$$

Theorem 1. Suppose that Assumptions 1, 2, 3 hold. Suppose $\{\theta_n\}$ satisfies $\sqrt{n}(\theta_n - \theta_0) \rightarrow \eta$. Then the following assertions hold.

1. Suppose $a > b > 0$ are fixed numbers, $L_a(\cdot; \mathbf{X}^n)$, $L_b(\cdot; \mathbf{X}^n)$, $L_a^{(0)}(\cdot; \mathbf{X}^n)$ and $L_b^{(0)}(\cdot; \mathbf{X}^n)$ are \sqrt{n} -consistent. Then

$$2 \log \Lambda_{a,b} + (p - p_0) \log \left(\frac{a}{b} \right) \overset{P_{\theta_n}^n}{\rightsquigarrow} (a - b) \chi^2(p - p_0, \delta).$$

2. Suppose $a > b > 0$, a is fixed and as $n \rightarrow \infty$, $b \rightarrow 0$, $nb \rightarrow b^* \in (c^*, +\infty)$. Then

$$2 \log \Lambda_{a,b} + (p - p_0) \log \left(\frac{an}{2\pi} \right) \overset{P_{\theta_n}^n}{\rightsquigarrow} a \chi^2(p - p_0, \delta) + 2 \log \left(\frac{|I_{\xi|\nu}(\theta_0)|^{-\frac{1}{2}} \pi(\theta_0)}{\pi_0(\nu_0)} \right) \\ - 2 \log \left(\frac{\int_{\Theta} \exp \{-b^* D_1(\theta_0 \parallel \theta)\} \pi(\theta) d\theta}{\log \int_{\tilde{\Theta}_0} \exp \{-b^* D_1(\theta_0 \parallel \nu, \xi_0)\} \pi_0(\nu) d\nu} \right).$$

3. Suppose $a > b > 0$, a is fixed and as $n \rightarrow \infty$, $b \rightarrow 0$, $bn \rightarrow \infty$. Then

$$2 \log \Lambda_{a,b} + (p - p_0) \log \left(\frac{a}{b} \right) \overset{P_{\theta_n}^n}{\rightsquigarrow} a \chi^2(p - p_0, \delta).$$

Theorem 1 gives the asymptotic distribution of $\Lambda_{a,b}$ under the null hypothesis and the local alternative hypothesis. It can be seen that when $nb \rightarrow \infty$, $\Lambda_{a,b}$ has Wilks phenomenon for any prior distributions. To formulate a test with asymptotic type I error rate α , the critical value of $2 \log \Lambda_{a,b}$ can be defined to be $-(p - p_0) \log(1 + a/b) + a \chi_\alpha^2(p - p_0)$ where $\chi_\alpha^2(p - p_0)$ is the $1 - \alpha$ quantile of a chi-squared random variable with $p - p_0$ degrees of freedom. By Theorem 1, the resulting test has local asymptotic power

$$\Pr(\chi^2(p - p_0, \delta) > \chi_\alpha^2(p - p_0)). \quad (6)$$

It is known that, under certain regular conditions, (6) is also the local asymptotic power of the likelihood ratio test. In this view, $\Lambda_{a,b}$ enjoys good frequentist properties.

LRT essentially require the MLE. For Bayes factor, we need not estimate any parameter.

3 General integrated likelihood ratio test

It is known that the computation of Bayes factor is not trivial even if it is easy to sample from the posterior distribution. Surprisingly, $\Lambda_{a,b}$ can be easily computed provided sampling from the posterior distribution is feasible. The computation of the ILRT statistic is relatively simple. It can be seen that $\Lambda_{a,b}$ can be written as

$$\Lambda_{a,b} = \frac{\int_{\Theta} [p_n(\mathbf{X}^n|\theta)]^{a-b} \pi_b(\theta|\mathbf{X}^n) d\theta}{\int_{\tilde{\Theta}_0} [p_n(\mathbf{X}^n|\nu, \xi_0)]^{a-b} \pi_{b,0}(\nu|\mathbf{X}^n) d\nu}.$$

We can independently generate $\theta_1, \dots, \theta_m$ and ν_1, \dots, ν_m according to $\pi_b(\theta|\mathbf{X}^n)$ and $\pi_{b,0}(\nu; \mathbf{X}^n)$ for a large m . Then the ILRT statistic can be approximated by

$$\frac{\sum_{i=1}^m [p_n(\mathbf{X}^n|\theta_i)]^a}{\sum_{i=1}^m [p_n(\mathbf{X}^n|\nu_i, \xi_0)]^a}.$$

Note that $\pi(\theta; \mathbf{X}^n)$ and $\pi_0(\nu; \mathbf{X}^n)$ are data dependent but does not need to be the posterior density. Under Assumption 4, we consider the ILRT statistic

$$\Lambda_{a,b}^* = \frac{\int_{\Theta} [p_n(\mathbf{X}^n|\theta)]^a \pi(\theta; \mathbf{X}^n) d\theta}{\int_{\tilde{\Theta}_0} [p_n(\mathbf{X}^n|\nu, \xi_0)]^a \pi_0(\nu; \mathbf{X}^n) d\nu}. \quad (7)$$

If we take the weight functions as the fractional posteriors

$$\begin{aligned} \pi(\theta; \mathbf{X}^n) &= \frac{[p_n(\mathbf{X}^n|\theta)]^b \pi(\theta)}{\int_{\Theta} [p_n(\mathbf{X}^n|\theta)]^b \pi(\theta) d\theta}, \\ \pi_0(\nu; \mathbf{X}^n) &= \frac{[p_n(\mathbf{X}^n|\nu, \xi_0)]^b \pi_0(\nu)}{\int_{\tilde{\Theta}_0} [p_n(\mathbf{X}^n|\nu, \xi_0)]^b \pi_0(\nu) d\nu}, \end{aligned} \quad (8)$$

For some moderately complex models, the fractional posterior (8) may be complicated, and it is not easy to sample from it. In this case, one may use some simple form weight function to approximate the fractional posterior (8). A popular method for approximating (8) is variational inference; see, e.g., Blei et al. (2017). In this case, the weight functions in the integrated likelihood ratio test statistic are the variational approximations of the fractional posteriors (8). Let $h = \sqrt{n}(\theta - \theta_0)$ be the local parameter and $\pi_n(h; \mathbf{X}^n) = \pi(\theta_0 + n^{-1/2}h; \mathbf{X}^n)$ be the weight function in terms of h . If $\pi(\theta; \mathbf{X}^n)$ is the posterior density of θ , then Bernstein-von Mises theorem asserts that under certain conditions, $\|\pi_n(h; \mathbf{X}^n) - \phi(h; \Delta_{n,\theta_0}, I_{\theta_0}^{-1})\|$ converges to 0 in $P_{\theta_0}^n$ probability, where for two densities $q_1(h)$ and $q_2(h)$, $\|q_1(h) - q_2(h)\| = \int |q_1(h) - q_2(h)| dh$ is their total variation distance. Similarly, if $\pi(\theta; \mathbf{X}^n)$ is the fractional posterior density of θ with fractional power b , it can be proved that under certain conditions, $\|\pi_n(h; \mathbf{X}^n) - \phi(h; \Delta_{n,\theta_0}, b^{-1}I_{\theta_0}^{-1})\|$ converges to 0 in $P_{\theta_0}^n$ probability. We shall assume that the weight function inherits such properties.

Assumption 4. Let $b \in (0, 1)$ be a fixed number. Assume that $\pi_n(h; \mathbf{X}^n)$ satisfies

$$\|\pi_n(h; \mathbf{X}^n) - \phi(h; \Delta_{n,\theta_0}, b^{-1}I_{\theta_0}^{-1})\| \xrightarrow{P_{\theta_0}^n} 0. \quad (9)$$

Similarly, let $h^{(0)} = \sqrt{n}(\nu - \nu_0)$. Define $\pi_n(h^{(0)}; \mathbf{X}^n) = n^{-1/2}\pi_0(\nu; \mathbf{X}^n)$. Assume that

$$\|\pi_n(h^{(0)}; \mathbf{X}^n) - \phi(h^{(0)}; \Delta_{n,\theta_0}^{(0)}, b^{-1}I_{\theta_0}^{(0)-1})\| \xrightarrow{P_{\theta_0}^n} 0, \quad (10)$$

where

$$\Delta_{n,\theta_0}^{(0)} = \frac{1}{\sqrt{n}} \sum_{i=1}^n I_{\theta_0}^{(0)-1} \dot{\ell}_{\theta_0}^{(0)}(X_i), \quad I_{\theta_0}^{(0)} = P_{\theta_0} \dot{\ell}_{\theta_0}^{(0)} \dot{\ell}_{\theta_0}^{(0)\top}, \quad \dot{\ell}^{(0)}(X) = \frac{\partial}{\partial \nu} \log p(X|\nu, \xi_0) \Big|_{\nu=\nu_0}.$$

Furthermore, assume that for every $\epsilon > 0$, there exist Lebesgue integrable functions $T(h)$ and $T^{(0)}(h)$ such that

$$\liminf_{n \rightarrow \infty} P_{\theta_0}^n \left\{ \sup_{h \in \mathbb{R}^p} (\pi_n(h; \mathbf{X}^n) - T(h)) \leq 0 \right\} \geq 1 - \epsilon. \quad (11)$$

$$\liminf_{n \rightarrow \infty} P_{\theta_0}^n \left\{ \sup_{h^{(0)} \in \mathbb{R}^{p_0}} (\pi_n(h^{(0)}; \mathbf{X}^n) - T^{(0)}(h^{(0)})) \leq 0 \right\} \geq 1 - \epsilon. \quad (12)$$

The conditions (11) and (12) assume that there is a function controlling the tail of the weight functions. We need to control the tail of the weight function since the behavior of the likelihood may be undesirable when θ is far away from θ_0 . In fact, even for some fairly regular models, the likelihood may tends to infinity, which invalidates the LRT; see, e.g., Le Cam (1990). So we control the tail of the weight function to avoid too much weights on the tail of likelihood. If the weight function $\pi_n(h; \mathbf{X}^n)$ is the normal density, then it can be shown that the conditions (9) and (10) implies (11) and (12).

The following theorem gives the asymptotic distribution of ILRT statistic.

Theorem 2. Suppose that Assumptions 1 and 4 hold with $a + b \leq 1$. Then for $\{\theta_n\}$ such that $\sqrt{n}(\theta_n - \theta_0) \rightarrow \eta$, we have

$$2 \log \Lambda_{a,b}^* \overset{P_{\theta_n}^n}{\rightsquigarrow} -(p - p_0) \log(1 + \frac{a}{b}) + a\chi^2(p - p_0, \delta),$$

where δ is defined as in Theorem 1.

Theorem 2 shows that even with approximate weight function, the ILRT statistic can still produce an asymptotic optimal test. A practical method to obtain simple form weight function $\pi_n(h; \mathbf{X}^n)$ is the variational inference; see, e.g., Blei et al. (2017). Next we shall consider a simple variational method which is guaranteed to yield a weight function satisfying Assumption 4. For comprehensive considerations of the statistical properties of variational methods; see the recent works of Wang and Blei (2017), Pati et al. (2017) and Yang et al. (2017).

Let \mathcal{Q} be the family of all p dimensional normal distribution. Let $\pi(\theta; \mathbf{X}^n)$ be the fractional posterior of order b and $\pi_n(h; \mathbf{X}^n) = n^{-1/2}\pi(\theta_0 + n^{-1/2}h; \mathbf{X}^n)$ be the corresponding fractional posterior of h . Suppose that $\pi_n(h; \mathbf{X}^n)$ satisfies

$$\|\pi_n(h; \mathbf{X}^n) - \phi(h; \Delta_{n,\theta_0}, b^{-1}I_{\theta_0}^{-1})\| \xrightarrow{P_{\theta_0}^n} 0. \quad (13)$$

Let the weight function $\pi^\dagger(\theta; \mathbf{X}^n)$ be the normal approximation of $\pi(\theta; \mathbf{X}^n)$ obtained from Rényi divergence variational inference (Li and Turner, 2016), that is,

$$\pi^\dagger(\theta; \mathbf{X}^n) = \arg \min_{q(\theta) \in \mathcal{Q}} -\frac{1}{1-\alpha} \log \int q(\theta)^\alpha \pi(\theta; \mathbf{X}^n)^{1-\alpha} d\theta,$$

where $0 < \alpha < 1$ is an arbitrary constant. Let $\pi_n^\dagger(h; \mathbf{X}^n) = n^{-1/2} \pi^\dagger(\theta_0 + n^{-1/2}h; \mathbf{X}^n)$ be the weight function of h . It can be seen that

$$\pi_n^\dagger(h; \mathbf{X}^n) = \arg \min_{q(h) \in \mathcal{Q}} -\frac{1}{1-\alpha} \log \int q(h)^\alpha \pi_n(h; \mathbf{X}^n)^{1-\alpha} dh.$$

Hence we have

$$\begin{aligned} & -\frac{1}{1-\alpha} \log \int \pi_n^\dagger(h; \mathbf{X}^n)^\alpha \pi_n(h; \mathbf{X}^n)^{1-\alpha} dh \\ & \leq -\frac{1}{1-\alpha} \log \int \phi(h; \Delta_{n, \theta_0}, I_{\theta_0}^{-1})^\alpha \pi_n(h; \mathbf{X}^n)^{1-\alpha} dh. \end{aligned} \tag{14}$$

Since Rényi divergence and total variation distance are topologically equivalent, (13) implies that the right hand side of (14) tends to 0 in $P_{\theta_0}^n$ -probability. Again by the topological equivalence of Rényi divergence and total variation distance, we have

$$\|\pi_n^\dagger(h; \mathbf{X}^n) - \phi(h; \Delta_{n, \theta_0}, b^{-1} I_{\theta_0}^{-1})\| \xrightarrow{P_{\theta_0}^n} 0.$$

Note that $\pi_n^\dagger(h; \mathbf{X}^n)$ and $\phi(h; \Delta_{n, \theta_0}, b^{-1} I_{\theta_0}^{-1})$ are both normal density functions. For normal distributions, the convergence in total variation implies the convergence of parameters. Hence the mean and covariance parameters of $\pi_n^\dagger(h; \mathbf{X}^n)$ are bounded in probability. Then a dominating function $T(h)$ exists and thus (11) holds.

4 \sqrt{n} -consistency of power posterior

The \sqrt{n} -consistency of $L_t(\cdot; \mathbf{X}^n)$ is a key assumption of Theorem 1. Hence we would like to give sufficient conditions for the \sqrt{n} -consistency of $L_t(\cdot; \mathbf{X}^n)$. First we consider the exponential family of distributions.

Proposition 2. *Suppose $p(X|\theta) = \exp[\theta^\top T(X) - A(\theta)]$, Θ is an open subset of \mathbb{R}^p , θ_0 is an interior point of Θ , $I(\theta_0)$ is positive-definite. Suppose the prior density $\pi(\theta)$ is continuous at θ_0 with $\pi(\theta_0) > 0$, and there exist $c^* > 0$ such that*

$$\int_{\Theta} \exp\{-c^* \|\theta - \theta_0\|\} \pi(\theta) d\theta < \infty.$$

Then $L_t(\cdot; \mathbf{X}^n)$ is consistent for any fixed $t > 0$.

Proposition 2 establishes the \sqrt{n} -consistency of $L_t(\cdot; \mathbf{X}^n)$ for all $t > 0$ under full-rank exponential family models. The success in the full-rank exponential family models is just a minimal requirement. We would like to consider more general models.

For general models, a fundamental question is whether the fractional posterior is always well defined with proper prior. The following theorem shows that $L_t(\Theta; \mathbf{X}^n)$ is always well defined for $t \leq 1$ and is not well defined for some model for $t > 1$.

Proposition 3. *If $t \leq 1$, $L_t(\Theta; \mathbf{X}^n) < +\infty$ $P_{\theta_0}^n$ -a.s. for any models. If $t > 1$, $L_t(\Theta; \mathbf{X}^n) = +\infty$ for some models.*

Because of the bad behavior of $L_t(\Theta; \mathbf{X}^n)$ for $t > 1$, we shall only consider $L_t(\Theta; \mathbf{X}^n)$ for $t \leq 1$. For $t = 1$, the \sqrt{n} -consistency of $L_t(\cdot; \mathbf{X}^n)$ is equivalent to the \sqrt{n} -consistency of the posterior distribution which is a well studied problem; see, e.g., Ghosal et al. (2000), Shen and Wasserman (2001), van der Vaart and Ghosal (2007). A popular and convenient way of establishing the consistency of posterior is through the condition that suitable test sequences exist. For example, Theorem 3.1 of Kleijn and van der Vaart (2012) assumes that for every $\epsilon > 0$, there exists a sequence of tests ϕ_n such that

$$P_{\theta_0}^n \phi_n \rightarrow 0, \quad \sup_{\|\theta - \theta_0\| \geq \epsilon} P_{\theta}^n (1 - \phi_n) \rightarrow 0. \quad (15)$$

This condition is satisfied when the parameter space is compact and the model is suitably continuous; see Theorem 3.2 of Kleijn and van der Vaart (2012). However, if the parameter space is not compact, one may have to manually construct a test sequence satisfying the condition (15).

The consistency of $L_t(\cdot; \mathbf{X}^n)$ for $0 < t < 1$ is different from $t = 1$. Walker and Hjort (2001) considered the Hellinger consistency of $L_{1/2}(\cdot; \mathbf{X}^n)$. They derived the consistency of $L_{1/2}(\cdot; \mathbf{X}^n)$ under simple conditions. Recently, Bhattacharya et al. (2019) further developed the idea of Walker and Hjort (2001) and derived a general bounds for the consistency of $L_t(\cdot; \mathbf{X}^n)$ for $0 < t < 1$. However, their result can not yield the \sqrt{n} -consistency for parametric models. We shall prove the \sqrt{n} -consistency of $L_t(\cdot; \mathbf{X}^n)$ for $0 < t < 1$ under certain conditions on the Rényi divergence between distributions in the family $\{P_{\theta} : \theta \in \Theta\}$.

For two parameters θ_1 and θ_2 , the α order Rényi divergence ($0 < \alpha < 1$) of P_{θ_1} from P_{θ_2} is defined to be

$$D_{\alpha}(\theta_1 || \theta_2) = -\frac{1}{1 - \alpha} \log \rho_{\alpha}(\theta_1, \theta_2),$$

where $\rho_{\alpha}(\theta_1, \theta_2) = \int_{\mathcal{X}} p(X|\theta_1)^{\alpha} p(X|\theta_2)^{1-\alpha} d\mu$ is the so-called Hellinger integral. The following assumption is needed for our \sqrt{n} -consistency result.

Assumption 5. *For some $\alpha \in (0, 1)$, there exist positive constants δ , ϵ and C such that, $D_{\alpha}(\theta_0 || \theta) \geq C \|\theta - \theta_0\|^2$ for $\|\theta - \theta_0\| \leq \delta$ and $D_{\alpha}(\theta_0 || \theta) \geq \epsilon$ for $\|\theta - \theta_0\| > \delta$.*

Remark 1. A remarkable property of Rényi divergence is the equivalence of all D_{α} . If $0 < \alpha < \beta < 1$, then

$$\frac{\alpha}{1 - \alpha} \frac{1 - \beta}{\beta} D_{\beta}(\theta_1 || \theta_2) \leq D_{\alpha}(\theta_1 || \theta_2) \leq D_{\beta}(\theta_1 || \theta_2).$$

See, e.g., van Erven and Harremoës (2014). As a result, if Assumption 5 holds for some $\alpha \in (0, 1)$, then it will hold for every $\alpha \in (0, 1)$.

Proposition 4. *Suppose θ_0 is an interior of Θ , $\pi(\theta)$ is continuous at θ_0 and $\pi(\theta_0) > 0$. Under Assumptions 1 and 2, for fixed $t \in (0, 1)$, $L_t(\cdot; \mathbf{X}^n)$ is consistent.*

Note that if the conditions of Theorem 1 are satisfied, the asymptotic power of $\Lambda_{a,b}$ is independent of a, b . Hence specific choices of a, b are not crucial provided $L_{a+b}(\cdot; \mathbf{X}^n)$, $L_b(\cdot; \mathbf{X}^n)$, $L_{a+b}^{(0)}(\cdot; \mathbf{X}^n)$ and $L_b^{(0)}(\cdot; \mathbf{X}^n)$ are \sqrt{n} -consistent. For some models, it is more convenient to verify Assumption 5 than to directly construct a test sequence satisfying the condition (15). In such cases, it can be recommended to use the generalized FBF with $a + b < 1$.

5 Exponential family

6 Normal mixture model

In this section, we apply the generalized fractional Bayes factor to testing the component number of normal mixture model. Normal mixture model is a highly irregular model. Due to partial loss of identifiability, the LRT has undesirable behavior. For example, if the component variances are totally unknown, the likelihood is unbounded and thus the LRT is not defined (Le Cam, 1990). See Chen (2017) for a review of the testing problems for mixture models. Since the integral of the likelihood can smooth the irregular behavior of the likelihood, it can be expected that Bayes methods may have better behavior than LRT. For example, for unknown variances case, ILRT is at least well defined.

Suppose X_1, \dots, X_n are iid distributed as a mixture of normal distributions

$$p(X|\omega, \xi, \sigma^2) = \frac{1-\omega}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}X^2\right) + \frac{\omega}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2}(X-\xi)^2\right),$$

where $0 \leq \omega \leq 1$, $\mu \in \mathbb{R}$ and $\sigma^2 \in \mathbb{R}^+$. First, we assume $\omega = 1/2$ is known and consider testing the hypotheses

$$H : \xi = 0, \sigma = 1 \quad \text{vs.} \quad K : \xi \neq 0 \text{ or } \sigma \neq 1. \quad (16)$$

For this testing problem, the likelihood function is unbounded under the alternative hypothesis. In fact, if we take $\xi = X_1$ and let $\sigma^2 \rightarrow 0$, then the likelihood tends to infinity. Thus, the LRT can not be defined. Figure 1 exhibits this phenomenon. Using Theorem 1 and Proposition 4, we can obtain the following proposition.

Proposition 5. *For hypotheses testing problem (16), if $\sqrt{n}((\xi, \sigma^2) - (0, 1))^\top \rightarrow (\eta_1, \eta_2)^\top$, then the generalized FBF with $a > 0$, $b > 0$ and $a + b < 1$ satisfies*

$$2 \log \Lambda_{a,b} \overset{P_{\theta_n}^n}{\rightsquigarrow} -2 \log\left(1 + \frac{a}{b}\right) + a\chi^2(2, \eta_1^2/4 + \eta_2^2/8).$$




Figure 1: Data X_1, \dots, X_n are iid from the mixture model $(1 - \omega)\mathcal{N}(0, 1) + \omega\mathcal{N}(\xi, \sigma^2)$ with $(\omega, \xi, \sigma^2)^\top = (1/2, 1, 10^{-2})$ and $n = 50$. We plot the likelihood function in $-\log(\sigma^2)$ with $\omega = 1/2$ and $\xi = X_1$. The likelihood tends to infinity as $-\log(\sigma^2)$ tends to infinity, i.e., σ^2 tends to 0. In contrast, the likelihood has a local maximum around the true parameter $-\log(\sigma^2) = -\log(10^{-2})$.

This example shows that even when the LRT fails, the ILRT can still be valid and has the expected asymptotic distribution. Thus, the ILRT methodology has a wider application scope than the LRT.

In the above example, we assume $\omega = 1/2$ is known. If ω is unknown, then the mixture model suffers from loss of identifiability and the behavior of the likelihood is fairly complicated. For simplicity, we assume $\sigma^2 = 1$ is known and consider testing the hypotheses

$$H : \omega\xi = 0 \quad \text{vs.} \quad K : \omega\xi \neq 0. \quad (17)$$

Although the LRT exists in this problem, its asymptotic behavior is complicated and its power behavior is not satisfactory. In fact, Hall and Stewart (2005) showed that the LRT has trivial power under $n^{-1/2}$ local alternative hypothesis. For this irregular problem, Theorem 1 and Proposition 4 cannot be directly applied. This is because the second part of Assumption (4) is violated due to loss of identifiability. However, this does not mean that the ILRT is not applicable. In fact, the following theorem shows that the generalized FBF with $a + b < 1$ has the desirable asymptotic properties.

Theorem 3. *Suppose $\pi(\omega, \xi) = \pi_\omega(\omega)\pi_\xi(\xi)$, $\pi_\xi(\xi)$ is positive and continuous at $\xi = 0$, $\pi_\omega(\omega) \sim \text{Beta}(\alpha_1, \alpha_2)$ with $\alpha_1 > 1$. Suppose $a + b < 1$. Then,*

(i) *under the null hypothesis,*

$$2 \log \Lambda_{a,b} \overset{P_{\theta_0}^n}{\rightsquigarrow} \log\left(1 + \frac{a}{b}\right) + a\chi^2(1);$$

(ii) *suppose for some $s < 1/4$, $\omega \geq n^{-s}$ for large n , $\sqrt{n}\omega\xi \rightarrow \eta$, then*

$$2 \log \Lambda_{a,b} \overset{P_{\theta_n}^n}{\rightsquigarrow} \log\left(1 + \frac{a}{b}\right) + a\chi^2(1, \eta^2).$$

Theorem 3 shows that the ILRT has nontrivial power if $\omega\xi$ is of order $n^{-1/2}$. In comparison, Hall and Stewart (2005) showed that the LRT has trivial power asymptotically if $\omega\xi = \gamma(n^{-1} \log \log n)^{1/2}$ with $|\gamma| < 1$.

7 Heteroscedastic regression model

Li and Chan (2000), Crisp and Burrige (1994).

8 Conclusion

In this paper, we proposed a flexible methodology ILRT which includes some existing methods as special cases. The asymptotic behaviors of the ILRT are investigated. It is shown that the ILRT

has the Wilks phenomenon similar to the LRT. The asymptotic local power is also given. We also apply the ILRT methodology to two submodels of the normal mixture model. These examples show that the ILRT can have good behavior even if the LRT is not defined or has poor properties.

The ILRT statistic is easy to implement provided sampling from weight functions is convenient. If the weight functions are fractional posterior densities, then Markov chain Monte Carlo (MCMC) methods can be used to sample from weight functions. Furthermore, if MCMC is not efficient, one can use approximation methods, such as variational inference, and the resulting test procedure is still valid. Thus, the ILRT methodology can also be recommended when the classical LRT is not easy to implement.

It is interesting to apply the ILRT methodology to specific complex testing problems. We leave it for future research.

Acknowledgements

This work was supported by the National Natural Science Foundation of China under Grant No. 11471035.

Appendices

Appendix A Preliminary lemmas

Lemma 1. *Suppose that Assumption 1 holds. Suppose $\{\theta_n\}$ satisfies $\sqrt{n}(\theta_n - \theta_0) \rightarrow \eta$. Then for any statistics T_n , $T_n = o_{P_{\theta_0}^n}(1)$ if and only if $T_n = o_{P_{\theta_n}^n}(1)$.*

Proof. Assumption 1 implies that $P_{\theta_0}^n$ and $P_{\theta_n}^n$ are mutually contiguous. Then the conclusion follows from Le Cam's first lemma (van der Vaart, 1998, Lemma 6.4). \square

Lemma 2. *Suppose that Assumption 1 holds. Suppose $\{\theta_n\}$ satisfies $\sqrt{n}(\theta_n - \theta_0) \rightarrow \eta$. Then*

$$\Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} - \Delta_{n,\theta_0}^{(0)\top} I_\nu(\theta_0) \Delta_{n,\theta_0}^{(0)} \stackrel{P_{\theta_n}^n}{\rightsquigarrow} \chi^2(p - p_0, \eta^\top \tilde{\mathbf{J}} I_{\xi|\nu}(\theta_0) \tilde{\mathbf{J}}^\top \eta).$$

Proof. It can be seen that $\Delta_{n,\theta_0}^{(0)} = (\mathbf{J}^\top I(\theta_0) \mathbf{J})^{-1} \mathbf{J}^\top I(\theta_0) \Delta_{n,\theta_0}$. Then

$$\begin{aligned} & \Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} - \Delta_{n,\theta_0}^{(0)\top} I_\nu(\theta_0) \Delta_{n,\theta_0}^{(0)} \\ &= \Delta_{n,\theta_0}^\top I(\theta_0)^{1/2} (\mathbf{I}_p - I(\theta_0)^{1/2} \mathbf{J} (\mathbf{J}^\top I(\theta_0) \mathbf{J})^{-1} \mathbf{J}^\top I(\theta_0)^{1/2}) I(\theta_0)^{1/2} \Delta_{n,\theta_0}, \end{aligned}$$

where $\mathbf{I}_p - I(\theta_0)^{1/2} \mathbf{J} (\mathbf{J}^\top I(\theta_0) \mathbf{J})^{-1} \mathbf{J}^\top I(\theta_0)^{1/2}$ is a projection matrix with rank $p - p_0$. It remains to derive the asymptotic distribution of Δ_{n,θ_0} . Let $h_n = \sqrt{n}(\theta_n - \theta_0)$. From Assumption 1 and the

central limit theorem, we have

$$\left(\frac{1}{\sqrt{n}} \sum_{i=1}^n \dot{\ell}_{\theta_0}(X_i) \right) \stackrel{P_{\theta_0}^n}{\rightsquigarrow} \mathcal{N} \left(\begin{pmatrix} 0 \\ -\frac{1}{2} \eta^\top I(\theta_0) \eta \end{pmatrix}, \begin{pmatrix} I(\theta_0) & I(\theta_0) \eta \\ \eta^\top I(\theta_0) & \eta^\top I(\theta_0) \eta \end{pmatrix} \right).$$

Then Le Cam's third lemma (van der Vaart, 1998, Example 6.7) implies that

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n \dot{\ell}_{\theta_0}(X_i) \stackrel{P_{\theta_0}^n}{\rightsquigarrow} \mathcal{N}(I(\theta_0) \eta, I(\theta_0)).$$

Consequently, Δ_{n,θ_0} weakly converges to $\mathcal{N}(\eta, I(\theta_0)^{-1})$ in $P_{\theta_0}^n$. It follows that

$$\Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} - \Delta_{n,\theta_0}^{(0)\top} I_\nu(\theta_0) \Delta_{n,\theta_0}^{(0)} \stackrel{P_{\theta_0}^n}{\rightsquigarrow} \chi^2(p - p_0, \delta),$$

where $\delta = \eta^\top (I(\theta_0) - I(\theta_0) \mathbf{J} (\mathbf{J}^\top I(\theta_0) \mathbf{J})^{-1} \mathbf{J}^\top I(\theta_0)) \eta$. Note that

$$(I(\theta_0) - I(\theta_0) \mathbf{J} (\mathbf{J}^\top I(\theta_0) \mathbf{J})^{-1} \mathbf{J}^\top I(\theta_0)) \mathbf{J} \mathbf{J}^\top = \mathbf{0}_{p \times p}.$$

Hence

$$\begin{aligned} \delta &= \eta^\top (\mathbf{J} \mathbf{J}^\top + \tilde{\mathbf{J}} \tilde{\mathbf{J}}^\top) (I(\theta_0) - I(\theta_0) \mathbf{J} (\mathbf{J}^\top I(\theta_0) \mathbf{J})^{-1} \mathbf{J}^\top I(\theta_0)) (\mathbf{J} \mathbf{J}^\top + \tilde{\mathbf{J}} \tilde{\mathbf{J}}^\top) \eta \\ &= \eta^\top \tilde{\mathbf{J}} \tilde{\mathbf{J}}^\top (I(\theta_0) - I(\theta_0) \mathbf{J} (\mathbf{J}^\top I(\theta_0) \mathbf{J})^{-1} \mathbf{J}^\top I(\theta_0)) \tilde{\mathbf{J}} \tilde{\mathbf{J}}^\top \eta \\ &= \eta^\top \tilde{\mathbf{J}} I_{\xi|\nu}(\theta_0) \tilde{\mathbf{J}}^\top \eta. \end{aligned}$$

This completes the proof. \square

Appendix B Proofs in Section 2

Proposition 6. Suppose that Assumption 1 holds, $t \in (0, +\infty)$ is fixed, $L_t(\cdot; \mathbf{X}^n)$ is \sqrt{n} -consistent, $\pi(\theta)$ is continuous at θ_0 with $\pi(\theta_0) > 0$. Then we have

$$\int_{\Theta} \exp \{-t R_n(\theta_0 \| \theta)\} \pi(\theta) d\theta = (1 + o_{P_{\theta_0}^n}(1)) \pi(\theta_0) \left(\frac{2\pi}{tn} \right)^{p/2} |I(\theta_0)|^{-1/2} \exp \left\{ \frac{t}{2} \Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} \right\}.$$

Proof. For any fixed $M > 0$, we have

$$\begin{aligned} & \int_{\{\theta: \|\theta - \theta_0\| \leq M/\sqrt{n}\}} \exp[-t R_n(\theta_0 \| \theta)] \pi(\theta) d\theta \\ &= (1 + o_{P_{\theta_0}^n}(1)) n^{-p/2} \pi(\theta_0) \int_{\{h: \|h\| \leq M\}} \exp \left[-t R_n(\theta_0 \| \theta_0 + n^{-1/2} h) \right] dh, \\ &= (1 + o_{P_{\theta_0}^n}(1)) n^{-p/2} \pi(\theta_0) \exp \left\{ \frac{t}{2} \Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} \right\} \int_{\{h: \|h\| \leq M\}} \exp \left[-\frac{t}{2} (h - \Delta_{n,\theta_0})^\top I(\theta_0) (h - \Delta_{n,\theta_0}) \right] dh, \end{aligned}$$

where the first equality follows from the continuity of $\pi(\theta)$ at θ_0 and the coordinate transformation $h = \sqrt{n}(\theta - \theta_0)$; and the second equality follows from the uniform expansion given by Assumption 1. This equality holds for every $M > 0$ and hence also for some $M_n \rightarrow \infty$. Since Δ_{n,θ_0} is bounded in probability, we have

$$\begin{aligned} & \int_{\{h: \|h\| \leq M_n\}} \exp \left[-\frac{t}{2} (h - \Delta_{n,\theta_0})^\top I(\theta_0) (h - \Delta_{n,\theta_0}) \right] dh \\ &= (1 + o_{P_{\theta_0}^n}(1)) \int_{\mathbb{R}^p} \exp \left[-\frac{t}{2} (h - \Delta_{n,\theta_0})^\top I(\theta_0) (h - \Delta_{n,\theta_0}) \right] dh \\ &= (1 + o_{P_{\theta_0}^n}(1)) \left(\frac{2\pi}{t} \right)^{p/2} |I(\theta_0)|^{-1/2}. \end{aligned}$$

Thus,

$$\begin{aligned} \int_{\Theta} \exp[-tR_n(\theta_0\|\theta)] \pi(\theta) d\theta &= (1 + o_{P_{\theta_0}^n}(1)) \int_{\{\theta: \|\theta - \theta_0\| \leq M/\sqrt{n}\}} \exp[-tR_n(\theta_0\|\theta)] \pi(\theta) d\theta \\ &= (1 + o_{P_{\theta_0}^n}(1)) \pi(\theta_0) \left(\frac{2\pi}{tn} \right)^{p/2} |I(\theta_0)|^{-1/2} \exp \left\{ \frac{t}{2} \Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} \right\}. \end{aligned}$$

This completes the proof. \square

Proof of Proposition 1. From Proposition 6, we have

$$\int_{\Theta} \exp \{-R_n(\theta_0\|\theta)\} \pi(\theta) d\theta = (1 + o_{P_{\theta_0}^n}(1)) \pi(\theta_0) \left(\frac{2\pi}{n} \right)^{p/2} |I(\theta_0)|^{-1/2} \exp \left\{ \frac{1}{2} \Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} \right\},$$

and

$$\int_{\tilde{\Theta}} \exp \{-R_n(\nu_0\|\nu)\} \pi_0(\nu) d\nu = (1 + o_{P_{\theta_0}^n}(1)) \pi_0(\nu_0) \left(\frac{2\pi}{n} \right)^{p_0/2} |I_{\theta_0}^{(0)}|^{-1/2} \exp \left\{ \frac{1}{2} \Delta_{n,\theta_0}^{(0)\top} I_{\nu}(\theta_0) \Delta_{n,\theta_0}^{(0)} \right\}.$$

It follows that

$$\begin{aligned} \log \text{BF}_1(\mathbf{X}^n) &= \log \int_{\Theta} \exp[-R_n(\theta_0\|\theta)] \pi(\theta) d\theta - \log \int_{\tilde{\Theta}} \exp[-R_n(\nu_0\|\nu)] \pi_0(\nu) d\nu \\ &= \frac{p-p_0}{2} \log \left(\frac{2\pi}{n} \right) + \frac{1}{2} \left(\Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} - \Delta_{n,\theta_0}^{(0)\top} I_{\nu}(\theta_0) \Delta_{n,\theta_0}^{(0)} \right) \\ &\quad + \log \frac{|I(\theta_0)|^{-\frac{1}{2}} \pi(\theta_0)}{|I_{\nu}(\theta_0)|^{-\frac{1}{2}} \pi_0(\nu_0)} + o_{P_{\theta_0}^n}(1) \\ &= \frac{p-p_0}{2} \log \left(\frac{2\pi}{n} \right) + \frac{1}{2} \left(\Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} - \Delta_{n,\theta_0}^{(0)\top} I_{\nu}(\theta_0) \Delta_{n,\theta_0}^{(0)} \right) \\ &\quad + \log \frac{|I_{\xi|\nu}(\theta_0)|^{-\frac{1}{2}} \pi(\theta_0)}{\pi_0(\nu_0)} + o_{P_{\theta_0}^n}(1). \end{aligned}$$

By the mutual contiguity of $P_{\theta_0}^n$ and $P_{\theta_n}^n$, the term $o_{P_{\theta_0}^n}(1)$ is also $o_{P_{\theta_n}^n}(1)$. \square

Proposition 7. Suppose that Assumptions 1, 2, 3 hold. Then the following assertions hold.

(a) If $t \rightarrow 0$, $tn \rightarrow \infty$, then

$$\int_{\Theta} \exp\{-tR_n(\theta_0\|\theta)\} \pi(\theta) d\theta = (1 + o_{P_{\theta_0}^n}(1)) \pi(\theta_0) \left(\frac{2\pi}{tn}\right)^{p/2} |I(\theta_0)|^{-1/2}.$$

(b) If $tn \rightarrow c \in (0, +\infty)$, then

$$\int_{\Theta} \exp\{-tR_n(\theta_0\|\theta)\} \pi(\theta) d\theta \xrightarrow{P_{\theta_0}^n} \int_{\Theta} \exp\{-cD_1(\theta_0\|\theta)\} \pi(\theta) d\theta.$$

Proof. Assertion (a) follows from the following Lemma 4 and Lemma 3 with $t^\dagger = 0$. Assertion (b) follows directly from Lemma 4. \square

Lemma 3. Suppose that Assumptions 1, 2, 3 hold. Suppose as $n \rightarrow \infty$, $t \rightarrow 0$, $tn \rightarrow \infty$. Then for any fixed $t^\dagger \in [0, t^*)$,

$$\int_{\Theta} \exp\{-tnD_{1-t^\dagger}(\theta_0\|\theta)\} \pi(\theta) d\theta = (1 + o(1)) \pi(\theta_0) \left(\frac{2\pi}{(1-t^\dagger)tn}\right)^{p/2} |I(\theta_0)|^{-1/2}.$$

Proof. Assumption 2 implies that

$$\begin{aligned} & \int_{\Theta} \exp\{-tnD_{1-t^\dagger}(\theta_0\|\theta)\} \pi(\theta) d\theta \\ & \geq \int_{\{\theta: \|\theta - \theta_0\| \leq (tn)^{-1/4}\}} \exp\{-tnD_{1-t^\dagger}(\theta_0\|\theta)\} \pi(\theta) d\theta \\ & = (1 + o(1)) \pi(\theta_0) \int_{\{\theta: \|\theta - \theta_0\| \leq (tn)^{-1/4}\}} \exp\left\{-\frac{tn(1-t^\dagger)}{2} (\theta - \theta_0)^\top I(\theta_0) (\theta - \theta_0)\right\} d\theta \\ & = (1 + o(1)) \pi(\theta_0) (tn)^{-p/2} \int_{\{\vartheta: \|\vartheta\| \leq (tn)^{1/4}\}} \exp\left\{-\frac{1-t^\dagger}{2} \vartheta^\top I(\theta_0) \vartheta\right\} d\vartheta \\ & = (1 + o(1)) \pi(\theta_0) \left(\frac{2\pi}{(1-t^\dagger)tn}\right)^{p/2} |I(\theta_0)|^{-1/2}. \end{aligned}$$

Now we prove the other direction of the inequality. Assumption 2 allows us to choose $\epsilon \in (0, 1)$ and $\delta > 0$ such that for $\|\theta - \theta_0\| \leq \delta$,

$$D_{1-t^\dagger}(\theta_0\|\theta) \geq (1 - \epsilon) \frac{1-t^\dagger}{2} (\theta - \theta_0)^\top I(\theta_0) (\theta - \theta_0), \quad \pi(\theta) \leq (1 + \epsilon) \pi(\theta_0).$$

Also by Assumption 2, there exists a $\epsilon^* > 0$ such that $D_{t^\dagger}(\theta_0\|\theta) \geq \epsilon^*$ for $\|\theta - \theta_0\| \geq \delta$. Hence for

sufficiently large n such that $tn > c^*$, we have

$$\begin{aligned}
& \int_{\Theta} \exp\{-tnD_{1-t^\dagger}(\theta_0\|\theta)\}\pi(\theta) d\theta \\
& \leq (1+\epsilon)\pi(\theta_0) \int_{\{\theta:\|\theta-\theta_0\|\leq\delta\}} \exp\left\{-tn(1-\epsilon)\frac{1-t^\dagger}{2}(\theta-\theta_0)^\top I(\theta_0)(\theta-\theta_0)\right\} d\theta \\
& \quad + \exp\{-tn\epsilon^*\} \int_{\{\theta:\|\theta-\theta_0\|>\delta\}} \exp\{-tn(D_{1-t^\dagger}(\theta_0\|\theta)-\epsilon^*)\}\pi(\theta) d\theta \\
& \leq (1+\epsilon)\pi(\theta_0) \int_{\Theta} \exp\left\{-tn(1-\epsilon)\frac{1-t^\dagger}{2}(\theta-\theta_0)^\top I(\theta_0)(\theta-\theta_0)\right\} d\theta \\
& \quad + \exp\{-tn\epsilon^*\} \int_{\Theta} \exp\{-c^*(D_{1-t^*}(\theta_0\|\theta)-\epsilon^*)\}\pi(\theta) d\theta \\
& = (1+\epsilon)(1-\epsilon)^{-p/2}\pi(\theta_0) \left(\frac{2\pi}{(1-t^\dagger)tn}\right)^{p/2} |I(\theta_0)|^{-1/2} \\
& \quad + \exp\{-(tn-c^*)\epsilon^*\} \int_{\Theta} \exp\{-c^*D_{1-t^*}(\theta_0\|\theta)\}\pi(\theta) d\theta \\
& = (1+o(1))(1+\epsilon)(1-\epsilon)^{-p/2}\pi(\theta_0) \left(\frac{2\pi}{(1-t^\dagger)tn}\right)^{p/2} |I(\theta_0)|^{-1/2}.
\end{aligned}$$

Note that ϵ can be arbitrarily small. This completes the proof. \square

Lemma 4. *Suppose that Assumptions 1, 2, 3 hold. Suppose as $n \rightarrow \infty$, $t \rightarrow 0$, $tn \rightarrow c \in (c^*, +\infty]$. Then*

$$\int_{\Theta} \exp\{-tR_n(\theta_0\|\theta)\}\pi(\theta) d\theta = (1 + o_{P_{\theta_0}^n}(1)) \int_{\Theta} \exp\{-tnD_1(\theta_0\|\theta)\}\pi(\theta) d\theta.$$

Proof. Without loss of generality, suppose $tn > c^*$. Define

$$w_n(\theta) = \frac{\exp\{-tnD_1(\theta_0\|\theta)\}\pi(\theta)}{\int_{\Theta} \exp\{-tnD_1(\theta_0\|\theta)\}\pi(\theta) d\theta}.$$

Note that

$$\int_{\Theta} \exp\{-tnD_1(\theta_0\|\theta)\}\pi(\theta) d\theta \leq \int_{\Theta} \exp\{-c^*D_{1-t^*}(\theta_0\|\theta)\}\pi(\theta) d\theta < \infty.$$

Hence $w_n(\theta)$ is well defined. It is easy to verify the following equality which will play an important role in our proof.

$$\begin{aligned}
D_1(w_n(\theta) d\theta \|\pi_t(\theta|\mathbf{X}^n) d\theta) &= \int_{\Theta} [tR_n(\theta_0\|\theta) - tnD_1(\theta_0\|\theta)] w_n(\theta) d\theta \\
&+ \log \frac{\int_{\Theta} \exp\{-tR_n(\theta_0\|\theta)\}\pi(\theta) d\theta}{\int_{\Theta} \exp\{-tnD_1(\theta_0\|\theta)\}\pi(\theta) d\theta}.
\end{aligned} \tag{18}$$

In view of (18), we only need to prove

$$P_{\theta_0}^n D_1 (w_n(\theta) d\theta \| \pi_t(\theta | \mathbf{X}^n) d\theta) \rightarrow 0 \quad (19)$$

and

$$P_{\theta_0}^n \left(\int_{\Theta} [tR_n(\theta_0 \| \theta) - tnD_1(\theta_0 \| \theta)] w_n(\theta) d\theta \right)^2 \rightarrow 0. \quad (20)$$

Proof of (19): From Fubini's theorem and the fact $P_{\theta_0}^n R_n(\theta_0 \| \theta) = nD_1(\theta_0 \| \theta)$, we have

$$P_{\theta_0}^n \int_{\Theta} [tR_n(\theta_0 \| \theta) - tnD_1(\theta_0 \| \theta)] w_n(\theta) d\theta = 0.$$

Jensen's inequality implies that

$$\begin{aligned} P_{\theta_0}^n \log \int_{\Theta} \exp\{-tR_n(\theta_0 \| \theta)\} \pi(\theta) d\theta &\leq \log P_{\theta_0}^n \int_{\Theta} \exp\{-tR_n(\theta_0 \| \theta)\} \pi(\theta) d\theta \\ &= \log \int_{\Theta} \exp\{-tnD_{1-t}(\theta_0 \| \theta)\} \pi(\theta) d\theta. \end{aligned}$$

Then from (18), we have the upper bound

$$P_{\theta_0}^n D_1 (w_n(\theta) d\theta \| \pi_t(\theta | \mathbf{X}^n) d\theta) \leq \log \frac{\int_{\Theta} \exp\{-tnD_{1-t}(\theta_0 \| \theta)\} \pi(\theta) d\theta}{\int_{\Theta} \exp\{-tnD_1(\theta_0 \| \theta)\} \pi(\theta) d\theta}. \quad (21)$$

If $tn \rightarrow c \in (0, +\infty)$, then the dominated convergence theorem implies that

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\Theta} \exp\{-tnD_{1-t}(\theta_0 \| \theta)\} \pi(\theta) d\theta &= \int_{\Theta} \exp\{-cD_1(\theta_0 \| \theta)\} \pi(\theta) d\theta, \\ \lim_{n \rightarrow \infty} \int_{\Theta} \exp\{-tnD_1(\theta_0 \| \theta)\} \pi(\theta) d\theta &= \int_{\Theta} \exp\{-cD_1(\theta_0 \| \theta)\} \pi(\theta) d\theta. \end{aligned}$$

Hence the right hand side of (21) converges to 0.

We turn to the case $tn \rightarrow \infty$. For any $t^\dagger \in (0, t^*)$, since $t < t^\dagger$ for sufficiently large n , we have

$$\begin{aligned} \limsup_{n \rightarrow \infty} \log \frac{\int_{\Theta} \exp\{-tnD_{1-t}(\theta_0 \| \theta)\} \pi(\theta) d\theta}{\int_{\Theta} \exp\{-tnD_1(\theta_0 \| \theta)\} \pi(\theta) d\theta} &\leq \limsup_{n \rightarrow \infty} \log \frac{\int_{\Theta} \exp\{-tnD_{1-t^\dagger}(\theta_0 \| \theta)\} \pi(\theta) d\theta}{\int_{\Theta} \exp\{-tnD_1(\theta_0 \| \theta)\} \pi(\theta) d\theta} \\ &= -\frac{p}{2} \log(1 - t^\dagger), \end{aligned}$$

where the last equality follows from Lemma 3. Let $t^\dagger \rightarrow 0$, then the right hand side of (21) converges to 0. This completes the proof of (19).

Proof of (20): It can be seen that

$$\begin{aligned} &P_{\theta_0}^n \left(\int_{\Theta} [tR_n(\theta_0 \| \theta) - tnD_1(\theta_0 \| \theta)] w_n(\theta) d\theta \right)^2 \\ &\leq \int_{\Theta} P_{\theta_0}^n [tR_n(\theta_0 \| \theta) - tnD_1(\theta_0 \| \theta)]^2 w_n(\theta) d\theta \\ &= t^2 n \frac{\int_{\Theta} V(\theta_0 \| \theta) \exp\{-tnD_1(\theta_0 \| \theta)\} \pi(\theta) d\theta}{\int_{\Theta} \exp\{-tnD_1(\theta_0 \| \theta)\} \pi(\theta) d\theta}. \end{aligned}$$

If $tn \rightarrow c \in (0, +\infty)$, the above expression obviously tends to 0. Now we assume $tn \rightarrow \infty$. Assumption 2 allows us to choose $\epsilon \in (0, 1)$ and $\delta > 0$ such that for $\|\theta - \theta_0\| \leq \delta$,

$$D_1(\theta_0\|\theta) \geq \frac{1-\epsilon}{2}(\theta - \theta_0)^\top I(\theta_0)(\theta - \theta_0), \quad V(\theta_0\|\theta) \leq C\|\theta - \theta_0\|^2, \quad \pi(\theta) \leq (1+\epsilon)\pi(\theta_0).$$

Also by Assumption 2, there exists a $\epsilon^* > 0$ such that $D_1(\theta_0\|\theta) \geq \epsilon^*$ for $\|\theta - \theta_0\| \geq \delta$. Then for sufficiently large n such that $tn > c^\dagger$, we have

$$\begin{aligned} & \int_{\Theta} V(\theta_0\|\theta) \exp\{-tnD_1(\theta_0\|\theta)\} \pi(\theta) d\theta \\ & \leq (1+\epsilon)\pi(\theta_0) \int_{\{\theta: \|\theta-\theta_0\| \leq \delta\}} C\|\theta - \theta_0\|^2 \exp\left\{-tn(1-\epsilon)\frac{1}{2}(\theta - \theta_0)^\top I(\theta_0)(\theta - \theta_0)\right\} d\theta \\ & \quad + \exp\{-tn\epsilon^*\} \int_{\{\theta: \|\theta-\theta_0\| > \delta\}} V(\theta_0\|\theta) \exp\{-tn(D_1(\theta_0\|\theta) - \epsilon^*)\} \pi(\theta) d\theta. \\ & \leq (1+\epsilon)\pi(\theta_0)(tn)^{-p/2-1} \int_{\mathbb{R}^p} C\|\vartheta\|^2 \exp\left\{-(1-\epsilon)\frac{1}{2}\vartheta^\top I(\theta_0)\vartheta\right\} d\vartheta \\ & \quad + \exp\left\{-(tn - c^\dagger)\epsilon^*\right\} \int_{\Theta} V(\theta_0\|\theta) \exp\left\{-c^\dagger D_1(\theta_0\|\theta)\right\} \pi(\theta) d\theta \\ & = O\left((tn)^{-p/2-1}\right). \end{aligned}$$

The last inequality, combined with Lemma 3, leads to

$$P_{\theta_0}^n \left(\int_{\Theta} [tR_n(\theta_0\|\theta) - tnD_1(\theta_0\|\theta)] w_n(\theta) d\theta \right)^2 = t^2 n \frac{O((tn)^{-p/2-1})}{\pi(\theta_0) (2\pi)^{p/2} (tn)^{-p/2} |I(\theta_0)|^{-1/2}} \rightarrow 0.$$

Hence (20) holds. This completes the proof. □

Proof of Theorem 1.

$$\begin{aligned} \log \Lambda_{a,b} &= \log \int_{\Theta} \exp\{-aR_n(\theta_0\|\theta)\} \pi(\theta) d\theta - \log \int_{\Theta} \exp\{-bR_n(\theta_0\|\theta)\} \pi(\theta) d\theta \\ & \quad - \log \int_{\tilde{\Theta}_0} \exp\{-aR_n(\theta_0\|\nu, \xi_0)\} \pi_0(\nu) d\nu + \log \int_{\tilde{\Theta}_0} \exp\{-bR_n(\theta_0\|\nu, \xi_0)\} \pi_0(\nu) d\nu. \end{aligned}$$

If a and b are fixed, Proposition 6 implies that

$$\log \Lambda_{a,b} = -\frac{p-p_0}{2} \log\left(\frac{a}{b}\right) + \frac{a-b}{2} \left(\Delta_{n,\theta_0}^\top I(\theta_0) \Delta_{n,\theta_0} - \Delta_{n,\theta_0}^{(0)T} I_\nu(\theta_0) \Delta_{n,\theta_0}^{(0)} \right) + o_{P_{\theta_0}^n}(1).$$

Then the assertion (a) follows from Lemma 2.

Similarly, assertion (b) and (b) follows from Proposition 7 and Lemma 2. □

Appendix C Proofs in Section 4

Proof of Proposition 2. For exponential family, we have

$$I(\theta_0)\Delta_{n,\theta_0} = n^{-1/2} \sum_{i=1}^n T(X_i) - \sqrt{n} \frac{\partial}{\partial \theta} A(\theta_0)$$

and

$$R_n(\theta_0 \| \theta_0 + n^{-1/2}h) = -h^\top I(\theta_0)\Delta_{n,\theta_0} + \frac{1}{2}h^\top I(\theta_0)h + g_n(h),$$

where

$$g_n(h) = n \left(A(\theta_0 + n^{-1/2}h) - A(\theta_0) - n^{-1/2}h \frac{\partial}{\partial \theta} A(\theta_0) - \frac{1}{2}h^\top I(\theta_0)h \right).$$

Without loss of generality, we assume $M_n \rightarrow \infty$ and $M_n^3/\sqrt{n} \rightarrow 0$. From Taylor's theorem and the continuity of the third derivative of $A(\theta)$,

$$\max_{\{h: \|h\| \leq M_n\}} |g_n(h)| = O\left(\frac{M_n^3}{\sqrt{n}}\right) \rightarrow 0.$$

This allows us to derive the following lower bound for $\int_{\Theta} \exp\{-tR_n(\theta_0 \| \theta)\} \pi(\theta) d\theta$.

$$\begin{aligned} & \int_{\Theta} \exp\{-tR_n(\theta_0 \| \theta)\} \pi(\theta) d\theta \\ & \geq \int_{\{\theta: \|\theta - \theta_0\| \leq M_n/\sqrt{n}\}} \exp\{-tR_n(\theta_0 \| \theta)\} \pi(\theta) d\theta \\ & = (1 + o_{P_{\theta_0}^n}(1)) \pi(\theta_0) n^{-p/2} \int_{\{h: \|h\| \leq M_n\}} \exp\left\{th^\top I(\theta_0)\Delta_{n,\theta_0} - \frac{t}{2}h^\top I(\theta_0)h\right\} dh \\ & = (1 + o_{P_{\theta_0}^n}(1)) \pi(\theta_0) n^{-p/2} \int_{\mathbb{R}^p} \exp\left\{th^\top I(\theta_0)\Delta_{n,\theta_0} - \frac{t}{2}h^\top I(\theta_0)h\right\} dh \\ & = (1 + o_{P_{\theta_0}^n}(1)) \pi(\theta_0) \left(\frac{2\pi}{tn}\right)^{p/2} |I(\theta_0)|^{-1/2} \exp\left\{\frac{t}{2}\Delta_{n,\theta_0}^\top I(\theta_0)\Delta_{n,\theta_0}\right\}. \end{aligned} \tag{22}$$

Next we lower bound $R_n(\theta_0 \| \theta)$ for $\|\theta - \theta_0\| \geq M_n/\sqrt{n}$. We have

$$\begin{aligned} \min_{\{\theta: \|\theta - \theta_0\| = M_n/\sqrt{n}\}} R_n(\theta_0 \| \theta) &= \min_{\{h: \|h\| = M_n\}} R_n(\theta_0 \| \theta_0 + n^{-1/2}h) \\ &\geq -\|I(\theta_0)\Delta_{n,\theta_0}\| M_n + \frac{\lambda_{\min}(I(\theta_0))}{2} M_n^2 - \max_{\{h: \|h\| = M_n\}} |g_n(h)|, \end{aligned}$$

where $\lambda_{\min}(I(\theta_0)) > 0$ is the minimum eigenvalue of $I(\theta_0)$. Also note that $I(\theta_0)\Delta_{n,\theta_0}$ is bounded in probability. Hence with probability tending to 1,

$$\min_{\{\theta: \|\theta - \theta_0\| = M_n/\sqrt{n}\}} R_n(\theta_0 \| \theta) \geq \frac{\lambda_{\min}(I(\theta_0))}{4} M_n^2.$$

Note that $R_n(\theta_0 \| \theta)$ is convex in θ and $R_n(\theta_0 \| \theta_0) = 0$. Then Jensen's inequality implies that for $\|\theta - \theta_0\| \geq M_n/\sqrt{n}$,

$$\frac{M_n/\sqrt{n}}{\|\theta - \theta_0\|} R_n(\theta_0 \| \theta) \geq R_n\left(\theta_0 \left\| \theta_0 + \frac{M_n/\sqrt{n}}{\|\theta - \theta_0\|} (\theta - \theta_0) \right\| \right).$$

The last two inequalities imply that with probability tending to 1,

$$\begin{aligned}
R_n(\theta_0|\theta) &\geq \frac{\sqrt{n}\|\theta - \theta_0\|}{M_n} R_n\left(\theta_0\left\|\theta_0 + \frac{M_n/\sqrt{n}}{\|\theta - \theta_0\|}(\theta - \theta_0)\right\|\right) \\
&\geq \frac{\sqrt{n}\|\theta - \theta_0\|}{M_n} \min_{\{\theta:\|\theta - \theta_0\|=M_n/\sqrt{n}\}} R_n(\theta_0|\theta) \\
&\geq \frac{\lambda_{\min}(I(\theta_0))}{4} \sqrt{n}\|\theta - \theta_0\| M_n.
\end{aligned}$$

Fix an $\epsilon > 0$ such that $\sup_{\|\theta - \theta_0\| < \epsilon} \pi(\theta) < +\infty$. For sufficiently large n , with probability tending to 1, we have

$$\begin{aligned}
&\int_{\{\theta:\|\theta - \theta_0\| > M_n/\sqrt{n}\}} \exp\{-tR_n(\theta_0|\theta)\} \pi(\theta) d\theta \\
&\leq \int_{\{\theta:\|\theta - \theta_0\| > M_n/\sqrt{n}\}} \exp\left\{-\frac{t\lambda_{\min}(I(\theta_0))}{4} \sqrt{n}\|\theta - \theta_0\| M_n\right\} \pi(\theta) d\theta \\
&= \int_{\{\theta:M_n/\sqrt{n} < \|\theta - \theta_0\| \leq \epsilon\}} \exp\left\{-\frac{t\lambda_{\min}(I(\theta_0))}{4} \sqrt{n}\|\theta - \theta_0\| M_n\right\} \pi(\theta) d\theta \\
&\quad + \exp\left\{-\frac{t\lambda_{\min}(I(\theta_0))}{4} \epsilon \sqrt{n} M_n\right\} \int_{\{\theta:\|\theta - \theta_0\| > \epsilon\}} \exp\left\{-\frac{t\lambda_{\min}(I(\theta_0))}{4} \sqrt{n}(\|\theta - \theta_0\| - \epsilon) M_n\right\} \pi(\theta) d\theta \\
&\leq \left(\sup_{\|\theta - \theta_0\| < \epsilon} \pi(\theta)\right) \int_{\{\theta:\|\theta - \theta_0\| \geq M_n/\sqrt{n}\}} \exp\left\{-\frac{t\lambda_{\min}(I(\theta_0))}{4} \sqrt{n}\|\theta - \theta_0\| M_n\right\} d\theta \\
&\quad + \exp\left\{-\frac{t\lambda_{\min}(I(\theta_0))}{4} \epsilon \sqrt{n} M_n\right\} \int_{\{\theta:\|\theta - \theta_0\| > \epsilon\}} \exp\{-c^*(\|\theta - \theta_0\| - \epsilon)\} \pi(\theta) d\theta \\
&= \left(\sup_{\|\theta - \theta_0\| < \epsilon} \pi(\theta)\right) (\sqrt{n} M_n)^{-p} \int_{\{h:\|h\| \geq M_n^2\}} \exp\left\{-\frac{t\lambda_{\min}(I(\theta_0))}{4} \|h\|\right\} dh \\
&\quad + \exp\left\{-\frac{t\lambda_{\min}(I(\theta_0))}{4} \epsilon \sqrt{n} M_n\right\} \int_{\{\theta:\|\theta - \theta_0\| > \epsilon\}} \exp\{-c^*(\|\theta - \theta_0\| - \epsilon)\} \pi(\theta) d\theta.
\end{aligned}$$

It follows that

$$\int_{\{\theta:\|\theta - \theta_0\| > M_n/\sqrt{n}\}} \exp\{-tR_n(\theta_0|\theta)\} \pi(\theta) d\theta = O_{P_{\theta_0}^n}((\sqrt{n} M_n)^{-p}).$$

Combining the last display and (22) leads to

$$\frac{\int_{\{\theta:\|\theta - \theta_0\| > M_n/\sqrt{n}\}} \exp\{-tR_n(\theta_0|\theta)\} \pi(\theta) d\theta}{\int_{\Theta} \exp\{-tR_n(\theta_0|\theta)\} \pi(\theta) d\theta} = O_{P_{\theta_0}^n}(M_n^{-p}) \xrightarrow{P_{\theta_0}^n} 0.$$

This completes the proof. □

Proof of Proposition 3. Note that $L_1(\Theta; \mathbf{X}^n)$ is well defined $P_{\theta_0}^n$ -a.s. since it has finite integral

$$\int_{\mathcal{X}^n} L_1(\Theta; \mathbf{X}^n) d\mu^n = \int_{\Theta} \left(\int_{\mathcal{X}^n} p_n(\mathbf{X}^n|\theta) d\mu^n \right) \pi(\theta) d\theta = 1.$$

For $0 < t < 1$, by Hölder's inequality, we have $L_t(\Theta; \mathbf{X}^n) \leq L_1^{1/t}(\Theta; \mathbf{X}^n)$. This proves the first part of the proposition.

We turn to the second part of the proposition. The following observation is critical in our proof. For $\epsilon \in (0, +\infty]$, $\gamma \in (0, +\infty)$,

$$\begin{aligned} \int_{\{x \in \mathbb{R}^p : \|x\| < \epsilon\}} \|x\|^{\gamma-p} \exp\{-\|x\|\} \, dx &= \frac{\pi^{p/2}}{\Gamma(p/2)} \int_0^{\epsilon^2} y^{\gamma/2-1} \exp\{-\sqrt{y}\} \, dy \\ &= \frac{2\pi^{p/2}}{\Gamma(p/2)} \int_0^\epsilon y^{\gamma-1} \exp\{-y\} \, dy, \end{aligned} \quad (23)$$

where the first equality follows from Fang et al. (1990), Lemma 1.4. If $\gamma \leq 0$, the above integral is infinity.

In view of (23), we can define a family of density functions indexed by $\theta \in \mathbb{R}^p$ as

$$p(X|\theta) = \frac{\Gamma(p/2)}{2\pi^{p/2}\Gamma(\gamma)} \|X - \theta\|^{\gamma-p} \exp\{-\|X - \theta\|\},$$

where $\gamma > 0$ is a known hyperparameter. Suppose $\Theta = \mathbb{R}^p$, $X_1, \dots, X_n \in \mathbb{R}^p$ are iid random vectors with density function $p(X|\theta_0)$. Let $\pi(\theta)$ be any proper prior density which is continuous and positive. Then

$$L_t(\Theta; \mathbf{X}^n) = \left(\frac{\Gamma(p/2)}{2\pi^{p/2}\Gamma(\gamma)} \right)^{tn} \int_{-\infty}^{+\infty} \left(\prod_{i=1}^n \|X_i - \theta\| \right)^{t(\gamma-p)} \exp\left\{-t \sum_{i=1}^n \|X_i - \theta\|\right\} \pi(\theta) \, d\theta.$$

Note that with probability 1, there is no tie among X_1, \dots, X_n . Consequently, there exists a sufficiently small $\epsilon > 0$ such that the sets $A_i := \{\theta : \|X_i - \theta\| < \epsilon\}$, $i = 1, \dots, n$, are disjoint. It can be seen that for $i = 1, \dots, n$, $p(X_i|\theta)$ is continuous and bounded for $\theta \notin A_i$. Hence

$$\int_{\mathbb{R}^p \setminus \bigcup_{i=1}^n A_i} [p_n(\mathbf{X}^n|\theta)]^t \pi(\theta) \, d\theta \leq \prod_{i=1}^n \sup_{\theta \notin A_i} [p(X_i|\theta)]^t < \infty.$$

On the other hand, for $i = 1, \dots, n$,

$$\begin{aligned} \int_{A_i} [p_n(\mathbf{X}^n|\theta)]^t \pi(\theta) \, d\theta &\leq \left(\prod_{j \neq i} \sup_{\theta \in A_i} [p(X_j|\theta)]^t \right) \left(\sup_{\theta \in A_i} \pi(\theta) \right) \int_{A_i} [p(X_i|\theta)]^t \, d\theta, \\ \int_{A_i} [p_n(\mathbf{X}^n|\theta)]^t \pi(\theta) \, d\theta &\geq \left(\prod_{j \neq i} \inf_{\theta \in A_i} [p(X_j|\theta)]^t \right) \left(\inf_{\theta \in A_i} \pi(\theta) \right) \int_{A_i} [p(X_i|\theta)]^t \, d\theta. \end{aligned}$$

Hence $\int_{A_i} [p_n(\mathbf{X}^n|\theta)]^t \pi(\theta) \, d\theta$ is finite if and only if $\int_{A_i} [p(X_i|\theta)]^t \, d\theta$ is finite. Note that

$$\int_{A_i} [p(X_i|\theta)]^t \, d\theta = \left(\frac{\Gamma(p/2)}{2\pi^{p/2}\Gamma(\gamma)} \right)^t \int_{\{\|\theta\| < \epsilon\}} \|\theta\|^{t(\gamma-p)+p-p} \exp\{-t\|\theta\|\} \, d\theta.$$

It follows that $\int_{A_i} [p(X_i|\theta)]^t \, d\theta$ is finite if and only if $t(\gamma-p)+p > 0$, or equivalently, $\gamma > p(t-1)/t$. Thus, $L_t(\theta; \mathbf{X}^n)$ is finite if and only if $\gamma > p(t-1)/t$. If $t > 1$, $p(t-1)/t > 0$, then $L_t(\theta; \mathbf{X}^n) = +\infty$ provided $\gamma \in (0, p(t-1)/t]$. This completes the proof. \square

Proof of Proposition 4. Without loss of generality, we assume $M_n/\sqrt{n} \rightarrow 0$.

It follows from Fubini's theorem that

$$\begin{aligned} & P_{\theta_0}^n \int_{\{\theta: \|\theta - \theta_0\| \geq M_n/\sqrt{n}\}} \exp\{-tR_n(\theta_0|\theta)\} \pi(\theta) d\theta \\ &= \int_{\{\theta: \|\theta - \theta_0\| \geq M_n/\sqrt{n}\}} \exp\{-tnD_{1-t}(\theta_0|\theta)\} \pi(\theta) d\theta. \end{aligned}$$

Assumption 2 implies that there exist $\delta > 0$, $C > 0$ and $\epsilon > 0$ such that $D_{1-t}(\theta_0|\theta) \geq C\|\theta - \theta_0\|^2$ for $\|\theta - \theta_0\| \leq \delta$ and $D_{1-t}(\theta_0|\theta) \geq \epsilon$ for $\|\theta - \theta_0\| > \delta$. Decompose the integral region into two parts $\{\theta : M_n/\sqrt{n} \leq \|\theta - \theta_0\| \leq \delta\}$ and $\{\theta : \|\theta - \theta_0\| > \delta\}$. Then for sufficiently large n ,

$$\begin{aligned} & \int_{\{\theta: \|\theta - \theta_0\| \geq M_n/\sqrt{n}\}} \exp\{-tnD_{1-t}(\theta_0|\theta)\} \pi(\theta) d\theta \\ & \leq \max_{\|\theta - \theta_0\| \leq \delta} \pi(\theta) \int_{\{\theta: \|\theta - \theta_0\| \geq M_n/\sqrt{n}\}} \exp\{-tCn\|\theta - \theta_0\|^2\} d\theta \\ & \quad + \exp\{-t\epsilon n\} \int_{\{\theta: \|\theta - \theta_0\| \geq M_n/\sqrt{n}\}} \exp\{-tn(D_{1-t}(\theta_0|\theta) - \epsilon)\} \pi(\theta) d\theta \\ & \leq \left(\max_{\|\theta - \theta_0\| \leq \delta} \pi(\theta) \right) n^{-p/2} \int_{\{h: \|h\| \geq M_n\}} \exp\{-tC\|h\|^2\} dh \\ & \quad + \exp\{-t\epsilon n\} \int_{\{\theta: \|\theta - \theta_0\| \geq M_n/\sqrt{n}\}} \exp\{-c^*(D_{1-t}(\theta_0|\theta) - \epsilon)\} \pi(\theta) d\theta. \end{aligned}$$

Hence

$$n^{p/2} \int_{\{\theta: \|\theta - \theta_0\| \geq M_n/\sqrt{n}\}} \left[\frac{p_n(\mathbf{X}^n|\theta)}{p_n(\mathbf{X}^n|\theta_0)} \right]^t \pi(\theta) d\theta = o_{P_{\theta_0}^n}(1). \quad (24)$$

Now we consider the denominator of (??).

$$\begin{aligned} & \int_{\Theta} \left[\frac{p_n(\mathbf{X}^n|\theta)}{p_n(\mathbf{X}^n|\theta_0)} \right]^t \pi(\theta) d\theta \geq \int_{\{\theta: \|\theta - \theta_0\| \leq n^{-1/2}\}} \left[\frac{p_n(\mathbf{X}^n|\theta)}{p_n(\mathbf{X}^n|\theta_0)} \right]^t \pi(\theta) d\theta \\ & \geq \left(\min_{\|\theta - \theta_0\| \leq n^{-1/2}} \left[\frac{p_n(\mathbf{X}^n|\theta)}{p_n(\mathbf{X}^n|\theta_0)} \right]^t \pi(\theta) \right) \int_{\{\theta: \|\theta - \theta_0\| \leq n^{-1/2}\}} 1 d\theta \\ & \geq \left(\exp \left[t \min_{\|\theta - \theta_0\| \leq n^{-1/2}} \log \frac{p_n(\mathbf{X}^n|\theta)}{p_n(\mathbf{X}^n|\theta_0)} \right] \right) \left(\min_{\|\theta - \theta_0\| \leq n^{-1/2}} \pi(\theta) \right) n^{-p/2} \frac{2\pi^{p/2}}{\Gamma(p/2)}. \end{aligned}$$

By Assumption 1,

$$\min_{\|\theta - \theta_0\| \leq n^{-1/2}} \log \frac{p_n(\mathbf{X}^n|\theta)}{p_n(\mathbf{X}^n|\theta_0)} \geq -\|I_{\theta_0}\Delta_{n,\theta_0}\| - \frac{1}{2}\|I_{\theta_0}\| + o_{P_{\theta_0}^n}(1),$$

where $I_{\theta_0}\Delta_{n,\theta_0}$ is bounded in probability. Also note that

$$\min_{\|\theta - \theta_0\| \leq n^{-1/2}} \pi(\theta) \rightarrow \pi(\theta_0) > 0.$$

Then for every $\epsilon' > 0$, there is a constant $c > 0$ such that for large n , with probability at least $1 - \epsilon'$,

$$\int_{\Theta} \left[\frac{p_n(\mathbf{X}^n|\theta)}{p_n(\mathbf{X}^n|\theta_0)} \right]^t \pi(\theta) d\theta \geq cn^{-p/2}.$$

This inequality, together with (24), implies the \sqrt{n} -consistency of $L_t(\cdot; \mathbf{X}^n)$. □

Proof of Theorem 2. It can be seen that

$$\Lambda_{a,b}^* = \frac{\int [p_n(\mathbf{X}^n|\theta_0 + n^{-1/2}h)/p_n(\mathbf{X}^n|\theta_0)]^a \pi_n(h; \mathbf{X}^n) dh}{\int [p_n(\mathbf{X}^n|\nu_0 + n^{-1/2}h^{(0)}, \xi_0)/p_n(\mathbf{X}^n|\theta_0)]^a \pi_n(h^{(0)}; \mathbf{X}^n) dh^{(0)}}.$$

Let $M > 0$ be any fixed number. We have

$$\begin{aligned} & \int_{\|h\| \leq M} \left[\frac{p_n(\mathbf{X}^n|\theta_0 + n^{-1/2}h)}{p_n(\mathbf{X}^n|\theta_0)} \right]^a \pi_n(h; \mathbf{X}^n) dh \\ &= \exp[o_{P_0^n}(1)] \int_{\|h\| \leq M} \exp \left[ah^\top I_{\theta_0} \Delta_{n,\theta_0} - \frac{a}{2} h^\top I_{\theta_0} h \right] \pi_n(h; \mathbf{X}^n) dh \\ &= \int_{\|h\| \leq M} \exp \left[ah^\top I_{\theta_0} \Delta_{n,\theta_0} - \frac{a}{2} h^\top I_{\theta_0} h \right] \phi(h; \Delta_{n,\theta_0}, b^{-1} I_{\theta_0}^{-1}) dh + o_{P_{\theta_0}^n}(1), \end{aligned} \quad (25)$$

where the first equality follows from Assumption 1 and the second equality follows from (9). This is true for every $M > 0$ and hence also for some $M_n \rightarrow \infty$.

Now we prove that for any $M_n \rightarrow +\infty$,

$$\int_{\|h\| > M_n} \left[\frac{p_n(\mathbf{X}^n|\theta_0 + n^{-1/2}h)}{p_n(\mathbf{X}^n|\theta_0)} \right]^a \pi_n(h; \mathbf{X}^n) dh \xrightarrow{P_{\theta_0}^n} 0. \quad (26)$$

By Assumption 4, for any $\epsilon > 0$, with probability at least $1 - \epsilon$,

$$\begin{aligned} & \int_{\|h\| > M_n} \left[\frac{p_n(\mathbf{X}^n|\theta_0 + n^{-1/2}h)}{p_n(\mathbf{X}^n|\theta_0)} \right]^a \pi_n(h; \mathbf{X}^n) dh \\ & \leq \int_{\|h\| > M_n} \left[\frac{p_n(\mathbf{X}^n|\theta_0 + n^{-1/2}h)}{p_n(\mathbf{X}^n|\theta_0)} \right]^a T(h) dh. \end{aligned} \quad (27)$$

By Hölder's inequality,

$$\begin{aligned} & P_{\theta_0}^n \left[\frac{p_n(\mathbf{X}^n|\theta_0 + n^{-1/2}h)}{p_n(\mathbf{X}^n|\theta_0)} \right]^a \\ &= \int p_n(\mathbf{X}^n|\theta_0 + n^{-1/2}h)^a p_n(\mathbf{X}^n|\theta_0)^{1-a} d\mu^n \\ & \leq 1. \end{aligned}$$

Hence the expectation of the right hand side of (27) satisfies

$$P_{\theta_0}^n \int_{\|h\| > M_n} \left[\frac{p_n(\mathbf{X}^n|\theta_0 + n^{-1/2}h)}{p_n(\mathbf{X}^n|\theta_0)} \right]^a T(h) dh \leq \int_{\|h\| > M_n} T(h) dh \rightarrow 0.$$

This verifies (26).

Combining (25) and (26) yields

$$\begin{aligned} & \int \left[\frac{p_n(\mathbf{X}^n | \theta_0 + n^{-1/2}h)}{p_n(\mathbf{X}^n | \theta_0)} \right]^a \pi_n(h; \mathbf{X}^n) dh \\ &= \left(1 + \frac{a}{b}\right)^{-p/2} \exp\left(\frac{a}{2} \Delta_{n,\theta_0}^\top I_{\theta_0} \Delta_{n,\theta_0}\right) + o_{P_{\theta_0}^n}(1). \end{aligned}$$

Similarly, we have

$$\begin{aligned} & \int \left[\frac{p_n(\mathbf{X}^n | \nu_0 + n^{-1/2}h^{(0)}, \xi_0)}{p_n(\mathbf{X}^n | \theta_0)} \right]^a \pi_n(h^{(0)}; \mathbf{X}^n) dh^{(0)} \\ &= \left(1 + \frac{a}{b}\right)^{-p_0/2} \exp\left(\frac{a}{2} \Delta_{n,\theta_0}^{(0)\top} I_{\theta_0}^{(0)} \Delta_{n,\theta_0}^{(0)}\right) + o_{P_{\theta_0}^n}(1). \end{aligned}$$

Hence

$$\begin{aligned} & 2 \log \Lambda_{a,b}^* \\ &= -(p - p_0) \log\left(1 + \frac{a}{b}\right) + a \left(\Delta_{n,\theta_0}^\top I_{\theta_0} \Delta_{n,\theta_0} - \Delta_{n,\theta_0}^{(0)\top} I_{\theta_0}^{(0)} \Delta_{n,\theta_0}^{(0)} \right) + o_{P_{\theta_0}^n}(1). \end{aligned}$$

Then the conclusion follows by the same argument as the last part of Theorem 1. \square

Appendix D Proofs in Section 3

Proof of Proposition 5. We shall verify Assumption 1 and Assumption 2. We use the parameterization $\theta = (\xi, \tau)^\top = (\xi, \sigma^{-2})^\top$. Then

$$p(X|\theta) = \frac{1}{2} \phi(X) + \frac{1}{2} \sqrt{\tau} \phi(\sqrt{\tau}(X - \xi)).$$

By direct calculation, we have

$$\dot{\ell}_{\theta_0}(X) = \left(\frac{1}{2}X, \frac{1}{4}(1 - X^2) \right)^\top.$$

Hence $P_{\theta_0} \dot{\ell}_{\theta_0} = \mathbf{0}_2$ and $I(\theta_0) = \text{diag}(1/4, 1/8)$.

Let $M > 0$ be a fixed constant. For $h = (h_1, h_2)^\top \in \mathbb{R}^2$ such that $\|h\| \leq M$ and $i = 1, \dots, n$, we have

$$\begin{aligned} \frac{p(X_i | \theta_0 + n^{-1/2}h)}{p(X_i | \theta_0)} &= \frac{1}{2} + \frac{1}{2} \sqrt{1 + \frac{h_2}{\sqrt{n}}} \exp \left\{ -\frac{h_2}{2\sqrt{n}} X_i^2 \right. \\ &\quad \left. + \left(1 + \frac{h_2}{\sqrt{n}}\right) \frac{h_1}{\sqrt{n}} X_i - \frac{1}{2} \left(1 + \frac{h_2}{\sqrt{n}}\right) \frac{h_1^2}{n} \right\}. \end{aligned}$$

It is well known that $\max_{1 \leq i \leq n} |X_i| = O_{P_{\theta_0}^n}(\sqrt{\log n})$. Then from Taylor expansion $\exp(x) = 1 + x + x^2/2 + O(x^3)$, we have, uniformly for $\|h\| \leq M$ and $i = 1, \dots, n$, that

$$\begin{aligned} & \exp \left\{ -\frac{h_2}{2\sqrt{n}} X_i^2 + \left(1 + \frac{h_2}{\sqrt{n}}\right) \frac{h_1}{\sqrt{n}} X_i - \frac{1}{2} \left(1 + \frac{h_2}{\sqrt{n}}\right) \frac{h_1^2}{n} \right\} \\ &= 1 - \frac{h_1^2}{2n} + \left(\frac{h_1}{\sqrt{n}} + \frac{h_1 h_2}{n} \right) X_i + \left(-\frac{h_2}{2\sqrt{n}} + \frac{h_1^2}{2n} \right) X_i^2 \\ &\quad - \frac{h_1 h_2}{2n} X_i^3 + \frac{h_2^2}{8n} X_i^4 + O_{P_{\theta_0}^n} \left(\frac{\log^3 n}{n^{3/2}} \right). \end{aligned}$$

On the other hand, for sufficiently large n such that $M/\sqrt{n} \leq 1/2$, we have, uniformly for $\|h\| \leq M$, that

$$\sqrt{1 + \frac{h_2}{\sqrt{n}}} = 1 + \frac{h_2}{2\sqrt{n}} - \frac{h_2^2}{8n} + O\left(\frac{1}{n^3}\right).$$

Multiplying the above two expansions yields

$$\begin{aligned} \frac{p(X_i|\theta_0 + n^{-1/2}h)}{p(X_i|\theta_0)} &= 1 + \frac{h_1}{2\sqrt{n}}X_i + \left(\frac{h_2}{4\sqrt{n}} - \frac{h_1^2}{4n}\right)(1 - X_i^2) + \frac{h_2^2}{16n}X_i^4 \\ &\quad - \frac{h_2^2}{8n}X_i^2 - \frac{h_2^2}{16n} + \frac{3h_1h_2}{4n}X_i - \frac{h_1h_2}{4n}X_i^3 + O_{P_{\theta_0}^n}\left(\frac{\log^3 n}{n^{3/2}}\right). \end{aligned}$$

From Taylor expansion $\log(1+x) = x - x^2/2 + O(x^3)$ for $x \in (-1, 1)$, we have, uniformly for $\|h\| \leq M$ and $i = 1, \dots, n$, that

$$\begin{aligned} &\log \frac{p(X_i|\theta_0 + n^{-1/2}h)}{p(X_i|\theta_0)} \\ &= \frac{h_1}{2\sqrt{n}}X_i + \left(\frac{h_2}{4\sqrt{n}} - \frac{h_1^2}{4n}\right)(1 - X_i^2) + \frac{h_2^2}{16n}X_i^4 - \frac{h_2^2}{8n}X_i^2 - \frac{h_2^2}{16n} + \frac{3h_1h_2}{4n}X_i \\ &\quad - \frac{h_1h_2}{4n}X_i^3 - \frac{h_1^2}{8n}X_i^2 - \frac{h_2^2}{32n}(1 - X_i^2)^2 - \frac{h_1h_2}{8n}X_i(1 - X_i^2) + O_{P_{\theta_0}^n}\left(\frac{\log^3 n}{n^{3/2}}\right). \end{aligned}$$

Thus,

$$\begin{aligned} \log \frac{p_n(\mathbf{X}^n|\theta_0 + n^{-1/2}h)}{p_n(\mathbf{X}^n|\theta_0)} &= \sum_{i=1}^n \log \frac{p(X_i|\theta_0 + n^{-1/2}h)}{p(X_i|\theta_0)} \\ &= \frac{h_1}{2\sqrt{n}} \sum_{i=1}^n X_i + \frac{h_2}{4\sqrt{n}} \sum_{i=1}^n (1 - X_i^2) - \frac{h_1^2}{8} - \frac{h_2^2}{16} + o_{P_{\theta_0}^n}(1), \end{aligned}$$

where the $o_{P_{\theta_0}^n}(1)$ term is uniform for $\|h\| \leq M$. This verifies Assumption 1.

Now we verify Assumption 5. We have

$$\begin{aligned} D_{1/2}(\theta_0||\theta) &= -2 \log \int \sqrt{p(X|\theta)p(X|\theta_0)} d\mu \\ &\geq 2 \left(1 - \int \sqrt{p(X|\theta)p(X|\theta_0)} d\mu\right) \\ &= \int \left(\sqrt{p(X|\theta)} - \sqrt{p(X|\theta_0)}\right)^2 d\mu \\ &\geq \frac{1}{4} \left(\int |p(X|\theta) - p(X|\theta_0)| d\mu\right)^2 \\ &= \frac{1}{16} \left(\int |\sqrt{\tau}\phi(\sqrt{\tau}(X - \xi)) - \phi(X)| d\mu\right)^2. \end{aligned}$$

Note that

$$\begin{aligned} \int |\sqrt{\tau}\phi(\sqrt{\tau}(X - \xi)) - \phi(X)| d\mu &\geq \left| \int \exp(iX)\sqrt{\tau}\phi(\sqrt{\tau}(X - \xi)) - \exp(itX)\phi(X) d\mu \right| \\ &= |\exp(i\xi - 1/(2\tau)) - \exp(-1/2)|. \end{aligned}$$

The last display has two consequences. On the one hand,

$$\int |\sqrt{\tau}\phi(\sqrt{\tau}(X - \xi)) - \phi(X)| d\mu \geq |\sin \xi| \exp(-1/(2\tau)).$$

Hence if $(\xi, \tau)^\top$ is close enough to $(0, 1)^\top$, then

$$\int |\sqrt{\tau}\phi(\sqrt{\tau}(X - \xi)) - \phi(X)| d\mu \gtrsim |\xi|.$$

On the other hand, it is not hard to see that

$$|\exp(i\xi - 1/(2\tau)) - \exp(-1/2)| \geq |\exp(-1/(2\tau)) - \exp(-1/2)|.$$

Hence if $(\xi, \tau)^\top$ close enough to $(0, 1)^\top$, then

$$\int |\sqrt{\tau}\phi(\sqrt{\tau}(X - \xi)) - \phi(X)| d\mu \gtrsim |\tau - 1|.$$

The above equalities imply that there exist $\delta > 0$ and $C > 0$ such that for $\sqrt{\xi^2 + (\tau - 1)^2} < \delta$,

$$D_{1/2}(\theta_0 || \theta) \geq C(\xi^2 + (\tau - 1)^2).$$

We turn to the case $\sqrt{\xi^2 + (\tau - 1)^2} \geq \delta$. We have

$$\begin{aligned} D_{1/2}(\theta_0 || \theta) &\geq \frac{1}{16} \left(\int |\sqrt{\tau}\phi(\sqrt{\tau}(X - \xi)) - \phi(X)| d\mu \right)^2 \\ &\geq \frac{1}{16} \left(\int \left(\sqrt{\sqrt{\tau}\phi(\sqrt{\tau}(X - \xi))} - \sqrt{\phi(X)} \right)^2 d\mu \right)^2 \\ &= \frac{1}{4} \left(1 - \sqrt{\frac{2\sqrt{\tau}}{1+\tau}} \exp \left\{ -\frac{1}{4} \frac{\tau\xi^2}{1+\tau} \right\} \right)^2. \end{aligned}$$

Note that if $\sqrt{\xi^2 + (\tau - 1)^2} \geq \delta$, $(\tau - 1)^2 \geq \delta^2/2$ or else $(\tau - 1)^2 < \delta^2/2$ and $\xi^2 \geq \delta^2/2$. If $(\tau - 1)^2 \geq \delta^2/2$, then

$$1 - \sqrt{\frac{2\sqrt{\tau}}{1+\tau}} \exp \left\{ -\frac{1}{4} \frac{\tau\xi^2}{1+\tau} \right\} \geq 1 - \sqrt{\frac{2\sqrt{\tau}}{1+\tau}},$$

which obviously has a positive lower bound. On the other hand, suppose $(\tau - 1)^2 < \delta^2/2$ and $\xi^2 \geq \delta^2/2$, then

$$1 - \sqrt{\frac{2\sqrt{\tau}}{1+\tau}} \exp \left\{ -\frac{1}{4} \frac{\tau\xi^2}{1+\tau} \right\} \geq 1 - \exp \left\{ -\frac{1}{4} \frac{\tau\xi^2}{1+\tau} \right\} \geq 1 - \exp \left\{ -\frac{\delta^2}{8} \frac{1 - \delta/\sqrt{2}}{2 - \delta/\sqrt{2}} \right\}.$$

Thus, $D_{1/2}(\theta_0 || \theta)$ has a positive lower bound for $\sqrt{\xi^2 + (\tau - 1)^2} \geq \delta$. This verifies Assumption 5.

If $\sqrt{n}((\xi, \sigma^2) - (0, 1))^\top \rightarrow (\eta_1, \eta_2)^\top$, then $\sqrt{n}((\xi, \tau) - (0, 1))^\top \rightarrow (\eta_1, -\eta_2)^\top$ and the conclusion follows from Theorem 1. \square

To prove Theorem 3, the following result is useful.

Proposition 8. *Suppose the conditions of Theorem 3 holds. Let $A(M_n) = \{(\omega, \xi) : \omega(2\Phi(|\xi|/2) - 1) \leq M_n n^{-1/2}\}$. Let $0 < t < 1$ be a constant. If $M_n \geq \sqrt{\log n / (2(t \wedge (1-t)))}$, then*

$$P_{\theta_0}^n \int_{A(M_n)^c} \left[\prod_{i=1}^n \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} \right]^t \pi(\omega, \xi) d\omega d\mu = o(n^{-1/2}).$$

Proof. It can be seen that

$$\begin{aligned} & P_{\theta_0}^n \int_{A(M_n)^c} \left[\prod_{i=1}^n \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} \right]^t \pi(\omega, \xi) d\omega d\xi \\ &= \int_{A(M_n)^c} \left(\int p(X_1|\omega, \xi)^t p(X_1|0, 0)^{1-t} d\mu \right)^n \pi(\omega, \xi) d\omega d\xi. \end{aligned}$$

Note that

$$\begin{aligned} & \int p(X_i|\omega, \xi)^t p(X_i|0, 0)^{1-t} d\mu \\ & \leq \left(\int \sqrt{p(X_i|\omega, \xi)p(X_i|0, 0)} d\mu \right)^{2(t \wedge (1-t))} \\ & = \left(1 - \frac{1}{2} \int (\sqrt{p(X_i|\omega, \xi)} - \sqrt{p(X_i|0, 0)})^2 d\mu \right)^{2(t \wedge (1-t))} \\ & \leq \exp \left(- (t \wedge (1-t)) \int (\sqrt{p(X_i|\omega, \xi)} - \sqrt{p(X_i|0, 0)})^2 d\mu \right) \\ & \leq \exp \left(- \frac{1}{4} (t \wedge (1-t)) \left(\int |p(X_i|\omega, \xi) - p(X_i|0, 0)| d\mu \right)^2 \right) \\ & = \exp \left(- \frac{1}{4} (t \wedge (1-t)) \omega^2 \left(\int |\phi(X_i - \xi) - \phi(X_i)| d\mu \right)^2 \right) \\ & = \exp \left(- (t \wedge (1-t)) \omega^2 (2\Phi(|\xi|/2) - 1)^2 \right). \end{aligned}$$

The last display implies that

$$\begin{aligned} & P_{\theta_0}^n \int_{A(M_n)^c} \left[\prod_{i=1}^n \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} \right]^t \pi_\omega(\omega) \pi_\xi(\xi) d\omega d\xi \\ & \leq \int_{A(M_n)^c} \exp \left[- (t \wedge (1-t)) n \omega^2 (2\Phi(|\xi|/2) - 1)^2 \right] \pi_\omega(\omega) \pi_\xi(\xi) d\omega d\xi \\ & \leq \int_{A(M_n)^c} \exp \left[- (t \wedge (1-t)) M_n^2 \right] \pi_\omega(\omega) \pi_\xi(\xi) d\omega d\xi \\ & \leq n^{-1/2} \int_{A(M_n)^c} \pi_\omega(\omega) \pi_\xi(\xi) d\omega d\xi \\ & = o(n^{-1/2}). \end{aligned}$$

This completes the proof. □

Proof of Theorem 3. We have

$$\sum_{i=1}^n \log \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} = \sum_{i=1}^n \log \left(1 + \omega \left(\exp(\xi X_i - \xi^2/2) - 1 \right) \right) = \sum_{i=1}^n \log(1 + \omega \xi Y_i),$$

where $Y_i = (\exp(\xi X_i - \xi^2/2) - 1)/\xi$ if $\xi \neq 0$ and $Y_i = X_i$ if $\xi = 0$.

Let $r > 1/2$ and $s < 1/4$, on $A((\log n)^r) \cap \{\omega \geq n^{-s}\}$, we have $|\xi| = O((\log n)^r/n^{1/2-s})$. It is known that $\max_{1 \leq i \leq n} |X_i| = O_P(\sqrt{\log n})$. On $A((\log n)^r) \cap \{\omega \geq n^{-s}\}$, we have $\max_{1 \leq i \leq n} |\xi X_i - \xi^2/2| \leq |\xi| \max_{1 \leq i \leq n} |X_i| + \xi^2/2 = O_P(|\xi|(\log n)^{1/2})$. Then on $A((\log n)^r) \cap \{\omega \geq n^{-s}\}$, uniformly for $i = 1, \dots, n$, we have

$$\begin{aligned} Y_i &= \xi^{-1} \left(\xi X_i - \xi^2/2 + \frac{1}{2}(\xi X_i - \xi^2/2)^2 + O_{P_{\theta_0}^n}(|\xi|^3(\log n)^{3/2}) \right) \\ &= X_i - \frac{1}{2}\xi + \frac{1}{2}\xi X_i^2 - \frac{1}{2}\xi^2 X_i + \frac{1}{8}\xi^3 + O_{P_{\theta_0}^n}(|\xi|^2(\log n)^{3/2}) \\ &= X_i + \frac{1}{2}\xi(X_i^2 - 1) + O_{P_{\theta_0}^n}(|\xi|^2(\log n)^{3/2}). \end{aligned}$$

In particular, on $A((\log n)^r) \cap \{\omega \geq n^{-s}\}$, we have $\max_{1 \leq i \leq n} |Y_i| = O_{P_{\theta_0}^n}(\sqrt{\log n})$. On $A((\log n)^r) \cap \{\omega \geq n^{-s}\}$, we have $\omega \xi = O((\log n)^r/\sqrt{n})$, then by Taylor expansion,

$$\begin{aligned} \sum_{i=1}^n \log(1 + \omega \xi Y_i) &= \omega \xi \sum_{i=1}^n Y_i - \frac{1}{2} \omega^2 \xi^2 \sum_{i=1}^n Y_i^2 + O_{P_{\theta_0}^n}(n \omega^3 \xi^3 (\log n)^{3/2}) \\ &= \omega \xi \sum_{i=1}^n Y_i - \frac{1}{2} \omega^2 \xi^2 \sum_{i=1}^n Y_i^2 + o_{P_{\theta_0}^n}(1). \end{aligned}$$

Note that

$$\begin{aligned} \omega \xi \sum_{i=1}^n Y_i &= \omega \xi \sum_{i=1}^n X_i + \frac{1}{2} \omega \xi^2 \sum_{i=1}^n (X_i^2 - 1) + O_{P_{\theta_0}^n}(n \omega |\xi|^3 (\log n)^{3/2}) \\ &= \omega \xi \sum_{i=1}^n X_i + O_{P_{\theta_0}^n} \left(\frac{(\log n)^{3r+3/2}}{n^{1/2-2s}} \right) \\ &= \omega \xi \sum_{i=1}^n X_i + o_{P_{\theta_0}^n}(1). \end{aligned}$$

On the other hand, $\omega^2 \xi^2 \sum_{i=1}^n Y_i^2 = n \omega^2 \xi^2 + o_{P_{\theta_0}^n}(1)$. Then uniformly on $A((\log n)^r) \cap \{\omega \geq n^{-s}\}$,

$$\sum_{i=1}^n \log \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} = \omega \xi \sum_{i=1}^n X_i - \frac{1}{2} n \omega^2 \xi^2 + o_{P_{\theta_0}^n}(1). \quad (28)$$

As a result,

$$\begin{aligned} &\int_{A((\log n)^r) \cap \{\omega \geq n^{-s}\}} \left[\prod_{i=1}^n \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} \right]^t \pi(\omega, \xi) d\omega d\xi \\ &= (1 + o_{P_{\theta_0}^n}(1)) \int_{A((\log n)^r) \cap \{\omega \geq n^{-s}\}} \exp \left\{ t \omega \xi \sum_{i=1}^n X_i - \frac{1}{2} n t \omega^2 \xi^2 \right\} \pi(\omega, \xi) d\omega d\xi. \end{aligned}$$

Note that on $A((\log n)^r) \cap \{\omega \geq n^{-s}\}$, $\pi_\xi(\xi) = (1 + o(1))\pi_\xi(0)$. Then

$$\begin{aligned}
& \int_{A((\log n)^r) \cap \{\omega \geq n^{-s}\}} \exp \left\{ t\omega\xi \sum_{i=1}^n X_i - \frac{1}{2}nt\omega^2\xi^2 \right\} \pi(\omega, \xi) d\omega d\xi \\
&= (1 + o_{P_{\theta_0}^n}(1))\pi_\xi(0) \int_{A((\log n)^r) \cap \{\omega \geq n^{-s}\}} \exp \left\{ t\omega\xi \sum_{i=1}^n X_i - \frac{1}{2}nt\omega^2\xi^2 \right\} \pi_\omega(\omega) d\omega d\xi \\
&= (1 + o_{P_{\theta_0}^n}(1))\pi_\xi(0) \int_{n^{-s}}^1 \pi_\omega(\omega) d\omega \int_{-2\Phi^{-1}((\log n)^r/(2\omega\sqrt{n})+1/2)}^{2\Phi^{-1}((\log n)^r/(2\omega\sqrt{n})+1/2)} \exp \left\{ t\omega\xi \sum_{i=1}^n X_i - \frac{1}{2}nt\omega^2\xi^2 \right\} d\xi.
\end{aligned}$$

By direct calculation, we have

$$\begin{aligned}
& \int_{-2\Phi^{-1}((\log n)^r/(2\omega\sqrt{n})+1/2)}^{2\Phi^{-1}((\log n)^r/(2\omega\sqrt{n})+1/2)} \exp \left\{ t\omega\xi \sum_{i=1}^n X_i - \frac{1}{2}nt\omega^2\xi^2 \right\} d\xi \\
&= \frac{1}{\omega} \sqrt{\frac{2\pi}{tn}} \exp \left\{ \frac{t}{2n} \left(\sum_{i=1}^n X_i \right)^2 \right\} \left[\Phi \left(2\sqrt{tn}\omega\Phi^{-1} \left(\frac{(\log n)^r}{2\omega\sqrt{n}} + \frac{1}{2} \right) + \frac{1}{2} \right) - \sqrt{\frac{t}{n}} \sum_{i=1}^n X_i \right] \\
&\quad - \Phi \left(-2\sqrt{tn}\omega\Phi^{-1} \left(\frac{(\log n)^r}{2\omega\sqrt{n}} + \frac{1}{2} \right) - \sqrt{\frac{t}{n}} \sum_{i=1}^n X_i \right).
\end{aligned}$$

Since

$$2\sqrt{tn}\omega\Phi^{-1} \left(\frac{(\log n)^r}{2\omega\sqrt{n}} + \frac{1}{2} \right) \geq \sqrt{2\pi t}(\log n)^r,$$

we have

$$\begin{aligned}
& \int_{-2\Phi^{-1}((\log n)^r/(2\omega\sqrt{n})+1/2)}^{2\Phi^{-1}((\log n)^r/(2\omega\sqrt{n})+1/2)} \exp \left\{ t\omega\xi \sum_{i=1}^n X_i - \frac{1}{2}nt\omega^2\xi^2 \right\} d\xi \\
&= \frac{1}{\omega} \sqrt{\frac{2\pi}{tn}} \exp \left\{ \frac{t}{2n} \left(\sum_{i=1}^n X_i \right)^2 \right\} (1 + o_{P_{\theta_0}^n}(1)),
\end{aligned}$$

where the $o_{P_{\theta_0}^n}(1)$ term is uniform for ω . Thus,

$$\begin{aligned}
& \int_{A((\log n)^r) \cap \{\omega \geq n^{-s}\}} \exp \left\{ t\omega\xi \sum_{i=1}^n X_i - \frac{1}{2}nt\omega^2\xi^2 \right\} \pi(\omega, \xi) d\omega d\xi \\
&= (1 + o_{P_{\theta_0}^n}(1))\pi_\xi(0) \sqrt{\frac{2\pi}{tn}} \exp \left\{ \frac{t}{2n} \left(\sum_{i=1}^n X_i \right)^2 \right\} \int_0^1 \frac{1}{\omega} \pi_\omega(\omega) d\omega.
\end{aligned}$$

Now we consider the event $A((\log n)^r) \cap \{\omega \leq n^{-s}\}$. By Theorem 2 of Liu and Shao (2004), we have

$$\sup_{\omega \in [0,1], t \in \mathbb{R}} \sum_{i=1}^n (\log p(X_i|\omega, \xi) - \log p(X_i|0, 0)) = O_{P_{\theta_0}^n}(\log \log n).$$

Thus,

$$\begin{aligned}
& \int_{A((\log n)^r) \cap \{\omega \leq n^{-s}\}} \left[\prod_{i=1}^n \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} \right]^t \pi(\omega, \xi) d\omega d\xi \\
&= \exp \left\{ O_{P_{\theta_0}^n}(\log(\log n)) \right\} \Pi(\omega(2\Phi(|\xi|/2) - 1) \leq (\log n)^r n^{-1/2}, \omega \leq n^{-s}).
\end{aligned}$$

We break the probability into two parts:

$$\begin{aligned} & \Pi(\omega(2\Phi(|\xi|/2) - 1) \leq (\log n)^r n^{-1/2}, \omega \leq n^{-s}) \\ & \leq \Pi(\omega(2\Phi(|\xi|/2) - 1) \leq (\log n)^r n^{-1/2}, \omega \leq 2(\log n)^r n^{-1/2}) \\ & \quad + \Pi(\omega(2\Phi(|\xi|/2) - 1) \leq (\log n)^r n^{-1/2}, 2(\log n)^r n^{-1/2} \leq \omega \leq n^{-s}). \end{aligned}$$

The first probability satisfies

$$\begin{aligned} & \Pi(\omega(2\Phi(|\xi|/2) - 1) \leq (\log n)^r n^{-1/2}, \omega \leq 2(\log n)^r n^{-1/2}) \\ & \leq \Pi(\omega \leq 2(\log n)^r n^{-1/2}) \\ & \lesssim \int_0^{2(\log n)^r n^{-1/2}} \omega^{\alpha_1-1} d\omega \\ & \lesssim \left(\frac{(\log n)^r}{\sqrt{n}} \right)^{\alpha_1}. \end{aligned}$$

Next we deal with the second probability. On the event of the second probability, we have $(2\Phi(|\xi|/2) - 1) \leq \omega^{-1}(\log n)^r n^{-1/2} \leq 1/2$, which implies the boundedness of ξ . It follows that $|\xi| \leq C\omega^{-1}(\log n)^r n^{-1/2}$ for some constant $C > 0$ on this event. Thus,

$$\begin{aligned} & \Pi(\omega(2\Phi(|\xi|/2) - 1) \leq (\log n)^r n^{-1/2}, 2(\log n)^r n^{-1/2} \leq \omega \leq n^{-s}) \\ & \leq \Pi(|\xi| \leq C\omega^{-1}(\log n)^r n^{-1/2}, 2(\log n)^r n^{-1/2} \leq \omega \leq n^{-s}) \\ & \leq \Pi(|\xi| \leq C\omega^{-1}(\log n)^r n^{-1/2}, \omega \leq n^{-s}) \\ & \lesssim \int_0^{n^{-s}} \omega^{\alpha_1-1} d\omega \int_{-C\omega^{-1}(\log n)^r n^{-1/2}}^{C\omega^{-1}(\log n)^r n^{-1/2}} \pi_\xi(\xi) d\xi. \end{aligned}$$

There exists $\epsilon > 0$ and $M > 0$ such that $\pi_\xi(\xi) \leq M$ for $\xi \in [-\epsilon, \epsilon]$. Then

$$\begin{aligned} & \int_0^{n^{-s}} \omega^{\alpha_1-1} d\omega \int_{-C\omega^{-1}(\log n)^r n^{-1/2}}^{C\omega^{-1}(\log n)^r n^{-1/2}} \pi_\xi(\xi) d\xi \\ & \leq \int_0^{C(\log n)^r/(\epsilon\sqrt{n})} \omega^{\alpha_1-1} d\omega + \int_{C(\log n)^r/(\epsilon\sqrt{n})}^{n^{-s}} 2MC\omega^{\alpha_1-2}(\log n)^r n^{-1/2} d\omega \\ & \lesssim \left(\frac{(\log n)^r}{\sqrt{n}} \right)^{\alpha_1} + \frac{(\log n)^r}{\sqrt{n}} \left(\left(\frac{(\log n)^r}{\sqrt{n}} \right)^{\alpha_1-1} \vee \left(\frac{1}{n^s} \right)^{\alpha_1-1} \right) \\ & = \left(\frac{(\log n)^r}{\sqrt{n}} \right)^{\alpha_1} \vee \frac{(\log n)^r}{n^{1/2+s(\alpha_1-1)}}. \end{aligned}$$

It follows that

$$\begin{aligned} & \int_{A((\log n)^r) \cap \{\omega \leq n^{-s}\}} \left[\prod_{i=1}^n \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} \right]^t \pi(\omega, \xi) d\omega d\xi \\ & = \exp \{ O_{P_{\theta_0}^2}(\log(\log n)) \} \left(\left(\frac{(\log n)^r}{\sqrt{n}} \right)^{\alpha_1} \vee \frac{(\log n)^r}{n^{1/2+s(\alpha_1-1)}} \right) = o_{P_{\theta_0}^n}(n^{-1/2}). \end{aligned}$$

Combine these arguments and Proposition 8, we have

$$\begin{aligned}
& \int \left[\prod_{i=1}^n \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} \right]^t \pi(\omega, \xi) d\omega d\xi \\
&= \left(\int_{A((\log n)^c)} + \int_{A((\log n)^r) \cap \{\omega < n^{-s}\}} + \int_{A((\log n)^r) \cap \{\omega \geq n^{-s}\}} \right) \left[\prod_{i=1}^n \frac{p(X_i|\omega, \xi)}{p(X_i|0, 0)} \right]^t \pi(\omega, \xi) d\omega d\xi \\
&= (1 + o_{P_{\theta_0}^n}(1)) \pi_{\xi}(0) \sqrt{\frac{2\pi}{tn}} \exp \left\{ \frac{t}{2n} \left(\sum_{i=1}^n X_i \right)^2 \right\} \int_0^1 \frac{1}{\omega} \pi_{\omega}(\omega) d\omega.
\end{aligned}$$

This implies that

$$2 \log \Lambda_{a,b} = -\log(1 + a/b) + \frac{a}{n} \left(\sum_{i=1}^n X_i \right)^2 + o_{P_{\theta_0}^n}(1).$$

Then the conclusion of (i) holds since $(\sum_{i=1}^n X_i)^2/n$ weakly converges to $\chi^2(1)$ under $P_{\theta_0}^n$.

Now we prove (ii). Suppose that $\theta_n = (\omega, \xi)$ satisfies that for some $s < 1/4$, $\omega \geq n^{-s}$ for large n and $\sqrt{n}\omega\xi \rightarrow \eta$. Then it follows from (28) and Le Cam's first lemma (van der Vaart, 1998, Theorem 6.4) that $P_{\theta_n}^n$ and $P_{\theta_0}^n$ are mutually contiguous. As a result,

$$2 \log \Lambda_{a,b} = -\log(1 + a/b) + \frac{a}{n} \left(\sum_{i=1}^n X_i \right)^2 + o_{P_{\theta_n}^n}(1).$$

Note that (28) implies that

$$\left(n^{-1/2} \sum_{i=1}^n X_i, \log \frac{p_n(\mathbf{X}^n|\theta)}{p_n(\mathbf{X}^n|\theta_0)} \right)^\top \overset{P_{\theta_0}^n}{\rightsquigarrow} \mathcal{N}_2 \left(\begin{pmatrix} 0 \\ -\eta^2/2 \end{pmatrix}, \begin{pmatrix} 1 & \eta \\ \eta & \eta^2 \end{pmatrix} \right).$$

By Le Cam's third lemma (van der Vaart, 1998, Example 6.7), we have

$$\sum_{i=1}^n X_i \overset{P_{\theta_n}^n}{\rightsquigarrow} \mathcal{N}(\eta, 1).$$

This proves the conclusion of (ii). □

References

- Aitkin, M. (1991). Posterior bayes factors. *Journal of the Royal Statistical Society: Series B (Methodological)*, 53(1):111–128.
- Bayarri, M. J. and Garca-Donato, G. (2008). Generalization of jeffreys divergence-based priors for bayesian hypothesis testing. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 70(5):981–1003.
- Berger, J. O. and Pericchi, L. R. (1996). The intrinsic bayes factor for model selection and prediction. *Journal of the American Statistical Association*, 91(433):109–122.

- Bhattacharya, A., Pati, D., and Yang, Y. (2019). Bayesian fractional posteriors. *The Annals of Statistics*, 47(1):39–66.
- Blei, D. M., Kucukelbir, A., and McAuliffe, J. D. (2017). Variational inference: A review for statisticians. *arxiv*.
- Chen, J. (2017). On finite mixture models. *Statistical Theory and Related Fields*, 1(1):15–27.
- Clarke, B. S. and Barron, A. R. (1990). Information-theoretic asymptotics of bayes methods. *IEEE Transactions on Information Theory*, 36(3):453–471.
- Crisp, A. and Burridge, J. (1994). A note on nonregular likelihood functions in heteroscedastic regression models. *Biometrika*, 81(3):585–587.
- De Santis, F. and Spezzaferri, F. (1997). Alternative bayes factors for model selection. *Canadian Journal of Statistics*, 25(4):503–515.
- Fang, K. T., Kotz, S., and Ng, K. W. (1990). *Symmetric multivariate and related distributions*, volume 36 of *Monographs on Statistics and Applied Probability*. Chapman and Hall, Ltd., London.
- Friel, N. and Wyse, J. (2012). Estimating the evidence - a review. *Statistica Neerlandica*, 66(3):288–308.
- Ghosal, S., Ghosh, J. K., and van der Vaart, A. W. (2000). Convergence rates of posterior distributions. *Ann. Statist.*, 28(2):500–531.
- Hall, P. and Stewart, M. (2005). Theoretical analysis of power in a two-component normal mixture model. *Journal of Statistical Planning and Inference*, 134(1):158 – 179.
- Jeffreys, H. (1931). *Scientific Inference*. Cambridge University Press, Cambridge, 1 edition.
- Jeffreys, H. (1961). *Theory of probability*. Third edition. Clarendon Press, Oxford.
- Kass, R. E. and Raftery, A. E. (1995). Bayes factors. *Journal of the American Statistical Association*, 90(430):773–795.
- Kass, R. E. and Wasserman, L. (1995). A reference bayesian test for nested hypotheses and its relationship to the schwarz criterion. *Journal of the American Statistical Association*, 90(431):928–934.
- Kleijn, B. J. K. and van der Vaart, A. W. (2012). The Bernstein-Von-Mises theorem under misspecification. *Electronic Journal of Statistics*, 6:354–381.
- Le Cam, L. (1990). Maximum likelihood: An introduction. *International Statistical Review*, 58(2):153–171.

- Li, K.-H. and Chan, N. N. (2000). Degeneracy in heteroscedastic regression models. *Journal of Multivariate Analysis*, 74(2):282–295.
- Li, Y. and Turner, R. E. (2016). Rényi divergence variational inference. In Lee, D. D., Sugiyama, M., Luxburg, U. V., Guyon, I., and Garnett, R., editors, *Advances in Neural Information Processing Systems 29*, pages 1073–1081. Curran Associates, Inc.
- Liu, X. and Shao, Y. (2004). Asymptotics for the likelihood ratio test in a two-component normal mixture model. *Journal of Statistical Planning and Inference*, 123(1):61 – 81.
- O’Hagan, A. (1995). Fractional Bayes factors for model comparison. *Journal of the Royal Statistical Society. Series B. Methodological*, 57(1):99–138. With discussion and a reply by the author.
- Pati, D., Bhattacharya, A., and Yang, Y. (2017). On Statistical Optimality of Variational Bayes. *ArXiv e-prints*.
- Perez, J. M. (2002). Expected-posterior prior distributions for model selection. *Biometrika*, 89(3):491–512.
- Shafer, G. (1982). Lindley’s paradox. *Journal of the American Statistical Association*, 77(378):325–351. With discussion and with a reply by the author.
- Shen, X. and Wasserman, L. (2001). Rates of convergence of posterior distributions. *Annals of Statistics*, 29(3):687–714.
- van der Vaart, A. and Ghosal, S. (2007). Convergence rates of posterior distributions for noniid observations. *Annals of Statistics*, 35(1):192–223.
- van der Vaart, A. W. (1998). *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge.
- van Erven, T. and Harremoës, P. (2014). Rényi divergence and kullback-leibler divergence. *IEEE Transactions on Information Theory*, 60(7):3797–3820.
- Walker, S. G. and Hjort, N. L. (2001). On bayesian consistency. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 63(4):811–821.
- Wang, Y. and Blei, D. M. (2017). Frequentist Consistency of Variational Bayes. *ArXiv e-prints*.
- Wilks, S. S. (1938). The large-sample distribution of the likelihood ratio for testing composite hypotheses. *Annals of Mathematical Statistics*, 9(1):60–62.
- Yang, Y., Pati, D., and Bhattacharya, A. (2017). α -Variational Inference with Statistical Guarantees. *ArXiv e-prints*.