

2022-2023 Spring  
MAT124 Midterm  
(08/05/2023)

1. Sketch the traces of the following surfaces with the coordinate planes  $x = 0$ ,  $y = 0$ , and  $z = 0$ , and then sketch the graphs of them.

(a)  $\frac{x^2}{16} + \frac{y^2}{4} - z^2 = 1$

(b)  $z = \frac{x^2}{4} + \frac{y^2}{4} - 6$

2. Show that the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{y^3 \sqrt{x}}{2(x^2 + y^4)}$$

does not exist.

3. Find the equation of the plane passing through the point  $P_0(0, 1, 2)$  and which is perpendicular to the line that is tangent to the curve of intersection of the surfaces

$$xz^2 - 2xy + y^2 = 2 \quad \text{and} \quad xz - x^2y + z^2 = 1$$

at the point  $P_1(0, \sqrt{2}, 1)$ .

4. Find  $\frac{\partial w}{\partial s}$  where  $w = xy \ln(1 + \sqrt{x^2 + y^2}) + xz$  and  $x = t + s$ ,  $y = e^s$ ,  $z = \ln(s^2 + t)$ .

5. Use incremental approximation to estimate the value  $\tan((0.97) \cdot (2.05)^2)$ .

6. Find the direction vector in which the function

$$f(x, y, z) = \sqrt{x + yz}$$

has the minimum rate of change at the point  $(1, 1, 3)$ . Also, find this rate of change.


7. Find the absolute extrema of

$$f(x, y) = x^2 + y - xy + 4$$


on the triangular region with vertices  $(0, 0)$ ,  $(4, 0)$ ,  $(0, 4)$ .


1.

(a)



(b)





2. Apply the Two-Path Test.

$$x = y \implies \lim_{(x,y) \rightarrow (0,0)} \frac{y^3 \sqrt{x}}{2(x^2 + y^4)} = \lim_{x \rightarrow 0} \frac{x^{7/2}}{2(x^2 + x^4)} = \lim_{x \rightarrow 0} \frac{x^{3/2}}{2(1 + x^2)} = \frac{0}{2} = 0$$

$$x = y^2 \implies \lim_{(x,y) \rightarrow (0,0)} \frac{y^3 \sqrt{x}}{2(x^2 + y^4)} = \lim_{y \rightarrow 0} \frac{y^4}{4y^4} = \lim_{y \rightarrow 0} \frac{1}{4} = \frac{1}{4}$$

Since  $0 \neq \frac{1}{4}$ , by the Two-Path Test, the limit does not exist.

3. Let  $f(x, y, z) = xz^2 - 2xy + y^2 = 2$  and  $g(x, y, z) = xz - x^2y + z^2 = 1$  be the level surfaces. The cross product of the gradient of these functions give us the vector that is parallel to the line of intersection. Compute the gradients.

$$\nabla f = \langle z^2 - 2y, -2x + 2y, 2xz \rangle, \quad \nabla g = \langle z - 2xy, -x^2, x + 2z \rangle$$

$$\nabla f|_{(0, \sqrt{2}, 1)} = \langle 1 - 2\sqrt{2}, 2\sqrt{2}, 0 \rangle, \quad \nabla g|_{(0, \sqrt{2}, 1)} = \langle 1, 0, 2 \rangle$$

Find the cross product.

$$\mathbf{n} = \nabla f \times \nabla g = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 - 2\sqrt{2} & 2\sqrt{2} & 0 \\ 1 & 0 & 2 \end{vmatrix}$$

$$= \mathbf{i} \begin{vmatrix} 2\sqrt{2} & 0 \\ 0 & 2 \end{vmatrix} - \mathbf{j} \begin{vmatrix} 1 - 2\sqrt{2} & 0 \\ 1 & 2 \end{vmatrix} + \mathbf{k} \begin{vmatrix} 1 - 2\sqrt{2} & 2\sqrt{2} \\ 1 & 0 \end{vmatrix}$$

$$= (2\sqrt{2} \cdot 2 - 0 \cdot 0) \mathbf{i} - [(1 - 2\sqrt{2}) \cdot 2 - 0 \cdot 1] \mathbf{j} + [(1 - 2\sqrt{2}) \cdot 0 - 2\sqrt{2} \cdot 1] \mathbf{k}$$

$$= 4\sqrt{2} \mathbf{i} + (4\sqrt{2} - 2) \mathbf{j} - 2\sqrt{2} \mathbf{k}$$

The line of intersection is the normal line of the plane. Therefore,  $\mathbf{n}$  is the normal vector of the plane. The equation of the plane with the normal vector  $\mathbf{n}$  and containing the point  $P(x_0, y_0, z_0)$  is

$$n_1(x - x_0) + n_2(y - y_0) + n_3(z - z_0) = 0$$

Therefore, the equation for this plane is

$$4\sqrt{2}(x - 0) + (4\sqrt{2} - 2)(y - 1) - 2\sqrt{2}(z - 2) = 0$$

$$\boxed{2x\sqrt{2} + y(2\sqrt{2} - 1) - z\sqrt{2} + 1 = 0}$$

4. Apply the chain rule.

$$\frac{\partial w}{\partial s} = \frac{\partial w}{\partial x} \cdot \frac{\partial x}{\partial s} + \frac{\partial w}{\partial y} \cdot \frac{\partial y}{\partial s} + \frac{\partial w}{\partial z} \cdot \frac{\partial z}{\partial s}$$

Compute the partial derivatives.

$$\frac{\partial w}{\partial x} = y \ln(1 + \sqrt{x^2 + y^2}) + xy \cdot \frac{1}{1 + \sqrt{x^2 + y^2}} \cdot \left( \frac{x}{\sqrt{x^2 + y^2}} \right) + z$$

$$\frac{\partial w}{\partial y} = x \ln(1 + \sqrt{x^2 + y^2}) + xy \cdot \frac{1}{1 + \sqrt{x^2 + y^2}} \cdot \left( \frac{y}{\sqrt{x^2 + y^2}} \right), \quad \frac{\partial w}{\partial z} = x$$

$$\frac{\partial x}{\partial s} = 1, \quad \frac{\partial y}{\partial s} = e^s, \quad \frac{\partial z}{\partial s} = \frac{2s}{s^2 + t}$$

$$\begin{aligned} \frac{\partial w}{\partial s} &= \left[ y \ln(1 + \sqrt{x^2 + y^2}) + xy \cdot \frac{1}{1 + \sqrt{x^2 + y^2}} \cdot \left( \frac{x}{\sqrt{x^2 + y^2}} \right) + z \right] \cdot 1 \\ &+ \left[ x \ln(1 + \sqrt{x^2 + y^2}) + xy \cdot \frac{1}{1 + \sqrt{x^2 + y^2}} \cdot \left( \frac{y}{\sqrt{x^2 + y^2}} \right) \right] \cdot e^s + \frac{2xs}{s^2 + t} \end{aligned}$$

Write in terms of  $s$  and  $t$  rigorously.

$$\begin{aligned} \frac{\partial w}{\partial s} &= e^s \ln(1 + \sqrt{(t+s)^2 + e^{2s}}) + \frac{(t+s)^2 \cdot e^s}{\sqrt{(t+s)^2 + e^{2s}} + (t+s)^2 + e^{2s}} + \ln(s^2 + t) \\ &+ (t+s)e^s \ln(1 + \sqrt{(t+s)^2 + e^{2s}}) + \frac{(t+s) \cdot e^{3s}}{\sqrt{(t+s)^2 + e^{2s}} + (t+s)^2 + e^{2s}} \\ &+ \frac{2s(t+s)}{s^2 + t} \end{aligned}$$

5. Let  $f(x, y) = \tan(xy^2)$ . The total differential of  $f$  is

$$df = f_x dx + f_y dy = \sec^2(xy^2) \cdot y^2 dx + \sec^2(xy^2) \cdot 2xy dy$$

Since  $x = 0.97 \approx 1$  and  $y = 2.05 \approx 2$ , we may approximate the value of  $\tan(0.97 \cdot 2.05^2)$  near  $\tan(1 \cdot 2^2) = \tan 4$ . Take  $x = 1$ ,  $y = 2$ ,  $dx = 0.97 - 1 = -0.03$ ,  $dy = 2.05 - 2 = 0.05$ .

$$df = \sec^2 4 \cdot 4 \cdot (-0.03) + \sec^2 4 \cdot 4 \cdot (0.05) = 0.08 \sec^2 4$$

Since  $f(x + \Delta x, y + \Delta y) \approx f(x, y) + df$ , the value of  $\tan(0.97 \cdot 2.05^2)$  is approximately

$$\boxed{\tan 4 + 0.08 \sec^2 4}$$

6. The function  $f$  has the minimum rate of change if the gradient vector of  $f$  and the unit direction vector  $\mathbf{u}$  are antiparallel. That is, they have opposite directions.

$$\begin{aligned} \nabla f &= \left\langle \frac{1}{2\sqrt{x+yz}}, \frac{z}{2\sqrt{x+yz}}, \frac{y}{2\sqrt{x+yz}} \right\rangle \\ (\nabla f \cdot \mathbf{u})_{\min} &= |\nabla f| |u| \cos \pi = -|\nabla f| \end{aligned}$$

$$\nabla f|_{(1,1,3)} = \left\langle \frac{1}{4}, \frac{3}{4}, \frac{1}{4} \right\rangle \implies -|\nabla f| = -\sqrt{\left(\frac{1}{4}\right)^2 + \left(\frac{3}{4}\right)^2 + \left(\frac{1}{4}\right)^2} = -\frac{\sqrt{11}}{4}$$

Since  $\mathbf{u}$  has the opposite direction to that of the gradient, we may also find the components of  $\mathbf{u}$ .

$$\mathbf{u} = -\frac{\nabla f}{|\nabla f|} = -\frac{\left\langle \frac{1}{4}, \frac{3}{4}, \frac{1}{4} \right\rangle}{\frac{\sqrt{11}}{4}} = \left\langle -\frac{1}{\sqrt{11}}, -\frac{3}{\sqrt{11}}, -\frac{1}{\sqrt{11}} \right\rangle$$

$$\boxed{\text{The minimum rate of change: } -\frac{\sqrt{11}}{4}, \text{ the direction vector: } \left\langle -\frac{1}{\sqrt{11}}, -\frac{3}{\sqrt{11}}, -\frac{1}{\sqrt{11}} \right\rangle}$$

7.  $f$  is continuous on a bounded and closed set on  $\mathbb{R}$ . By the Extreme Value Theorem, the extrema must exist in the region or on the boundary.

First, determine where  $f_x = f_y = 0$  to find the critical points.

$$\begin{aligned} f_x &= 2x - y, & f_y &= 1 - x \\ f_x = f_y = 0 &\implies 2x - y = 0, & 1 - x = 0 &\implies x = 1, & y = 2 \\ && f(1, 2) &= 5 \end{aligned}$$

Take a look at the boundary. From  $(0, 0)$  to  $(4, 0)$ , we have  $y = 0$ .

$$y = 0 \implies f(x, 0) = x^2 + 4 \quad \rightarrow \quad \frac{d}{dx}(x^2 + 4) = 2x = 0 \implies x = 0$$

$$f(0,0) = 4$$

From  $(0,0)$  to  $(0,4)$ , we have  $x = 0$ .

$$x = 0 \implies f(0,y) = y + 4 \quad \rightarrow \quad \frac{d}{dy}(y + 4) = 1 \neq 0$$

From  $(4,0)$  to  $(0,4)$ , we have  $x = y$ .

$$y = 4 - x \implies f(x, 4 - x) = x^2 + (4 - x) - x(4 - x) + 4 = 2x^2 - 5x + 8$$

$$\frac{d}{dx}(2x^2 - 5x + 8) = 4x - 5 = 0 \implies x = \frac{5}{4}, \quad y = \frac{11}{4}$$

$$f\left(\frac{5}{4}, \frac{11}{4}\right) = \frac{39}{8}$$

We also have  $f(0,0) = 5$ ,  $f(4,0) = 20$ ,  $f(0,4) = 8$  from the vertices of the triangular region.

Compare all the values  $f(0,0)$ ,  $f(1,2)$ ,  $f(4,0)$ ,  $f(0,4)$ ,  $f\left(\frac{5}{4}, \frac{11}{4}\right)$ .

<p>Absolute minimum: <math>f(0,0) = 4</math>,    absolute maximum: <math>f(4,0) = 20</math>.</p>
--------------------------------------------------------------------------------------------------