

Chapter 5 Part 1 Group Work

SOLUTIONS

Problem 1

Suppose that you randomly select a student from a large population of 4th grade students and record the student's sex and number of siblings. Let $M = 1$ if the selected student is male and $M = 0$ if the selected student is female. Let S record the student's number of siblings. The proportion of the population of students falling into each category of M and S is recorded in the table below.

# of siblings:	0	1	2	3	4
Female	.10	.18	.12	.07	.04
Male	.12	.18	.14	.03	.02

- a. Verify that the above table is a valid joint pmf.
- b. Find the probability that the randomly selected student will be male with two siblings
- c. Find $P(M = 0, S \leq 2)$
- d. Find the probability that the randomly selected student will be an only child
- e. Find $F(1, 1)$

Solution

Part a

$$0.10 + 0.12 + 0.18 + 0.18 + 0.12 + 0.14 + 0.07 + 0.03 + 0.04 + 0.02 = 1$$

Part b

$$P(M = 1, S = 2) = 0.14$$

Part c

$$P(M = 0, S \leq 2) = 0.12 + 0.18 + 0.14 = 0.44$$

Part d

$$P(S = 0) = 0.10 + 0.12 = 0.22$$

Part e

$$F(1, 1) = P(M \leq 1, S \leq 1) = P(S \leq 1) = 0.10 + 0.12 + 0.18 + 0.18 = 0.58$$

Problem 2

Of nine executives in a business firm, four are married, three have never married, and two are divorced. Three of the executives are to be selected for promotion. Let Y_1 denote the number of married executives and Y_2 denote the number of never married executives among the three selected for promotion. Assuming that the three are randomly selected from the nine available, find the joint pmf of Y_1 and Y_2 .

Solution

Note that using material from Chapter 3, the joint probability function is given by

$$p(y_1, y_2) = P(Y_1 = y_1, Y_2 = y_2) = \frac{\binom{4}{y_1} \binom{3}{y_2} \binom{2}{3-y_1-y_2}}{\binom{9}{3}}, \quad 0 \leq y_1, 0 \leq y_2, y_1 + y_2 \leq 3$$

Given table format, this is

y2	y1	0	1	2	3
0	0	3/84	6/84	1/84	
1	4/84	24/84	12/84	0	
2	12/84	18/84	0	0	
3	4/84	0	0	0	

Problem 3

Let Y_1 and Y_2 have the joint pdf

$$f(y_1, y_2) = \begin{cases} e^{-(y_1+y_2)}, & y_1 > 0, y_2 > 0 \\ 0, & \text{elsewhere} \end{cases}$$

- a. What is $P(Y_1 < 1, Y_2 > 5)$? b. What is $P(Y_1 + Y_2 < 3)$?

Part a

$$\begin{aligned} P(Y_1 < 1, Y_2 > 5) &= \int_0^1 \int_5^\infty e^{-(y_1+y_2)} dy_2 dy_1 \\ &= \int_0^1 \left(-e^{-(y_1+y_2)} \Big|_{y_2=5}^\infty \right) dy_1 \\ &= \int_0^1 e^{-(y_1+5)} dy_1 \\ &= -e^{-(1+5)} + e^{-5} \\ &= .00426. \end{aligned}$$

Part b

$$\begin{aligned} P(Y_1 + Y_2 < 3) &= P(Y_1 < 3 - Y_2) \\ &= \int_0^3 \left(\int_0^{3-y_1} e^{-(y_1+y_2)} dy_2 \right) dy_1 \\ &= 1 - 4e^{-3} \\ &= .8009. \end{aligned}$$

Problem 4

Return to the scenario in Problem 1

Return to the scenario in Problem 1.

- Find the marginal distribution of M
- Find the marginal distribution of S
- Find $P(S \leq 2 | M = 1)$
- Find $P(M = 1 | S \leq 2)$

Solutions

Part a

	P(M = m)
Female (M = 0)	0.51
Male (M = 1)	0.49

Part b

# of siblings (S):	0	1	2	3	4
P(S = s)	.22	.36	.26	.10	.06

Part c

$$P(S \leq 2 | M = 1) = \frac{P(S \leq 2, M = 1)}{P(M = 1)}$$

- $P(M = 1) = 0.49$
- $P(S \leq 2, M = 1) = P(S = 0, M = 1) + P(S = 1, M = 1) + P(S = 2, M = 1) = 0.12 + 0.18 + 0.14 = 0.44$

$$P(S \leq 2 | M = 1) = \frac{0.44}{0.49} \approx 0.898$$

Part d

$$P(M = 1 | S \leq 2) = \frac{P(S \leq 2, M = 1)}{P(S \leq 2)}$$

- $P(S \leq 2) = P(S = 0) + P(S = 1) + P(S = 2) = 0.22 + 0.36 + 0.26 = 0.84$
- $P(S \leq 2, M = 1) = 0.44$ (from above)

$$P(M = 1 | S \leq 2) = \frac{0.44}{0.84} \approx 0.524$$

Problem 5

Let $f(x, y) = \frac{4}{3}(1 - xy)$, $0 \leq x \leq 1$, $0 \leq y \leq 1$.

- a. Find the marginal pdfs of X and Y .
- b. Find $P(X \leq Y/2)$

$$f(x, y) = \frac{4}{3}(1 - xy), \quad 0 \leq x \leq 1, \quad 0 \leq y \leq 1$$

$$f_X(x) = \int_0^1 f(x, y) dy = \int_0^1 \frac{4}{3}(1 - xy) dy = \frac{4}{3} \left[y - \frac{1}{2}xy^2 \right]_0^1 = \frac{4}{3} \left(1 - \frac{1}{2}x \right) = \frac{4}{3} - \frac{2}{3}x, \quad 0 < x < 1$$

$$f_Y(y) = \int_0^1 f(x, y) dx = \int_0^1 \frac{4}{3}(1 - xy) dx = \frac{4}{3} \left[x - \frac{1}{2}yx^2 \right]_0^1 = \frac{4}{3} \left(1 - \frac{1}{2}y \right) = \frac{4}{3} - \frac{2}{3}y, \quad 0 < y < 1$$

$$P(X \leq Y/2) = \int_0^1 \int_0^{y/2} f(x, y) dx dy = \int_0^1 \int_0^{y/2} \frac{4}{3}(1 - xy) dx dy = \frac{7}{24}$$

Problem 6

Let the joint pmf of X and Y be

$$f(x, y) = \frac{xy^2}{30}, \quad x = 1, 2, 3, \quad y = 1, 2$$

- a. What are the marginal distributions of X and Y ?
- b. Are X and Y independent?
- c. What is $P(X > 1)$?
- d. What is $E(X)$?

Part a

$$f_X(x) = \sum_{y=1}^2 \frac{xy^2}{30} = \frac{x \cdot 1^2}{30} + \frac{x \cdot 2^2}{30} = \frac{x}{30} + \frac{4x}{30} = \frac{5x}{30} = \frac{x}{6}, \quad x = 1, 2, 3$$

$$f_Y(y) = \sum_{x=1}^3 \frac{xy^2}{30} = \frac{1 \cdot y^2}{30} + \frac{2 \cdot y^2}{30} + \frac{3 \cdot y^2}{30} = \frac{6y^2}{30} = \frac{y^2}{5}, \quad y = 1, 2$$

Part b

Since $f_X(x)f_Y(y) = \frac{x}{6} \frac{y^2}{5} = \frac{xy^2}{30} = f(x, y)$ for all x, y , X and Y are independent.

Part c

$$P(X > 1) = f_X(2) + f_X(3) = \frac{2}{6} + \frac{3}{6} = \frac{5}{6}$$

Part d

$$E(X) = \sum_{x=1}^3 xf_X(x) = 1 \cdot \frac{1}{6} + 2 \cdot \frac{2}{6} + 3 \cdot \frac{3}{6} = \frac{14}{6} = \frac{7}{3}$$

Problem 7

Let X and Y be two continuous random variables with joint pdf $f(x, y) = \frac{3}{16}xy^2$, $0 \leq x \leq 2$, $0 \leq y \leq 2$. Are the two random variables independent?

Solution

Both are bounded between constants, and we can define $g(x) = \frac{3}{16}x$ and $h(y) = y^2$, so by Theorem 5.5 they are independent.

Problem 8

Let X and Y have the joint pdf $f(x, y) = x + y$, $0 \leq x \leq 1$, $0 \leq y \leq 1$. Find the marginal pdfs $f_X(x)$ and $f_Y(y)$ and show that X and Y are dependent.

Solution

$$f_x(x) = \int_0^1 (x + y) dy = x + \frac{1}{2}, \quad 0 \leq x \leq 1$$

$$f_y(y) = \int_0^1 (x + y) dx = y + \frac{1}{2}, \quad 0 \leq y \leq 1$$

$$f_x(x)f_y(y) = (x + \frac{1}{2})(y + \frac{1}{2}) \neq x + y$$