

HW 04 SOLUTIONS

Practice Problems

3.1

$P(Y = 0) = P(\text{no impurities}) = 0.2$, $P(Y = 1) = P(\text{exactly one impurity}) = 0.7$, $P(Y = 2) = 0.1$.

3.3

$p(2) = P(DD) = 1/6$, $p(3) = P(DGD) + P(GDD) = 2(2/4)(2/3)(1/2) = 2/6$, $p(4) = P(GGDD) + P(DGGD) + P(GDGD) = 3(2/4)(1/3)(1) = 1/2$.

3.5

There are $3! = 6$ possible ways to assign the words to the pictures. Of these, one is a perfect match, three have one match, and two have zero matches. Thus, $p(0) = 2/6$, $p(1) = 3/6$, $p(3) = 1/6$.

3.15

a. $p(0) = P(Y = 0) = (.48)^3 = .1106$, $p(1) = P(Y = 1) = 3(.48)^2(.52) = .3594$,
 $p(2) = P(Y = 2) = 3(.48)(.52)^2 = .3894$, $p(3) = P(Y = 3) = (.52)^3 = .1406$.

b. The graph is omitted.

c. $P(Y = 1) = .3594$.

3.21

Note that $E(N) = E(8\pi R^2) = 8\pi E(R^2)$. So, $E(R^2) = 21^2(.05) + 22^2(.20) + \cdots + 26^2(.05) = 549.1$. Therefore $E(N) = 8\pi(549.1) = 13,800.388$.

3.31

- a. The mean of W will be larger than the mean of Y if $\mu > 0$. If $\mu < 0$, the mean of W will be smaller than μ . If $\mu = 0$, the mean of W will equal μ .
- b. $E(W) = E(2Y) = 2E(Y) = 2\mu$.
- c. The variance of W will be larger than σ^2 , since the spread of values of W has increased.
- d. $V(X) = E[(X - E(X))^2] = E[(2Y - 2\mu)^2] = 4E[(Y - \mu)^2] = 4\sigma^2$.

3.37

- a. Not a binomial random variable.
- b. Not a binomial random variable.
- c. Binomial with $n = 100$, $p =$ proportion of high school students who scored above 1026.
- d. Not a binomial random variable (not discrete).
- e. Not binomial, since the sample was not selected among all female HS grads.

3.39

Let $Y =$ number of components failing in less than 1000 hours. Then $Y \sim \text{Binomial}(n = 4, p = .2)$.

- a. $P(Y = 2) = \binom{4}{2}(.2)^2(.8)^2 = 0.1536$.
- b. The system operates if 0, 1, or 2 components fail. $P(\text{system operates}) = .4096 + .4096 + .1536 = .9728$.

3.51

- a. $P(\text{at least one 6 in four rolls}) = 1 - P(\text{no 6's in four rolls}) = 1 - (5/6)^4 = 0.51775$.
- b. In a single toss of two dice, $P(\text{double 6}) = 1/36$. Then $P(\text{at least one double 6 in 24 rolls}) = 1 - P(\text{no double 6's in 24 rolls}) = 1 - (35/36)^{24} = 0.4914$.

3.55

We use the identity

$$\begin{aligned} E[Y(Y-1)(Y-2)] &= \sum_{y=0}^n \frac{y(y-1)(y-2)n!}{y!(n-y)!} p^y (1-p)^{n-y} \\ &= \sum_{y=3}^n \frac{n(n-1)(n-2)(n-3)!}{(y-3)!(n-3-(y-3))!} p^y (1-p)^{n-y} \\ &= n(n-1)(n-2)p^3 \sum_{z=0}^{n-3} \binom{n-3}{z} p^z (1-p)^{n-3-z} \\ &= n(n-1)(n-2)p^3 \end{aligned}$$

Equating this to $E(Y^3) - 3E(Y^2) + 2E(Y)$, it is found that $E(Y^3) = 3n(n-1)p^2 - n(n-1)(n-2)p^3 + np$.

3.59

If Y = number of defective motors, then $Y \sim \text{Binomial}(n = 10, p = .08)$. Then $E(Y) = .8$. The seller's expected net gain is $1000 - 200E(Y) = 840$.

Submitted Problems

3.2

We know $P(HH) = P(TT) = P(HT) = P(TH) = 0.25$. So $P(Y = -1) = 0.5$, $P(Y = 1) = 0.25$, $P(Y = 2) = 0.25$.

3.6

There are 10 sample points, all equally likely: $(1, 2), (1, 3), (1, 4), (1, 5), (2, 3), (2, 4), (2, 5), (3, 4), (3, 5), (4, 5)$.

a. $p(2) = 0.1, p(3) = 0.2, p(4) = 0.3, p(5) = 0.4$. b. $p(3) = 0.1, p(4) = 0.1, p(5) = 0.2, p(6) = 0.2, p(7) = 0.2, p(8) = 0.1, p(9) = 0.1$.

3.10

Let R = rental on a given day, N = no rental. Thus the sequence of interest is $RR, RNR, RNNR, RNNNR, \dots$. Consider the position immediately following the first R : it is filled by an R with prob 0.2, N with prob 0.8. Thus, $P(Y = 0) = 0.2, P(Y = 1) = 0.8(0.2) = 0.16, P(Y = 2) = 0.128$, etc. In general, $P(Y = y) = 0.2(0.8)^y, y = 0, 1, 2, \dots$

3.12

$$E(Y) = 1(.4) + 2(.3) + 3(.2) + 4(.1) = 2.0$$

$$E(1/Y) = 1(.4) + \frac{1}{2}(.3) + \frac{1}{3}(.2) + \frac{1}{4}(.1) = 0.6417$$

$$E(Y^2 - 1) = E(Y^2) - 1 = [1^2(.4) + 2^2(.3) + 3^2(.2) + 4^2(.1)] - 1 = 5 - 1 = 4$$

$$V(Y) = E(Y^2) - [E(Y)]^2 = 5 - 2^2 = 1$$

3.20

With probability 0.3 the volume is $8 \cdot 10 \cdot 30 = 2400$. With probability 0.7 the volume is $8 \cdot 10 \cdot 40 = 3200$. Mean = $0.3(2400) + 0.7(3200) = 2960$

3.22

$p(y) = P(Y = y) = 1/6$ for $y = 1, 2, \dots, 6$. So $E(Y) = 3.5$ and $V(Y) = 2.9167$.

3.24

Distribution of Y = number of bottles with serious flaws:

y	0	1	2
$p(y)$	0.81	0.18	0.01

Thus $E(Y) = 0(.81) + 1(.18) + 2(.01) = 0.20$ and $V(Y) = 0^2(.81) + 1^2(.18) + 2^2(.01) - (0.20)^2 = 0.18$

3.30

- a. Mean of X is larger than mean of Y .
- b. $E(X) = E(Y + 1) = E(Y) + 1 = \mu + 1$.
- c. Variances of X and Y are the same.
- d. $V(X) = E[(X - E(X))^2] = E[(Y + 1 - \mu - 1)^2] = E[(Y - \mu)^2] = \sigma^2$