

Supporting information for:
Hydration of alkali and halide ions. I. Structure and energetics based on simulations with a polarizable force field

Guillaume Lamoureux^{a)†} and Benoît Roux^{b)*}

^{a)}Département de physique, Université de Montréal,
C.P. 6128, succ. centre-ville, Montréal, Québec H3C 3J7, Canada

^{b)}Department of Biochemistry, Weill Medical College of Cornell University,
1300 York Avenue, New York, NY 10021

[†]Current address: Center for Molecular Modeling, Department of Chemistry, University of Pennsylvania

*Corresponding author: Benoit.Roux@med.cornell.edu

Table 1: Hydration free energy data and computations from the literature

a) Alkali ions

Source ^b	Standard state ^a		Li ⁺	Na ⁺	K ⁺	Rb ⁺	Cs ⁺	Comments
	Gas	Solution						
Extrathermodynamic hypothesis								
Noyes62	1 atm	1 M	-120.8	-97.0	-79.3	-74.2	-66.5	$\Delta F_{\text{el}}^{\circ}$ from Table I, plus $\Delta F_{\text{neut}}^{\circ} = 1.325$ kcal/mol
Marcus86-87	1 MPa	1 M	-116.8	-91.5	-74.5	-68.9	-67.6	
Electrochemistry								
Randles56	1 atm	1 M	-122.1	-98.2	-80.6	-75.5	-67.8	Table 1
Gomer77	1 atm	1 M	-118.1	-90.6	-73.1			$\Delta G_{\text{solv}}^{\circ}$ from Table IV
Cluster measurements								
Klots81	1 atm	1 M	-124.0	-100.1	-82.5	-77.4	-69.7	ΔG° from Table I
Tissandier98	1 atm	1 M	-126.5	-101.3	-84.1	-78.7		$\Delta G_{\text{aq}}^{\circ}$ from Table 3, with $X = -264.0$ kcal/mol
Theory								
Zhan2001	1 atm	1 M	-124.9	-99.7	-82.5	-77.1	-69.7	Derived from the hydration free energy of H ⁺
Asthagiri2003	1 M	1 M	-112.7	-88.7				Column SPC/E from Table II
	1 atm	1 M	-110.8	-86.8				Column SPC/E from Table II, plus 1.9 kcal/mol
Grossfield2003	1 atm	1 M		-89.9	-72.6			Table 4
This work	1 M	1 M	-125.0	-98.5	-81.3	-75.7	-68.4	ΔG from TI with SSBP
	1 atm	1 M	-123.1	-96.6	-79.4	-73.8	-66.5	ΔG from TI with SSBP +1.9 kcal/mol = $\Delta G_{\text{hydr}}^{\text{real}}$

b) Halide ions

Source ^b	Standard state ^a		F ⁻	Cl ⁻	Br ⁻	I ⁻	Comments
	Gas	Solution					
Extrathermodynamic hypothesis							
Noyes62	1 atm	1 M	-88.2	-74.8	-67.9	-59.0	$\Delta F_{\text{el}}^{\circ}$ from Table I, plus $\Delta F_{\text{neut}}^{\circ} = 1.325$ kcal/mol
Marcus86-87	1 MPa	1 M	-112.1	-82.4	-76.1	-67.0	
Electrochemistry							
Randles56	1 atm	1 M	-99.1	-70.7	-64.9	-57.2	Table 2
Gomer77	1 atm	1 M	-110.7	-81.4	-76.1		$\Delta G_{\text{solv}}^{\circ}$ from Table IV
Cluster measurements							
Klots81	1 atm	1 M	-101.9	-73.9	-70.6	-59.5	ΔG° from Table I
Tissandier98	1 atm	1 M	-102.5	-72.7	-66.3	-57.4	$\Delta G_{\text{aq}}^{\circ}$ from Table 3, with $X = -264.0$ kcal/mol
Theory							
Zhan2001	1 atm	1 M	-104.1	-74.3	-67.9	-59.0	Derived from the hydration free energy of H ⁺
Zhan2004	1 atm	1 M	-104.3				
Grossfield2003	1 atm	1 M		-84.6			Table 4
This work	1 M	1 M	-108.7	-79.1	-72.6	-64.0	ΔG from TI with SSBP
	1 atm	1 M	-106.8	-77.2	-70.7	-62.1	ΔG from TI with SSBP + 1.9 kcal/mol = $\Delta G_{\text{hydr}}^{\text{real}}$

^aFree energies in the (1 M, 1 M) standard state are converted to the (1 atm, 1 M) standard state by adding 1.9 kcal/mol, the entropic contribution associated with confining 1 mol of ions from a volume of 24.465 ℓ to a volume of 1 ℓ .

^bSource for all values: Noyes62,¹ Marcus86-87,^{2,3,4} Randles56,⁵ Gomer77,⁶ Klots81,⁷ Tissandier98,⁸ Zhan2001,⁹ Zhan2004,¹⁰ Asthagiri2003,¹¹ Grossfield2003.¹²

References

- [1] Richard M. Noyes. Thermodynamics of ion hydration as a measure of effective dielectric properties of water. *J. Am. Chem. Soc.*, 84(4):513–522, 1962.
- [2] Yizhak Marcus. The hydration entropies of ions and their effects on the structure of water. *J. Chem. Soc., Faraday Trans. 1*, 82(1):233–242, 1986.
- [3] Yizhak Marcus. The thermodynamics of solvation of ions. Part 2.—The enthalpy of hydration at 298.15 k. *J. Chem. Soc., Faraday Trans. 1*, 83(2):339–349, 1987.
- [4] Yizhak Marcus. The thermodynamics of solvation of ions. Part 4.—Application of the tetraphenylarsonium tetraphenylborate (TATB) extrathermodynamic assumption to the hydration of ions and to properties of hydrated ions. *J. Chem. Soc., Faraday Trans. 1*, 83(9), 1987.
- [5] J. E. B. Randles. The real hydration energies of ions. *Trans. Faraday Soc.*, 52:1573–1581, 1956.
- [6] R. Gomer and G. Tryson. An experimental determination of absolute half-cell emf’s and single ion free energies of solvation. *J. Chem. Phys.*, 66(10):4413–4424, 1977.
- [7] Cornelius E. Klotz. Solubility of protons in water. *J. Phys. Chem.*, 85(24):3585–3588, 1981.
- [8] Michael D. Tissandier, Kenneth A. Cowen, Wan Yong Feng, Ellen Gundlach, Michael H. Cohen, Alan D. Earhart, James V. Coe, and Thomas R. Tuttle, Jr. The proton’s absolute aqueous enthalpy and Gibbs free energy of solvation from cluster-ion solvation data. *J. Phys. Chem. A*, 102(40):7787–7794, 1998.
- [9] Chang-Guo Zhan and David A. Dixon. Absolute hydration free energy of the proton from first-principles electronic structure calculations. *J. Phys. Chem. A*, 105(51):11534–11540, 2001.
- [10] Chang-Guo Zhan and David A. Dixon. Hydration of the fluoride anion: Structure and absolute hydration free energy from first-principles electronic structure calculations. *J. Phys. Chem. A*, 108(11):2020–2029, 2004.
- [11] D. Asthagiri, Lawrence R. Pratt, and H. S. Ashbaugh. Absolute hydration free energies of ions, ion–water clusters, and quasichemical theory. *J. Chem. Phys.*, 119(5):2702–2708, 2003.
- [12] Alan Grossfield, Pengyu Ren, and Jay W. Ponder. Ion solvation thermodynamics from simulation with a polarizable force field. *J. Am. Chem. Soc.*, 125(50):15671–15682, 2003.