


Supplemental Data

Coordinate File in PDB Format

The file kcsa_inac.pdb contains the atomic coordinates of the non-conducting conformational state of broken symmetry that is shown in Fig. 1d in the article. Ions are present in the binding site S_1 and S_3 ; there is an additional ion in the cavity. Coordinates will be deposited in the PDB.


Quick Time Animation

Illustration of the complete transition from the symmetric ion-conducting conformation to the non-conducting conformation of broken symmetry. A targeted MD trajectory was generated using a RMSD energy restraint applied to all non-hydrogen atoms of residues 72 to 77 (Young et al., 2001, Yang et al., 2002). The full KcsA system (channel, solvent and lipid bilayer) was simulated for 10 ps with a force constant of 500 kcal/mol/Å².



Supplemental Figure S1. Backbone Dihedral Angles in the Selectivity Filter

Φ - Ψ map of the Val76-Gly77 amide plane. The MD data were extracted from a simulation of 1.5 ns in which a backbone transition occurred (Bernèche and Roux, 2000). The conformations of the amide plane corresponding to the conducting and non-conducting states are respectively identified by the letter A and B. The conformation of these residues as observed in the X-ray structure determined in presence of 2 mM KCl and 148 mM NaCl (pdb id 1K4D) (Zhou et al., 2001) is also plotted. The structure corresponds only to a slight deviation from the high K conducting structure (pdb id 1K4C) (Zhou et al., 2001).



Supplemental Figure S2. Hydrogen Bonding Network Surrounding the Selectivity Filter

View from the extra-cellular side showing the inter-subunit interactions involving residues Tyr78-Trp68 and Asp80-Arg89.

References

- Bernèche, S., and Roux, B. Molecular dynamics of the KcsA K⁺ channel in a bilayer membrane (2000). *Biophys. J.* **78**, 2900-2917
- Yang, L., Beard, W., Wilson, S., Roux, B., Broyde, S., and Schlick, T. (2002). Local deformations revealed by dynamics simulations of DNA polymerase Beta with DNA mismatches at the primer terminus. *J Mol Biol.* **321** 459-478
- Young, M.A., Gonfloni, S., Superti-Furga, G., Roux, B., and Kuriyan, J. (2001) Dynamic Coupling Between the SH2 and SH3 Domains of c-Rc and Hck Underlies Their Inactivation by C-terminal Tyrosine Phosphorylation, *Cell* **105**, 115-126
- Zhou, Y., Morais-Cabral, J. H., Kaufman, A. & MacKinnon, R. Chemistry of ion coordination and hydration revealed by a K⁺ channel-Fab complex at 2.0 Å resolution (2001). *Nature* **414**, 43-48