

PSYCHIATRIC MORBIDITY AND ASSOCIATED SOCIO DEMOGRAPHIC FACTORS IN YOUNG UNEMPLOYED PEOPLE

Pakistan Armed Forces Medical Journal

September 30, 2016 Friday

Copyright 2016 PAFMJ All Rights Reserved

Section: No. 4

Length: 3070 words

Byline: Usama Bin Zubair, Sawera Mansoor, Mowaddat Hussain Rana, Usman Ali, Abdul Raafeh Jamal and Huzaifa Qureshi

Body

Abstract

Objective: To determine the prevalence of psychiatric morbidity in young unemployed people and analyze associated socio demographic factors.

Study Design: Cross sectional study.

Place and Duration of Study: Enrollment camps at Punjab and Kashmir, from Jan 2014 to March 2014.

Material and Methods: The sample population comprised of unemployed people who came for recruitment in Pakistan Army at different parts of the country. General Health Questionnaire 12 (GHQ12) was used to screen the population for psychiatric morbidity, 3 was used as cut off score. Age, province, education, level of family income, tobacco smoking, naswar (a tobacco based substance) use, marital status, dependent family members, worrying about the future and social support status were correlated with high GHQ score. Descriptive statistics were used to describe the characteristics of participants and the distribution of GHQ score.

Result: Out of 2511 people 1887 (75.1%) had GHQ score more than 3 showing some psychiatric morbidity. A total of 696 were Kashmiris out of which 540 (77.5%) were GHQ positive, 1329 were Punjabis out of which 978 (73.5%) were GHQ positive, 339 were Pakhtoons out of which 258 (76.1%) were GHQ positive, 129 were Balti out of which 102 (79%) were GHQ positive and 18 were from other ethnicities out of which 9 (50%) were GHQ positive. With logistic regression we found that family size, smoking, naswar use, family income, family history of psychiatric disorder, lack of social support, and low education were significantly correlated with high GHQ score while age of individual, worry about future and marital status had no significant relation with high GHQ score among unemployed people in our study.

Conclusion: This study showed a high prevalence of psychiatric morbidity among unemployed youth of our country. Special attention should be paid to poor people, people with low education and large families. Adequate social support should be provided to young people with no jobs and use of cigarette and naswar should be discouraged.

Keywords: Prevalence, Psychiatric morbidity, Socio demographic factors, Unemployment.

INTRODUCTION

Pakistan is a developing country with an unstable political and social environment. Low literacy rate, poverty, inadequate health facilities and unemployment are basic challenges which our country is facing nowadays. A total

PSYCHIATRIC MORBIDITY AND ASSOCIATED SOCIO DEMOGRAPHIC FACTORS IN YOUNG UNEMPLOYED PEOPLE

of 55.6% of our population is below the age of 25 out of which 21.6% are between the age of 15 and 24 who need sources for employment in near future¹.

Employment status has been found to be related to psychiatric symptoms. A study done in US suggests that the impetus for unemployment be it voluntary or involuntary, may significantly impact a person's mental health². In the few studies that focused on employment status the results consistently show that employed participants were less depressed than unemployed subjects^{3,4}. A study done in France revealed that unemployed men are found to have significantly higher prevalence of depression than the working population⁵. A study done in our neighboring country India showed that unemployment is a risk factor for psychological symptoms among alcohol dependent people⁶.

A study done in a big city of our own country also revealed that unemployment is a risk factor for anxiety and depression⁷.

Many studies have reported a correlation with psychiatric problems and such factors as smoking⁸⁻⁹, drinking¹⁰, family structure^{11,12}, parental relationships^{13,14}, family income¹⁵ and family history of depression¹⁶.

Table-1: Characteristics of the study group and their General Health Questionnaire 12 scores.

Socio demographic factors	No psychiatric morbidity (0-3)		Psychiatric morbidity (score >3)		p-value
	N	%	N	%	
Total	624	24.9	1887	75.1	
Age					
less than 20	234	37.5%	669	35.4%	0.356
20-23	390	62.5%	1218	64.6%	
Education					
10 or less	513	82.2%	1653	87.5%	0.001
>10	111	17.8%	234	12.5%	
Marital status					
Married	45	7.2%	150	7.9%	0.551
Un married	579	92.8%	1737	92.1%	
Family income					
less than Rs.12000	423	67.7%	1656	87.76%	
Rs.12000 or more	201	32.3%	231	12.24%	less
than 0.001					
Tobacco smoking					
Smoker	3	0.48	54	2.86%	less
than 0.001					
Non smoker	621	99.52	1833	97.14%	
Naswar use					
User	18	2.9%	123	6.5%	less
than 0.001					
Non user	606	97.1%	1764	93.5%	
Social support					
Adequate	246	39.4%	306	16.2%	less
than 0.001					
Inadequate	378	60.6%	1581	83.8%	
Worry about future					
Yes	495	79.3%	1641	86.96%	less
than 0.001					
No	129	20.7%	246	13.04%	
Family size					
less than 5 members	117	18.7%	252	13.3%	
5 or more members	507	81.3%	1635	86.7%	0.001
Family hx of psychiatric disorder					
Yes	6	0.96%	123	6.5%	less
than 0.001					
No	618	99.04%	1764	93.5%	

PSYCHIATRIC MORBIDITY AND ASSOCIATED SOCIO DEMOGRAPHIC FACTORS IN YOUNG UNEMPLOYED PEOPLE

Stressors are also found to be correlated with psychiatric illness. They include the military¹⁷, family¹⁸, love relationship¹⁹, finances²⁰, worrying about the future¹⁸, poor coping skills²¹⁻²⁴ and inadequate social support²⁵⁻²⁷.

There is evidence to support a link between unemployment and lower levels of psychological well-being, but debate continues as to whether unemployment results in psychological morbidity, or whether the association is due to those who are more vulnerable to mental illness becoming unemployed.

Here we assess the effect of unemployment in young men on the risk of developing psychiatric illness and to further evaluate and analyze factors associated with psychiatric morbidity.

MATERIAL AND METHODS

This cross sectional study was planned from January 2014 to March 2014. All the applicants who came for enrollment and met the inclusion criteria were included in the study. Males of age greater than 18 years who were unemployed and came for enrollment in Pakistan Army and gave written informed consent were included in the study.

Table-2: The correlated factors relating to psychiatric morbidity: binary logistic regression.

	B	p-value	Odds ratio	95% C.I for EXP(B)	
				Lower	Upper
Education (reference is higher than matriculate)	.381	.005	1.467	1.121	1.910
Marriage (reference is unmarried)	.214	.266	1.238	.83150	1.804
Family size (reference is less than 5 members)	.383	.004	1.467	1.131	1.903
Family income (reference is low income)	1.118	.000	3.059	2.425	3.859
Smoking (reference is no smoking)	1.667	.007	5.295	1.580	17.739
Naswar use (reference is no use of naswar)	.701	.010	2.016	1.186	3.427
Family history of psychiatric disorder (reference is negative family hx)	2.015	.000	7.504	3.197	17.616
Worry about future (reference is not worried)	.260	.049	1.297	1.001	1.681
Social support (reference is adequate support)	1.134	0.000	3.109	2.511	3.849

Non consenting applicants or applicants with past or current history of any psychiatric illness or with past or current psychoactive substance use or those who were unable to understand/complete the required questionnaires were also excluded.

Non probability convenient sampling was done. After the application of inclusion and exclusion criteria, 2511 subjects were included in the analyses.

Different cultures use various methods or screening questionnaires for assessment of mental disorder or psychiatric morbidity. We used validated Urdu version of General Health Questionnaire 12 (GHQ 12)^{28,29}. It is a standardized psychometric test for assessing the general health status of individuals and is used as a screening test. It is 12-item rating screening instrument. The cut off score is greater than 3 by Likert scoring. Severe psychiatric morbidity is suspected above score of 15.

PSYCHIATRIC MORBIDITY AND ASSOCIATED SOCIO DEMOGRAPHIC FACTORS IN YOUNG UNEMPLOYED PEOPLE

The sample was drawn from unemployed applicants who came to get enrolled in Pakistan Army at enrollment camps at different areas of Pakistan. The applicants were gathered in a quiet field with complete reassurance of confidentiality. The subjects were provided with a detailed description of the study and were inducted into the study after written informed consent. The socio demographic data of the full sample of subjects participating in the research was entered in a structured Proforma. The confounding variables were taken care of by detailed history taking about any current or previous psychiatric illness and any current or previous evidence of illicit substance/drug use. Those subjects with confounding variables were excluded from the study. GHQ 12 was administered to the subjects and 3 was taken as cut off value.

Descriptive statistics were used to describe the characteristics of participants and the distribution of GHQ score. Between-group variances in categorical correlates were determined using chi-square. Binary logistic regression analysis was done to evaluate factors related to depressive symptoms.

All statistical analysis was performed using Statistics Package for Social Sciences version 20.0. Differences between groups were considered significant if p-values were less than 0.05

RESULTS

Out of 2511 people 1887 (75.1%) had GHQ score more than 3 showing some psychiatric morbidity. A total of 696 were Kashmiris out of which 540 (77.5%) were GHQ positive, 1329 were Punjabis out of which 978 (73.5%) were GHQ positive, 339 were Pakhtoons out of which 258(76.1%) were GHQ positive, 129 were Bulti out of which 102(79%) were GHQ positive and 18 were from other ethnicities out of which 9(50%) were GHQ positive. With logistic regression we found that family size, smoking, naswar use, family income, family history of psychiatric disorder, lack of social support and low education were significantly correlated with high GHQ score

DISCUSSION

Our study is unique in a sense that it helps in the understanding of psychiatric morbidity among young people who are unemployed and struggling to survive in a competitive environment where making both ends meet is a problem. Using GHQ 12 we found that more than 75% of our sample population showed psychiatric morbidity. This is similar to other studies and surveys showing mental health issues among unemployed people^{30,31}. We used a screening tool so result may show a higher reflection and needs some diagnostic tool to ascertain the problem among the positive individuals. Reason for high frequency may also be inflation and political instability in the country.

Low education was found significantly correlated with psychiatric morbidity in our sample similar to the studies done in the past⁷.

Large family size and low family income is also associated with psychiatric morbidity in accordance with foreign literature^{3,7}. Reason is very clear as large family with less income needs more earning hands to earn adequate to meet the needs. Lack of social support is also found associated with psychiatric morbidity in accordance with the studies done in the past²⁻⁴. Smoking and naswar use also have a strong link with psychiatric morbidity in our study sample. Use of these may give a temporary relief but in long term they have adverse effect on mental health. A study done in France showed the similar results⁵.

Our study has few limitations as well. Randomized selection of study subjects from all the unemployed people was not done. Therefore, the results of the present study cannot be generalized. We used the cross-sectional study method.

CONCLUSION

This study showed a high prevalence of psychiatric morbidity among unemployed youth of our country.

CONFLICT OF INTEREST

This study has no conflict of interest to declare by any author.

PSYCHIATRIC MORBIDITY AND ASSOCIATED SOCIO DEMOGRAPHIC FACTORS IN YOUNG UNEMPLOYED PEOPLE

REFERENCES

1. CIA World Factbook. Unless otherwise noted, information in this page is accurate as of February 21, 2013
2. Pharr JR, Moonie S, Bungum TJ. The Impact of Unemployment on Mental and Physical Health, Access to Health Care and Health Risk Behaviors. ISRN Public Health Volume 2012; (2012).
3. Libby A. M, Ghushchyan, V, McQueen, R. B, and Campbell, J.D: Economic Grand Rounds: Psychological Distress and Depression Associated With Job Loss and Gain: The Social Costs of Job Instability. Psychiatric Services, 2010; VOL. 61, No. 12.
4. Makri-Botsari, E. and Makri, E. (2003). Maternal employment: Effects on her mental health and children's functional status. Psychological Studies, 2005; 48(1), 36-46.
5. Khlait M, Sermet C, Le Pape A. Increased prevalence of depression, smoking, heavy drinking and use of psychoactive drugs among unemployed men in France. Eur J Epidemiol. 2004; 19(5): 445-51.
6. Aswal S, Verma KK, Mathur A, Singh H, Lokesh Jain, Thalor Kapur. Study of psychiatric morbidity and psychosexual dysfunctions in patients of alcohol dependence. Delhi Psychiatry Journal Vol. 15 No.2.
7. Jadoon NA, Munir W, Choudhry ZS. risk factors and prevalence of anxiety and depression in urban multan. Nishtar Medical Journal. 2009; Vol 1, No 2.
8. Keuthen NJ, Niaura RS, Borrelli B: Comorbidity, smoking behavior and treatment outcome. Psychother Psychosom 2000; 69: 244-50.
9. Acton GS, Prochaska JJ, Kaplan AS, Small T, Hall SM: Depression and stages of change for smoking in psychiatric outpatients. Addict Behav 2001; 26: 621-31.
10. Fergusson DM, Boden JM, Horwood LJ: Tests of causal links between alcohol abuse or dependence and major depression. Arch Gen Psychiatry 2009; 66: 260-6.
11. Barrett AE, Turner RJ: Family structure and mental health: the mediating effects of socioeconomic status, family process, and social stress. J Health Soc Behav 2005; 46: 156-69.
12. Patten CA, Gillin JC, Farkas AJ, Gilpin EA, Berry CC, Pierce JP: Depressive symptoms in California adolescents: family structure and parental support. J Adolesc Health 1997; 20: 271-8.
13. Gilman SE, Kawachi I, Fitzmaurice GM, Buka SL: Family disruption in childhood and risk of adult depression: Am J Psychiatry 2003; 160: 939-46.
14. Stivers C: Parent-adolescent communication and its relationship to adolescent depression and suicide proneness. Adolescence 1988; 23: 291-5.
15. Whooley MA, Kiefe CI, Chesney MA, Markovitz JH, Matthews K, Hulley SB: Depressive symptoms, unemployment, and loss of income: The **CARDIA study**. Arch Intern Med 2002; 162: 2614-20.
16. Biederman J, Faraone SV, Hirshfeld-Becker DR, Friedman D, Robin JA, Rosenbaum JF: Patterns of psychopathology and dysfunction in high-risk children of parents with panic disorder and major depression. Am J Psychiatry 2001; 158: 49-57.
17. Pflanz SE, Ogle AD: Job stress, depression, work performance, and perceptions of supervisors in military personnel. Mil Med 2006; 171: 861-5.
18. Phelan J, Schwartz JE, Bromet EJ, Work stress, family stress and depression in professional and managerial employees. Psychol Med 1991; 21: 999-1012.

PSYCHIATRIC MORBIDITY AND ASSOCIATED SOCIO DEMOGRAPHIC FACTORS IN YOUNG UNEMPLOYED PEOPLE

19. Maltby J, Day L: Romantic acts and depression. *Psychol Rep* 2000; 86: 260-2.
20. Eisenberg L: Psychiatry and health in low-income populations. *Compr Psychiatry* 1997; 38: 69-73.
21. Cronkite RC, Moos RH, Twohey J, Cohen C, Swindle R Jr: Life circumstances and personal resources as predictors of the ten- year course of depression. *Am JCommunity Psychol* 1998; 26: 255-80.
22. Goodwin RD: Association between coping with anger and feelings of depression among youths. *Am J Public Health* 006; 96: 664-9.
23. Vitaliano PP, DeWolfe DJ, Maiuro RD, Russo J, Katon W: Appraised changeability of stressor as a modifier of the relationship between coping and depression: a test of the hypothesis of fit. *J Pers Soc Psychol* 1990; 59: 585-92.
24. Powers DV, Gallagher-Thompson D, Kraemer HC: Coping and depression in Alzheimer's caregivers: longitudinal evidence of stability. *J Gerontol BPsychol Sci Soc Sci* 2002; 57: 205-11.
25. Clovis Alexandrino-Silva, Tania Ferraz Alves, Luis Fernando Tofoli, Yuan-Pang Wang, Laura Helena Andrade. Psychiatry - life events and social support in late life depression. *Clinics (Sao Paulo)*. 2011; 66(2): 233-238.
26. Zubair UB, Mansoor S, Rana MH. Prevalence of depressive symptoms and associated socio-demographic factors among recruits duringmilitary training. *J R Army Med Corps* 2014; 0:1-5.
27. Kouvonen A, Roberto De Vogli, Stafford M. Social support and the likelihood of maintaining and improving levels of physical activity: the Whitehall II Study. *Eur J Public Health*. 2012; 22(4): 514-518.
28. Lesage FX, Martens-Resende S, Deschamps F, Berjot S. Validation of the General Health Questionnaire (GHQ-12) adapted to a work-related context. *OJPM> Vol.1 No.2*, 44-48 2011.
29. Khan A, Shah IM, Khan F, Suhail S. Reliability and Validity Assessment of 12 Items General Health Questionnaire (GHQ: 12) among Pakistani University Teachers. *World Applied Sciences Journal* 24 (5): 603-608, 2013.
30. OECD 2012. Sick on the Job?Myths and Realities about Mental Health and Work, OECD Publishing.

Classification

Language: ENGLISH

Publication-Type: Magazine

Subject: DEMOGRAPHIC GROUPS (90%); UNEMPLOYED PERSONS (90%); POPULATION & DEMOGRAPHICS (90%); SMOKING (89%); UNEMPLOYMENT RATES (89%); POVERTY & HOMELESSNESS (78%); MENTAL HEALTH (78%); MILITARY DEPENDENTS (76%); LITERACY & ILLITERACY (73%); ARMIES (72%); LITERACY RATES (69%); RACE & ETHNICITY (65%); DISEASES & DISORDERS (64%); MENTAL ILLNESS (64%)

PSYCHIATRIC MORBIDITY AND ASSOCIATED SOCIO DEMOGRAPHIC FACTORS IN YOUNG
UNEMPLOYED PEOPLE

Industry: MILITARY DEPENDENTS (76%); TOBACCO PRODUCTS (75%); ARMIES (72%)

Geographic: KASHMIR (88%); PAKISTAN (92%); FRANCE (50%); Pakistan

Load-Date: September 20, 2016

End of Document