

1.

a.

Let $x, y \in U$ such that $f(x) = f(y)$ and $\xi = y - x$. From the Taylor's theorem and the fact that $x + \theta\xi$ is in the convex set U for $\theta \in [0, 1]$, we have that for each $j = 1, \dots, N$, there exists a number $\theta_j \in [0, 1]$ such that

$$f_j(y) = f_j(x + \xi) = f_j(x) + \sum_{k=1}^N \frac{\partial f_j}{\partial x_k}(x + \theta_j \xi) \xi_k = f_j(x)$$

It follows that

$$\sum_{k=1}^N \frac{\partial f_j}{\partial x_k}(x + \theta_j \xi) \xi_k = 0$$

for $j = 1, \dots, N$. Let

$$A := \begin{bmatrix} \frac{\partial f_1}{\partial x_1}(x + \theta_1 \xi) & \dots & \frac{\partial f_1}{\partial x_N}(x + \theta_1 \xi) \\ \vdots & \ddots & \vdots \\ \frac{\partial f_N}{\partial x_1}(x + \theta_N \xi) & \dots & \frac{\partial f_N}{\partial x_N}(x + \theta_N \xi) \end{bmatrix}$$

so that $A\xi = 0$. However, since the points in the set $\{x + \theta\xi | \theta \in [0, 1]\}$ are collinear points. We have that $\det(A) \neq 0$ and therefore $\xi = 0$ which means that $x = y$.

b.

For any $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, we have that

$$\det \begin{bmatrix} \frac{\partial f_1}{\partial x_1}(x_1, y_1) & \frac{\partial f_1}{\partial x_2}(x_1, y_1) \\ \frac{\partial f_2}{\partial x_1}(x_1, y_1) & \frac{\partial f_2}{\partial x_2}(x_1, y_1) \end{bmatrix} = \det \begin{bmatrix} 3x_1^2 & -1 \\ e^{x_1+y_1} & e^{x_2+y_2} \end{bmatrix} = 3x_1^2 \cdot e^{x_2+y_2} - e^{x_1+y_1}$$

which is > 0 for all $(x_1, y_1), (x_2, y_2)$ and hence f is injective.

2.

a.

$$\begin{aligned} \det J_f(x, y) &= \det \begin{bmatrix} \frac{\sqrt{x^2 + y^2} - x \cdot \frac{2x}{2\sqrt{x^2 + y^2}}}{x^2 + y^2} & \frac{-xy}{(x^2 + y^2)^{3/2}} \\ \frac{-xy}{x^2 + y^2} & \frac{\sqrt{(x^2 + y^2)^{3/2}} - y \cdot \frac{2y}{2\sqrt{x^2 + y^2}}}{x^2 + y^2} \end{bmatrix} \\ &= \det \begin{bmatrix} \frac{y^2}{(x^2 + y^2)^{3/2}} & \frac{-xy}{(x^2 + y^2)^{3/2}} \\ \frac{-xy}{(x^2 + y^2)^{3/2}} & \frac{x^2}{(x^2 + y^2)^{3/2}} \end{bmatrix} = 0 \end{aligned}$$

b.

For all (x, y) , we have that

$$|f(x, y)| = \sqrt{\frac{x^2}{x^2 + y^2} + \frac{y^2}{x^2 + y^2}} = 1$$

Therefore, $f(U)$ is the circle around the origin with radius 1. It does not contain any non-empty open subset as for all point $(x, y) \in f(U) : \forall \epsilon > 0 : (x + \epsilon, y) \notin f(U)$

3.

Consider

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad x \rightarrow x^3$$

Then $f \in \mathcal{C}^1(U, \mathbb{R}^N)$ where $U = \mathbb{R}$ and $N = 1$.

$\det J_f(0) = 0$ but every neighborhood $(-\epsilon, \epsilon)$ is mapped to $(-\epsilon^3, \epsilon^3)$ which is open.

4.

a.

We have that for all x satisfies $f(x) \neq (0, \dots, 0)$,

$$\frac{\partial g}{\partial x_j}(x_0) = \frac{\sum_{j=1}^N 2f_j(x) \frac{\partial f_j(x)}{\partial x_i}}{\sqrt{\sum_{j=1}^N f_j(x)^2}}$$

and hence $\nabla g(x) = \frac{J_f(x)f(x)}{\sqrt{\sum_{j=1}^N f_j(x)^2}}$ which = 0 if and only if $J_f(x)f(x) = 0$

which means that $f(x) = 0$ as $\det J_f(x) \neq 0$ (its inverse exists) and therefore is a contradiction.

At $f(x_0) = (0, \dots, 0)$, $g(x_0) = 0$ and hence it is clearly the global minimum as norm is always ≥ 0 . Therefore, there is no local maximum.

b.

Since \overline{U} is compact $g(\overline{U})$ is also compact and hence $g(\overline{U})$ must attain its maximum since every converges sequence converges to a point in the set and hence the supremum is the maximum.

Since \tilde{f} does not attain its local maximum and hence global maximum on $\text{int } \overline{U}$, the global maximum must be attained on the boundary $\partial U = \partial \overline{U}$.