

MATH 217 (Fall 2022)
Honors Advanced Calculus, I

Assignment #6

1. Determine the Jacobians of

$$\mathbb{R}^3 \rightarrow \mathbb{R}^3, \quad (r, \theta, \phi) \mapsto (r \sin \theta \cos \phi, r \sin \theta \sin \phi, r \cos \theta)$$

and

$$\mathbb{R}^3 \rightarrow \mathbb{R}^3, \quad (r, \theta, z) \mapsto (r \cos \theta, r \sin \theta, z).$$

2. An $N \times N$ matrix X is *invertible* if there is $X^{-1} \in M_N(\mathbb{R})$ such that $XX^{-1} = X^{-1}X = I_N$ where I_N denotes the unit matrix.

(a) Show that $U := \{X \in M_N(\mathbb{R}) : X \text{ is invertible}\}$ is open. (*Hint:* $X \in M_N(\mathbb{R})$ is invertible if and only if $\det X \neq 0$.)

(b) Show that the map

$$f: U \rightarrow M_N(\mathbb{R}), \quad X \mapsto X^{-1}$$

is totally differentiable on U , and calculate $Df(X_0)$ for each $X_0 \in U$. (*Hint:* You may use that, by Cramer's Rule, f is continuous.)

3. Let

$$p: (\mathbb{R} \setminus \{0\}) \times \mathbb{R} \rightarrow \mathbb{R}^2, \quad (r, \theta) \mapsto (r \cos \theta, r \sin \theta),$$

let $\emptyset \neq U \subset \mathbb{R}^2$ be open, and let $f: U \rightarrow \mathbb{R}$ be twice continuously partially differentiable. Show that

$$(\Delta f) \circ p = \frac{\partial^2(f \circ p)}{\partial r^2} + \frac{1}{r} \frac{\partial(f \circ p)}{\partial r} + \frac{1}{r^2} \frac{\partial^2(f \circ p)}{\partial \theta^2}$$

on $p^{-1}(U)$. (*Hint:* Apply the Chain Rule twice.)

4. Let

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}, \quad (x, y) \mapsto \begin{cases} \frac{xy^3}{x^2+y^4}, & (x, y) \neq (0, 0), \\ 0, & (x, y) = (0, 0). \end{cases}$$

Show that:

- (a) f is continuous at $(0, 0)$;
- (b) for each $v \in \mathbb{R}^2$ with $\|v\| = 1$, the directional derivative $D_v f(0, 0)$ exists and equals 0;
- (c) f is not totally differentiable at $(0, 0)$.

(*Hint for (c)*: Assume towards a contradiction that f is totally differentiable at $(0, 0)$, and compute the first derivative of $\mathbb{R} \ni t \mapsto f(t^2, t)$ at 0 first directly and then using the chain rule. What do you observe?)

5. Let $U \subset \mathbb{R}^N$ be open and convex and contain 0, let $n \in \mathbb{N}_0$, and let $f: \mathbb{R}^N \rightarrow \mathbb{R}$ be $n + 1$ times continuously partially differentiable such that $\frac{\partial^\alpha f}{\partial x^\alpha} \equiv 0$ on U for all multiindices $\alpha \in \mathbb{N}_0^N$ with $|\alpha| = n + 1$. Show that, there are $(c_\alpha)_{\alpha \in \mathbb{N}_0^N, |\alpha| \leq n}$ in \mathbb{R} such that

$$f(x) = \sum_{|\alpha| \leq n} c_\alpha x^\alpha$$

for all $x \in U$.

- 6*. Let $\emptyset \neq C \subset \mathbb{R}^N$ be open and connected, and let $f: C \rightarrow \mathbb{R}$ be differentiable such that $\nabla f \equiv 0$. Show that f is constant. (*Hint*: First, treat the case where C is convex using the chain rule; then, for general C , assume that f is not constant, let $x, y \in C$ such that $f(x) \neq f(y)$, and show that $\{U, V\}$ with $U := \{z \in C : f(z) = f(x)\}$ and $V := \{z \in C : f(z) \neq f(x)\}$ is a disconnection for C .)

Due Monday, October 31, 2020, at 10:00 a.m.; no late assignments.