

Question 1

Correct

Marked out of 3.00

🚩 [Flag question](#)

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[i] - A[j] = k$, $i \neq j$.

Input Format

1. First line is number of test cases T. Following T lines contain:
2. N, followed by N integers of the array
3. The non-negative integer k

Output format

Print 1 if such indices exist and 0 if

Print 1 if such a pair exists and 0 if it doesn't.

Example

Input:

1

3 1 3 5

4

Output:

1

Input:

1

3 1 3 5

99

Output:

0

Answer: (penalty regime: 0 %)

```
1 #include<stdio.h>
2 int main(){
3     int t;
4     scanf("%d"
5 while(t--){
6     int n;
7     scanf("%d"
8     int a[n];
9     for (int i=0; i<n; i++){
10         scanf("%d", &a[i]);
11     }
12     int k;
13     scanf("%d", &k);
14     int flag=0;
15     for(int i=0; i<n; i++){
16         for(int j=i+1; j<n; j++){
17             if(a[i]-a[j]==k || a[j]-a[i]==k){
18                 flag=1; break;
19             }
```

```

20     }
21     }
22     if(flag)break;}
23     printf("%d\n",flag);
24     }
25     }

```

	Input	Expected	Got
✓	1 3 1 3 5 4	1	1
✓	1 3 1 3 5 99	0	0

Passed all tests! ✓

Question **2**

Correct

Marked out of 5.00

Flag question

Question 2

Correct

Marked out of 5.00

🚩 [Flag question](#)

Sam loves chocolates and starts buying them on the 1st day of the year. Each day of the year, x , is numbered from 1 to Y . On days when x is odd, Sam will buy x chocolates; on days when x is even, Sam will not purchase any chocolates.

Complete the code in the editor so that for each day N_i (where $1 \leq x \leq N \leq Y$) in array `arr`, the number of chocolates Sam purchased (during days 1 through N) is printed on a new line. This is a function-only challenge, so input is handled for you by the locked stub code in the editor.

Input Format

The program takes an array of integers as a parameter.

The locked code in the editor handles reading the following input from stdin, assembling it into an array of integers (arr), and calling calculate(arr).

The first line of input contains an integer, T (the number of test cases). Each line i of the T subsequent lines describes the i th test case as an integer, N_i (the number of days).

Constraints

$$1 \leq T \leq 2 \times 10^5$$

$$1 \leq N \leq 2 \times 10^6$$

$$1 \leq x \leq N \leq Y$$

Output Format

For each test case, T_i in arr, your calculate method should print the total number of chocolates Sam purchased by day N_i on a new line.

Sample Input 0

3

1

2

3

Sample Output 0

1

1

Explanation

Test Case 0: $N = 1$

Sam buys 1 chocolate on day 1, giving us a total of 1 chocolate. Thus, we print 1 on a new line.

Test Case 1: $N = 2$

Sam buys 1 chocolate on day 1 and 0 on day 2. This gives us a total of 1 chocolate. Thus, we print 1 on a new line.

Test Case 2: $N = 3$

Sam buys 1 chocolate on day 1, 0 on day 2, and 3 on day 3. This gives us a total of 4 chocolates. Thus, we print 4 on a new line.

Answer: (penalty regime: 0 %)

```
1  #include<stdio.h>
2  ▼ int main(){
3      int t;
4      scanf("%d",
5  ▼     while(t--){
6          int n,c=0;
7          scanf("%d",
8  ▼     for(int i=0;
9         if(i%2!=0)c=
10        }printf("%d
11    }
12    }
```

Question 3

Correct

Marked out of 7.00

 [Flag question](#)

The number of goals achieved by two football teams in matches in a league is given in the form of two lists. Consider:

- Football team A, has played three matches, and has scored $\{ 1, 2, 3 \}$ goals in each match respectively.
- Football team B, has played two matches, and has scored $\{ 2, 4 \}$ goals in each match respectively.
- Your task is to compute, for each match of team B, the total number of matches of team A, where team A has scored less than or equal to the number of

where team A has scored less than or equal to the number of goals scored by team B in that match.

- In the above case:
- For 2 goals scored by team B in its first match, team A has 2 matches with scores 1 and 2.
- For 4 goals scored by team B in its second match, team A has 3 matches with scores 1, 2 and 3.

Hence, the answer: {2, 3}.

Complete the code in the editor below. The program must return an array of m positive integers, one for each $\text{maxes}[i]$ representing the total number of elements $\text{nums}[j]$ satisfying $\text{nums}[j] \leq \text{maxes}[i]$ where $0 \leq j < n$ and $0 \leq i < m$, in the given order.

It has the following:

`nums[nums[0],...nums[n-1]]`: first array of positive integers

`maxes[maxes[0],...maxes[n-1]]`: second array of positive integers

Constraints

- $2 \leq n, m \leq 10^5$
- $1 \leq \text{nums}[j] \leq 10^9$, where $0 \leq j < n$.
- $1 \leq \text{maxes}[i] \leq 10^9$, where $0 \leq i < m$.

Input Format For Custom Testing

Input from stdin will be processed as follows and passed to the function.

The first line contains an integer n ,

The first line contains an integer n , the number of elements in `nums`.

The next n lines each contain an integer describing `nums[j]` where $0 \leq j < n$.

The next line contains an integer m , the number of elements in `maxes`.

The next m lines each contain an integer describing `maxes[i]` where $0 \leq i < m$.

Sample Case 0

Sample Input 0

4

1

4

2

4

2

3

5

Sample Output 0

2

4

Explanation 0

We are given $n = 4$, $\text{nums} = [1, 4, 2, 4]$, $m = 2$, and $\text{maxes} = [3, 5]$.

1. For $\text{maxes}[0] = 3$, we have 2 elements in nums ($\text{nums}[0] = 1$ and $\text{nums}[2] = 2$) that are $\leq \text{maxes}[0]$.

2. For $\text{maxes}[1] = 5$, we have 4 elements in nums ($\text{nums}[0] = 1$, $\text{nums}[1] = 4$, $\text{nums}[2] = 2$, and

2. For $\text{maxes}[1] = 5$, we have 4 elements in nums ($\text{nums}[0] = 1$, $\text{nums}[1] = 4$, $\text{nums}[2] = 2$, and $\text{nums}[3] = 4$) that are $\leq \text{maxes}[1]$.

Thus, the function returns the array $[2, 4]$ as the answer.

Sample Case 1

Sample Input 1

5

2

10

5

4

8

4

3

1

7

8

Sample Output 1

1

0

3

4

Explanation 1

We are given, $n = 5$, $\text{nums} = [2, 10, 5, 4, 8]$, $m = 4$, and $\text{maxes} = [3, 1, 7, 8]$.

1. For $\text{maxes}[0] = 3$, we have 1 element in nums ($\text{nums}[0] = 2$) that is $\leq \text{maxes}[0]$.

2. For $\text{maxes}[1] = 1$, there are 0 elements in nums that are \leq

maxes[1].

3. For maxes[2] = 7, we have 3 elements in nums (nums[0] = 2, nums[2] = 5, and nums[3] = 4) that are \leq maxes[2].

4. For maxes[3] = 8, we have 4 elements in nums (nums[0] = 2, nums[2] = 5, nums[3] = 4, and nums[4] = 8) that are \leq maxes[3].

Thus, the function returns the array [1, 0, 3, 4] as the answer.

Answer: (penalty regime: 0 %)

```
1  #include<stdio.h>
2  int main(){
3  int m,n,co=0;
4  scanf("%d\n",&m);
5  int a[m];
6  for(int i=0;i<m;i++){
7  scanf("%d",&n);
8  int b[n],c[n];
9  for(int j=0;j<n;j++){
10  for(int k=0;k<n;k++){
11  for(int p=0;p<m;p++){
```

Answer: (penalty regime: 0 %)

```
1  #include<stdio.h>
2  ▼ int main(){
3    int m,n,co=0;
4    scanf("%d\n",&m);
5    int a[m];
6    for(int i=0;i<m;i++)
7    scanf("%d",&n);
8    int b[n],c[n];
9    for(int j=0;j<n;j++)
10   for(int k=0;k<n;k++)
11   {for(int p=0;p<m;p++)
12    c[k]=co;
13    co=0;
14   }
15  ▼ for(int i=0;i<n;i++)
16    printf("%d\n",c[i]);
17  }
18  return 0;
19  }
20
```
