

Report 8

Khaled Hasan

2024-05-12

Data

```
GLOBALTEMPERATURE = read.csv(file = "C:\\Users/ss/Desktop/Time_series_Analysis/MyGlobalTemperetures.csv")
global_temp = ts(GLOBALTEMPERATURE[,1], start = c(1850, 1), frequency = 12)
northern_temp = ts(GLOBALTEMPERATURE[,2], start = c(1850, 1), frequency = 12)
southern_temp = ts(GLOBALTEMPERATURE[,3], start = c(1850, 1), frequency = 12)
Regional_temp = read.csv(file = "C:/Users/ss/Desktop/Time_series_Analysis/Regional_temperetures_data.csv")
Europe_temp = ts(Regional_temp[, 3], start = c(1850, 1), frequency = 12)
time = ts(Regional_temp[, 1], start = c(1850, 1), frequency = 12)
```


Plots

```
library(ggplot2)
library(gridExtra)
```


```
## Warning: package 'gridExtra' was built under R version 4.3.3
```

```
autoplot1 <- autoplot(window(global_temp, start = c(2000, 1), freq = 12), ylab = "global")
autoplot2 <- autoplot(window(northern_temp, start = c(2000, 1), freq = 12), ylab = "Northern")
autoplot3 <- autoplot(window(southern_temp, start = c(2000, 1), freq = 12), ylab = "Southern")


par(mfrow = c(1, 3))
plot(autoplot1)
```




```
plot(autoplot2)
```



```
plot(autoplot3)
```



```
layout(matrix(c(1, 2, 3), nrow = 1))  
grid.arrange(autoplot1, autoplot2, autoplot3)
```



Notice that the temperature averages in the southern hemi-sphere are out of phase with those from the northern Hemisphere, and that the global average is dominated by the northern hemisphere (northern hemisphere has more land mass).

plotting p-value

Finding best fitting model:

```
temp_1980_afterwards = window(global_temp, start=c(1980, 1), freq=12)

global_fitting_sarima = auto.arima(window(global_temp, start=c(1980, 1), freq=12), approximation = FALSE)


summary(global_fitting_sarima)
```

```
## Series: window(global_temp, start = c(1980, 1), freq = 12)
## ARIMA(2,0,2)(1,1,1)[12]
##
## Coefficients:
##          ar1      ar2      ma1      ma2      sar1      sma1
##      -0.0283  0.7152  0.2870 -0.4169 -0.0832 -0.8624
## s.e.   0.1128  0.1198  0.1323  0.1336  0.0511  0.0386
##
## sigma^2 = 0.2214: log likelihood = -341.91
## AIC=697.83   AICc=698.05   BIC=727.4
```

```
##
## Training set error measures:
##           ME      RMSE      MAE      MPE      MAPE      MASE
## Training set 0.06138175 0.4623172 0.3587063 0.1622547 5.140828 0.6865682
##           ACF1
## Training set 0.0006093937
```

```
checkresiduals(global_fitting_sarima)
```


Residuals from ARIMA(2,0,2)(1,1,1)[12]



```
##
## Ljung-Box test
##
## data: Residuals from ARIMA(2,0,2)(1,1,1)[12]
## Q* = 15.772, df = 18, p-value = 0.6085
##
## Model df: 6. Total lags used: 24
```

```
global_fitting_sarima %>% forecast(h = 24) %>% autoplot()
```

Forecasts from ARIMA(2,0,2)(1,1,1)[12]



```
#minimum = global_fitting_sarima$aic
#ord = c(2, 0, 2)
#for (i in 0:5) {
#  for (j in 0:5) {
#    tord = c(i, 0, j)
#    if(Arima(temp_1980_afterwards, tord, c(1, 1, 1), include.drift = TRUE)$aic < minimum){
#
#      #minimum = Arima(temp_1980_afterwards, tord, c(1, 1, 1), include.drift = TRUE)$aic
#      #ord = tord
#    }
#  }
#}
#}
```

```
ord = c(2, 0, 0)
model = Arima(temp_1980_afterwards, ord, c(1, 1, 1), include.drift = TRUE)
summary(model)
```


```
## Series: temp_1980_afterwards
## ARIMA(2,0,0)(1,1,1)[12] with drift
##
## Coefficients:
##          ar1      ar2      sar1      sma1      drift
##          0.2349  0.1870  -0.0407  -0.9680  0.0026
## s.e.    0.0437  0.0442   0.0481   0.0406  0.0002
##
```

```
## sigma^2 = 0.2039: log likelihood = -329.29
## AIC=670.57 AICc=670.74 BIC=695.92
##
## Training set error measures:
##           ME      RMSE      MAE      MPE      MAPE      MASE
## Training set -0.01208849 0.4440469 0.3418198 -0.8092102 4.972688 0.6542473
##           ACF1
## Training set 0.00154746
```

Notice that the manually fitted model has a lower aic, a higher loglikelihood and included a drift, that agrees with the trend regression fitting in the following section

```
checkresiduals(model)
```

Residuals from ARIMA(2,0,0)(1,1,1)[12] with drift



```
##
## Ljung-Box test
##
## data: Residuals from ARIMA(2,0,0)(1,1,1)[12] with drift
## Q* = 10.995, df = 20, p-value = 0.9464
##
## Model df: 4. Total lags used: 24
```


Fitting with regression:

```
Arima_fittng <- function(timeseries, startingPoint = start(timeseries), endingPoint = end(timeseries)){  
  cutted_data = window(timeseries, start = startingPoint, end = endingPoint, freq = 12)  
  t = seq_along(cutted_data)  
  regressors = cbind(sin(pi/6*t), cos(pi/6*t), t)  
  arima_fit = auto.arima(cutted_data, xreg = regressors, approximation = FALSE, seasonal = TRUE)  
  return(arima_fit)  
}
```


```
global_fitting_Arimareg = Arima_fittng(global_temp, startingPoint = c(1987, 1))
```

```
summary(global_fitting_Arimareg)
```

```
## Series: cutted_data  
## Regression with ARIMA(2,0,1) errors  
##  
## Coefficients:  
##          ar1      ar2      ma1  intercept              t  
##      -0.5406  0.2813  0.7924     9.3065    -3.1557    -4.5193    0.0026  
## s.e.   0.1670  0.0490  0.1702     0.0623     0.0404     0.0403    0.0003  
##  
## sigma^2 = 0.211: log likelihood = -274.1  
## AIC=564.2   AICc=564.54   BIC=596.77  
##  
## Training set error measures:  
##              ME      RMSE      MAE      MPE      MAPE      MASE  
## Training set -0.0002543913 0.4556364 0.354646 -0.662263 4.962195 0.6954969  
##              ACF1  
## Training set -0.00761437
```

```
checkresiduals(global_fitting_Arimareg)
```

Residuals from Regression with ARIMA(2,0,1) errors



```
##
##  Ljung-Box test
##
## data:  Residuals from Regression with ARIMA(2,0,1) errors
## Q* = 27.979, df = 21, p-value = 0.1408
##
## Model df: 3.    Total lags used: 24
```

doing the same with the data obtained from Europe

```
Europe_fitting_Arimareg = Arima_fitnng(Europe_temp, startingPoint = c(1987, 1))


summary(Europe_fitting_Arimareg)
```

```
## Series: cutted_data
## Regression with ARIMA(1,0,0) errors
##
## Coefficients:
##          ar1  intercept              t
##          0.2246    5.7010   -6.6958   -9.6269   4e-03
## s.e.    0.0469    0.1091    0.0736    0.0735   4e-04
##
## sigma^2 = 0.7838:  log likelihood = -559.17
## AIC=1130.34  AICc=1130.54  BIC=1154.77
##
```

```
## Training set error measures:
##           ME      RMSE      MAE      MPE      MAPE      MASE
## Training set 0.0008772039 0.8801977 0.6430352 -27.01633 62.86381 0.6869426
##           ACF1
## Training set -0.007694757
```

```
checkresiduals(Europe_fitting_Arimareg)
```

Residuals from Regression with ARIMA(1,0,0) errors



```
##
## Ljung-Box test
##
## data: Residuals from Regression with ARIMA(1,0,0) errors
## Q* = 19.403, df = 23, p-value = 0.6776
##
## Model df: 1. Total lags used: 24
```

Plotting global warming coefficients

```
linear_coef <- function(DATA, x, Ord, sOrd, radius = 2){
  temporary_data = window(DATA, start = c(x-radius, 1), end = c(x+radius, 1))
  new_t <- seq_along(temporary_data)
```

```

temporary_xreg = cbind(sin(new_t*pi/6), cos(new_t*pi/6), new_t)

temporary_model = arima(temporary_data, order = Ord, seasonal = sOrd, xreg = temporary_xreg)

std_error <- sqrt(diag(vcov(temporary_model)))

return(c(as.numeric(temporary_model$coef["new_t"]), as.numeric(sqrt(diag(vcov(temporary_model))))["new_t"]))
}

plot_Global_warming <- function(timeseries, arima_fit){

  #arima_fit = Arima_fitting(timeseries, startingPoint = c(1980, 1))

  ord = arima_fit$arma

  p = ord[1]; q = ord[2]; P = ord[3]; Q = ord[4]; period = ord[5]; d = ord[6]; D = ord[7];

  parameters = c()

  errors = c()

  rad = 20

  sp = 1850

  fp = 2023

  for (i in (sp + rad):(fp - rad)){
    u = linear_coef(timeseries, i, c(p, d, q), c(P, D, Q) , rad)
    parameters <- cbind(parameters, u[1])
    errors <- cbind(errors, u[2])
  }

  t_vals = c((sp + rad):(fp - rad))


  MyPlot = (plot(t_vals, parameters, type='b', main=paste("c\\nradius: ", toString(rad)), ylim = c(-0.01, 0.01),
    arrows(x0=t_vals, y0=parameters-2*errors, x1 = t_vals, y1=parameters+2*errors, code=3, angle = 90,
    return(list(arima_fit, parameters, errors, MyPlot))
}

Results_for_global = plot_Global_warming(global_temp, arima_fit = global_fitting_Arimareg)


## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced

```

```
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
```



```
Results_for_Europe = plot_Global_warming(Europe_temp, arima_fit = Europe_fitting_Arimareg)
```



Plotting p-values

using the standard errors and assuming normal distribution (due to the big number of data points)

```
plot_p_vals <- function(timeseries, arima_fit){
  #arima_fit = Arima_fittn(timeseries, startingPoint = c(1980, 1))

  ord = arima_fit$arma

  p = ord[1]; q = ord[2]; P = ord[3]; Q = ord[4]; period = ord[5]; d = ord[6]; D = ord[7];

  parameters = c()

  errors = c()

  p_values = c()

  rad = 20

  sp = 1850

  fp = 2023

  for (i in (sp + rad):(fp - rad)){
```

```

u = linear_coef(timeseries, i, c(p, d, q), c(P, D, Q) , rad)
parameters <- cbind(parameters, u[1])
errors <- cbind(errors, u[2])

p = 2*pnorm(min(0, 2*u[1]), mean = u[1], sd = u[2], lower.tail = TRUE)
p_values = cbind(p_values, p)
}

t_vals = c((sp + rad):(fp - rad))


plot(t_vals, -log10(p_values), type='b', main=paste("c\nradius: ", toString(rad)))
}

plot_p_vals(global_temp, arima_fit = global_fitting_Arimareg)


```

```
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
```

```
## Warning in sqrt(diag(vcov(temporary_model))): NaNs produced
```



```
plot_p_vals(Europe_temp, arima_fit = Europe_fitting_Arimareg)
```



There is no evidence that Europe had more global warming!!!