

COSC6364 Adv. Numerical Analysis

Wu, Panruo

COSC 6364 Adv. Numerical Analysis

Lecture 1: Matrix-Vector Multiplication; Orthogonal Vectors and Matrices

P. Wu¹

¹Computer Science University of Houston

Spring 2020

To accompany the Numerical Linear Algebra book, Lectures 1 and 2



Table of Contents

COSC6364 Adv. Numerical Analysis

Wu, Panruc



COSC6364 Adv. Numerical Analysis

Wu, Panruc



COSC6364 Adv. Numerical Analysis

Wii. Panriio

New interpretation that is essential for numerical linear algebra: if b = Ax, then b is the linear combination of the columns of A.



COSC6364 Adv. Numerical Analysis

Wii. Panriio

- New interpretation that is essential for numerical linear algebra: if b = Ax, then b is the linear combination of the columns of A.
 - Let a_i denote the i^{th} column of A, m-vector. Then we can write



COSC6364 Adv. Numerical Analysis

Wii Panriio

- New interpretation that is essential for numerical linear algebra: if b = Ax, then b is the linear combination of the columns of A.
 - Let a_i denote the i^{th} column of A, m-vector. Then we can write

$$\boldsymbol{b} = X = \sum_{i=1}^{n} x_i \boldsymbol{a}_i$$



COSC6364 Adv. Numerical Analysis

Wii. Panriid

- New interpretation that is essential for numerical linear algebra: if b = Ax, then b is the linear combination of the columns of A.
 - Let a_i denote the i^{th} column of A, m-vector. Then we can write

$$\boldsymbol{b} = X = \sum_{i=1}^{n} x_i \boldsymbol{a}_i$$

The equation can be displayed schematically as follows:



COSC6364 Adv. Numerical Analysis

Wu. Panruo

- New interpretation that is essential for numerical linear algebra: if b = Ax, then b is the linear combination of the columns of A.
 - Let a_i denote the i^{th} column of A, m-vector. Then we can write

$$\boldsymbol{b} = X = \sum_{i=1}^{n} x_i \boldsymbol{a}_i$$

The equation can be displayed schematically as follows:

$$egin{aligned} egin{bmatrix} b \end{bmatrix} = egin{bmatrix} oldsymbol{a}_1 & oldsymbol{a}_2 & \dots & oldsymbol{a}_n \end{bmatrix} egin{bmatrix} x_1 \ x_2 \ \vdots \ x_n \end{bmatrix} \end{aligned}$$



COSC6364 Adv Numerical Analysis

- New interpretation that is essential for numerical linear algebra: if b = Ax, then b is the linear combination of the columns of A.
 - Let a_i denote the i^{th} column of A, m-vector. Then we can write

$$\boldsymbol{b} = X = \sum_{i=1}^{n} x_i \boldsymbol{a}_i$$

The equation can be displayed schematically as follows:

$$\begin{bmatrix} b \end{bmatrix} = \begin{bmatrix} a_1 & a_2 & \dots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$= x_1 \boldsymbol{a}_1 + x_2 \boldsymbol{a}_2 + \ldots + x_n \boldsymbol{a}_n$$



COSC6364 Adv. Numerical Analysis

Wu, Panruc



COSC6364 Adv. Numerical Analysis

Wii Panrii

■ Mathematician: Ax = b means matrix A (representing linear transformation) acts on vector x to produce b.



COSC6364 Adv. Numerical Analysis

Wii. Panrii

- Mathematician: Ax = b means matrix A (representing linear transformation) acts on vector x to produce b.
- Numerical analyst: Ax = b means vector x acts on A to produce b (as linear combination of columns of A).



OSC6364 Ac Numerical Analysis

Wu. Panru

- Mathematician: Ax = b means matrix A (representing linear transformation) acts on vector x to produce b.
- Numerical analyst: Ax = b means vector x acts on A to produce b (as linear combination of columns of A).
- Function can be seen as an infinite dimensional vector. Matrix A's columns can be function



COSC6364 Adv. Numerical Analysis

Wu, Panruo



COSC6364 Adv. Numerical Analysis

Nu. Panruo

■ For a matrix product B = AC, each column of B is a linear combination of the columns of A.



COSC6364 Adv. Numerical Analysis

Wii Panriio

- For a matrix product B = AC, each column of B is a linear combination of the columns of A.
- To see this, write:



COSC6364 Adv. Numerical Analysis

Wu, Panruo

- For a matrix product B = AC, each column of B is a linear combination of the columns of A.
- To see this, write:

$$AC = \underbrace{\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}}_{m \times n} \underbrace{\begin{bmatrix} c_{11} & c_{12} & \dots & c_{1k} \\ c_{21} & c_{22} & \dots & a_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ c_{n1} & c_{n2} & \dots & c_{nk} \end{bmatrix}}_{n \times k}$$

COSC6364 Adv. Numerical Analysis

Wu. Panrue

- For a matrix product B = AC, each column of B is a linear combination of the columns of A.
- To see this, write:

$$AC = \underbrace{\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}}_{m \times n} \underbrace{\begin{bmatrix} c_{11} & c_{12} & \dots & c_{1k} \\ c_{21} & c_{22} & \dots & a_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ c_{n1} & c_{n2} & \dots & c_{nk} \end{bmatrix}}_{n \times k}$$

$$B = \underbrace{\begin{bmatrix} \mathbf{b_1} \mid \mathbf{b_2} \mid \dots \mid \mathbf{b_k} \end{bmatrix}}_{m \times k}$$



COSC6364 Adv. Numerical Analysis

Vu, Panruo

Then column $\mathbf{b_i}$ is just

COSC6364 Adv. Numerical Analysis

Vu. Panruo

Then column $\mathbf{b_i}$ is just

$$\mathbf{b_i} = \begin{bmatrix} a_{11}c_{1i} + a_{12}c_{2i} + \dots + a_{1n}c_{ni} \\ a_{21}c_{1i} + a_{22}c_{2i} + \dots + a_{2n}c_{ni} \\ \vdots \\ a_{m1}c_{1i} + a_{m2}c_{2i} + \dots + a_{mn}c_{ni} \end{bmatrix}$$

COSC6364 Adv. Numerical Analysis

Vu, Panruc

Then column b_i is just

$$\mathbf{b_{i}} = \begin{bmatrix} a_{11}c_{1i} + a_{12}c_{2i} + \dots + a_{1n}c_{ni} \\ a_{21}c_{1i} + a_{22}c_{2i} + \dots + a_{2n}c_{ni} \\ \vdots \\ a_{m1}c_{1i} + a_{m2}c_{2i} + \dots + a_{mn}c_{ni} \end{bmatrix}$$

$$= c_{1i} \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} + c_{2i} \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} + \dots + c_{ni} \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

COSC6364 Adv Numerical Analysis

Then column b_i is just

$$\mathbf{b_{i}} = \begin{bmatrix} a_{11}c_{1i} + a_{12}c_{2i} + \dots + a_{1n}c_{ni} \\ a_{21}c_{1i} + a_{22}c_{2i} + \dots + a_{2n}c_{ni} \\ \vdots \\ a_{m1}c_{1i} + a_{m2}c_{2i} + \dots + a_{mn}c_{ni} \end{bmatrix}$$

$$= c_{1i} \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} + c_{2i} \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} + \dots + c_{ni} \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

$$= c_{1i}\mathbf{a}_{1} + c_{2i}\mathbf{a}_{2} + \dots + c_{ni}\mathbf{a}_{n}$$

$$=c_{1i}\mathbf{a_1}+c_{2i}\mathbf{a_2}+\ldots+c_{ni}\mathbf{a_n}$$



COSC6364 Adv. Numerical Analysis

Nu, Panruo



COSC6364 Adv. Numerical Analysis

Wu, Panruc

$$oldsymbol{u}oldsymbol{v}^* = oldsymbol{u}ig[\overline{v_1} \quad \overline{v_2} \quad \dots \quad \overline{v_n}ig]$$



COSC6364 Adv. Numerical Analysis

Wu, Panruo

$$egin{aligned} oldsymbol{u}oldsymbol{v}^* &= oldsymbol{u}ig[\overline{v_1}oldsymbol{v} & \overline{v_2}oldsymbol{u} & \ldots & \overline{v_n}ig] \ &= ig[\overline{v_1}oldsymbol{u} & \overline{v_2}oldsymbol{u} & \ldots & \overline{v_n}oldsymbol{u} \end{bmatrix}$$



COSC6364 Adv. Numerical Analysis

Wu, Panruc

$$uv^* = u \begin{bmatrix} \overline{v_1} & \overline{v_2} & \dots & \overline{v_n} \end{bmatrix}$$

$$= \begin{bmatrix} \overline{v_1}u & \overline{v_2}u & \dots & \overline{v_n}u \end{bmatrix}$$

$$= \begin{bmatrix} u_1\overline{v_1} & u_1\overline{v_2} & \dots & u_1\overline{v_n} \\ u_2\overline{v_1} & u_2\overline{v_2} & \dots & u_2\overline{v_n} \\ \vdots & \vdots & \ddots & \vdots \\ u_n\overline{v_1} & u_n\overline{v_2} & \dots & u_n\overline{v_n} \end{bmatrix}$$