

Ů Š Ā Ŵ Ă Ė Š Ž Ÿ Ź Ğ

داده ها، بازه ها / ...

$\chi^{\circ} \acute{H}_t$, $\underline{\psi}^{\circ} \grave{H}\check{Y}$ $\overset{\circ}{\text{?}}\grave{A}_t \grave{H}$ `` $\underline{\psi}^{\check{J}}$?A ?P , ?

' e ' e ' , m , v j c o . , ' ~ - ' ' ? ? ? ? - ? H Ž Đ Ā Ī P D G Ÿ D Ğ Ē Ž Ď Ā d Z Ů Ā L H ? ? ? ~ ~ ~ ~ ~ . k , ? D ? ? /
? E ' > ^ ,

"... / ... [?] [?] ...
[?] [?] [?] [?] [?] [?]

1. $\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$

ÄËĞŦZĚŠĞĚ

[illegible]
$$g\dot{H}^{\circ}\tilde{A}_{\square}''\boxtimes G'_{\alpha o}\boxtimes\tilde{A}_{\square}\underline{y}$$

ZŽŮGŮVĚLJ

$\zeta_j = \zeta_j^0 - \frac{1}{2} \frac{\partial \zeta_j^0}{\partial x^i} x^i + \frac{1}{6} \frac{\partial^2 \zeta_j^0}{\partial x^i \partial x^k} x^i x^k - \frac{1}{24} \frac{\partial^3 \zeta_j^0}{\partial x^i \partial x^k \partial x^l} x^i x^k x^l + \dots$

$gH^{\circ} \tilde{A}_\bullet'' \boxtimes G' \rightarrow G \boxtimes \tilde{A} \xrightarrow{\psi}$

s ŽďAdZ ůĎL zÐΘ žŷ

$\mathbb{C}^n$ 上 の 線形 変換 T が $T^2 = 0$ を 満たす とき T は $\mathbb{C}^n$ 上 の 零 変換 である ことを 示す .

$g\acute{H}^{\circ}\tilde{A}_{\square}''\boxtimes\circ'\tau G\boxtimes\tilde{A}\wr$

:ŝŌĖǺ|| Ǽ ĞǺĖŝŶŌ

[illegible]

$g\dot{H}^{\circ}\tilde{A}_{\square}''\boxtimes_{\mathcal{O}}'\mathcal{G}\tilde{A}_{\square}\mathcal{U}$