On the set function in $\mathbb{R}^n$

measure and integral

Nguyen Ngoc Khanh

October 2023

In $\mathbb{R}^n$, let u be a set function on $\mathbb{R}^n$ which is a non-negative function on a subcollection of the collection of all subsets of $\mathbb{R}^n$, i.e. $u : \mathcal{U} \to [0, +\infty]$, $\mathcal{U} \subseteq \mathcal{P}(\mathbb{R}^n)$.

We will construct the smallest set of assumptions on u such that the following limit converges.

$$Du(x) = \lim_{B \searrow x} \frac{u(B)}{|B|}$$

where $B \searrow x$ denotes the contraction of B into a singleton x, and $|B|$ denotes the Lebesgue measure of B.

Assumption 1 (Closed balls) $\overline{\mathcal{B}_r(x)} \in \mathcal{U}$ *for all* $x \in \mathbb{R}^n$ *and* $r > 0$

Consider a simpler version where B is a closed ball of radius $r > 0$ centered at x, i.e. $B = \overline{\mathcal{B}_r(x)}$. Define lim sup, lim inf as follows

- $\overline{D}u(x) = \limsup_{B \searrow x} \frac{u(B)}{|B|} = \lim_{r \to 0^+} \left(\sup_{t \in (0,r)} \frac{u(B_t(x))}{|B_t(x)|} \right)$
- $\underline{D}u(x) = \liminf_{B \searrow x} \frac{u(B)}{|B|} = \lim_{r \to 0^+} \left(\inf_{t \in (0,r)} \frac{u(B_t(x))}{|B_t(x)|} \right)$

The necessary and sufficient condition for the existence of $Du(x)$ is $\overline{D}u(x) = \underline{D}u(x)$. If $\overline{D}u(x) > \underline{D}u(x)$, let $\alpha, \beta \in \mathbb{Q}$ such that $\overline{D}u(x) > \alpha > \beta > \underline{D}u(x) \geq 0$. Then $Du(x)$ exists for almost every $x \in \mathbb{R}^n$ if $A_{\alpha\beta} = \{x \in \mathbb{R}^n : \overline{D}u(x) > \alpha > \beta > \underline{D}u(x)\}$ has measure zero for all $\alpha, \beta \in \mathbb{Q}$. If $x \in A_{\alpha\beta}$, then given any $\delta > 0$, there exist balls $B^{(\alpha)}, B^{(\beta)}$ centered at x with radius less than δ such that $\frac{u(B^{(\alpha)})}{|B^{(\alpha)}|} > \alpha > \beta > \frac{u(B^{(\beta)})}{|B^{(\beta)}|}$. That is, the collections $\{B^{(\alpha)}\}$ and $\{B^{(\beta)}\}$ each cover $A_{\alpha\beta}$ finely.

Lemma 1 (Vitali Covering Theorem for Lebesgue measure) *A collection $\mathcal{V}$ of closed balls is said to cover a set E finely if for every $x \in E$ and $\eta > 0$, there is a ball B containing x whose radius is less than η. Suppose E is covered finely by a collection $\mathcal{V}$ of balls such that $0 < \mu_e(E) < +\infty$ [1]. Then, given any $\epsilon > 0$, there is a countable collection M of disjoint balls such that*

$$\mu\left(E \setminus \bigcup_{B_k \in M} B_k\right) = 0 \text{ and } \sum_{B_k \in M} \mu(B_k) < \mu_e(E) + \epsilon$$

Claim 1 *Given any open ball U contains a subset $E \subseteq A_{\alpha\beta}$, the $u(\overline{U}) \geq \alpha |E|_e$*

Given any open ball U contains a subset $E \subseteq A_{\alpha\beta}$, since U is open, for each $x \in E$, we can find a collection of balls $\{B_U^{(\alpha)}\}$ that finely covers E and is contained in U. By Vitali Covering Theorem for Lebesgue measure on finite outer measure E with $\{B_U^{(\alpha)}\}$, given any $\eta > 0$, there exists a finite collection of pairwise disjoint balls $\{B_i\}_{i=1}^N$ contained in U such that $\left| E \setminus \bigcup_{i=1}^N B_i \right|_e < \eta/\alpha$ and $u(B_i) > \alpha |B_i|$

Assumption 2 (Additivity) $A, B \in \mathcal{U}$ *and* $A \cap B = \varnothing \implies A \cup B \in \mathcal{U}$ *and* $u(A \cup B) = u(A) + u(B)$

Assumption 3 (Subset) $A, B \in \mathcal{U}$ *and* $A \subseteq B \implies u(A) \leq u(B)$

[1] if $\mu_e(E) = 0$, the result is immediate

Then,

$$
\begin{aligned}
u(\overline{U}) &\geq u\left(\bigcup_{i=1}^{N} B_i\right) && \text{(subset)} \\
&= \sum_{i=1}^{N} u(B_i) && \text{(additivity)} \\
&> \alpha \sum_{i=1}^{N} |B_i| \\
&= \alpha \left|\bigcup_{i=1}^{N} B_i\right| && \text{(additivity of Lebesgue measure)} \\
&= \alpha \left|E \cap \bigcup_{i=1}^{N} B_i\right|_e && \text{(subset of Lebesgue measure)} \\
&= \alpha \left(|E|_e - \left|E \setminus \bigcup_{i=1}^{N} B_i\right|_e\right) && \text{(Caratheodory condition)} \\
&= \alpha |E|_e - \eta
\end{aligned}
$$

As the inequality is true for all $\eta > 0$, $u(U) \geq \alpha |E|_e$

Assumption 4 (Finite outer measure of $A_{\alpha\beta}$) $|A_{\alpha\beta}|_e < +\infty$ *for all* $\alpha, \beta \in \mathbb{Q}$

By Vitali Covering Theorem for Lebesgue measure on finite outer measure $A_{\alpha\beta}$ with $\{B^{(\beta)}\}$, given any $\epsilon > 0$, there exists a finite collection of pairwise disjoint balls $\{B_i\}_{i=1}^N$ such that $\left|A_{\alpha\beta} \setminus \bigcup_{i=1}^N B_i\right| < \epsilon$, $\sum_{i=1}^N |B_i| < |A_{\alpha\beta}|_e + \epsilon$, and $|B_i| > \frac{1}{\beta} u(B^{(\beta)})$
Then

$$
\begin{aligned}
|A_{\alpha\beta}| + \epsilon &> \sum_{i=1}^{N} |B_i| \\
&> \frac{1}{\beta} \sum_{i=1}^{N} \mu(B_i) \\
&\geq \frac{\alpha}{\beta} \sum_{i=1}^{N} |B_i^o \cap A_{\alpha\beta}|_e && \text{(by claim 1 as open ball } B_i^o \text{ contains } B_i^o \cap A_{\alpha\beta} \subseteq A_{\alpha\beta}\text{)} \\
&\geq \frac{\alpha}{\beta} \left|A_{\alpha\beta} \cap \bigcup_{i=1}^{N} B_i^o\right|_e && \text{(subadditivity of Lebesgue outer measure)} \\
&\geq \frac{\alpha}{\beta} \left|A_{\alpha\beta} \cap \bigcup_{i=1}^{N} B_i\right|_e && (|A \cap B|_e \leq |A \cap B^o|_e + |A \cap B^b|_e \ ^2) \\
&\geq \frac{\alpha}{\beta} \left(|A_{\alpha\beta}|_e - \left|A_{\alpha\beta} \setminus \bigcup_{i=1}^{N} B_i\right|_e\right) && \text{(Caratheodory condition)} \\
&> \frac{\alpha}{\beta} |A_{\alpha\beta}|_e - \frac{\alpha}{\beta}\epsilon
\end{aligned}
$$

Send ϵ to 0^+ implies $|A_{\alpha\beta}| \geq \frac{\alpha}{\beta}|A_{\alpha\beta}|$. Hence $|A_{\alpha\beta}| = 0$

Theorem 1 *Given a set function u on $\mathbb{R}^n$ that is defined for all closed balls $\overline{\mathcal{B}_r(x)}, r > 0$ that is equipped with (1) additivity and (2) subset, Then*

$$Du(x) = \lim_{B \searrow x} \frac{u(B)}{|B|}$$

is defined for almost every $x \in \mathbb{R}^n$ where $B \searrow x$ denotes the contraction of closed balls $\overline{\mathcal{B}_r(x)}, r > 0$ into a singleton x

Remark 1 *Some remarks*

- *If u is the Lebesgue measure, then $Du(x) = 1$ everywhere.*
- *If u is any positive measure on $\mathbb{R}^n$, $Du(x)$ is defined everywhere.*
- *If $f : \mathbb{R}^n \to [0, +\infty]$ is Lebesgue measurable, let $u(E) = \int_E f$, then we have the Lebesgue Differentiation Theorem*

Definition 1 (Absolute Continuous Set Function) $u \geq 0$, $\lim_{B \searrow x} u(B) = 0$ *a.e*

Lemma 2 *If u is an absolute continuous set function and $\lim_{B \searrow x} \frac{u(B)}{|B|} = 0$. Then, $u(E) = 0$ for all E*

Theorem 2 *u is an absolute continuous set function if and only if*

- $u' = \lim_{B \searrow x} \frac{u(B)}{|B|}$ *is defined a.e*
- $u' \in L$
- $u(E) = \int_E u'$